

The new Checklist of the Italian Fauna: Formicidae

ENRICO SCHIFANI¹ *

¹ Department of Chemistry, Life Sciences and Environmental Sustainability (SCVSA), University of Parma, Parco Area delle Scienze 11/A, I-43124 Parma (Italy)

* corresponding author, email: enrico.schifani@unipr.it

Keywords: Biogeography, insect diversity, myrmecology, species list, species distribution

SUMMARY

I present the updated version of the ‘Checklist of the Italian Fauna’ for what concerns ants (Insecta: Hymenoptera: Formicidae), which is part of the broader effort to produce an updated comprehensive checklist of the Italian fauna about 25 years after the first edition. The present list is the fourth Italian checklist of ants to be published since 1916, and refers to the state of art on November 2020. A simplified version of the data set is available as a supplementary file to this paper, while the full data set will be accessible in a regularly updated form from the LifeWatch Italy Data Portal (<https://dataportal.lifewatchitaly.eu/data>). Compared to the previous list by Poldi and others, published in 1995, the new one contains changes retrieved from 86 literature sources, including 17 published between 1921 and 1995 (which were missed in the previous checklist) and 69 published from 1996 to 2020. These references add 50 new species, including 9 new endemic species, as well as 68 nomenclatural changes and 88 distribution novelties at the regional level to the previous checklist. A total of 267 species and subspecies belonging to 7 ant subfamilies and 42 genera are part of the new list. The knowledge of the Italian ant fauna is rapidly improving on several fronts, and such dynamism is well-testified by the several novelties from 10 articles published after this dataset was compiled, to be included in the first future update of the on-line checklist. A further new list with extensive comments, detailed species distribution and biogeographic consideration will be desirable as soon as the situation stabilizes.

INTRODUCTION

About 25 years ago, a complete list of all the species of animals known to occur in the Italian territory was compiled thanks to the involvement of several specialists within the project ‘Checklist of the Italian Fauna’ (Minelli et al. 1993-1995), probably making

Italy the first country to have a list of this kind. Starting from 2020, an important effort is now being conducted in the attempt to update this list, making a new data set available in a new form that allows continuous updates to be made thanks to the LifeWatch Italy Data Portal (<https://dataportal.lifewatchitaly.eu/da>)

ta; see Bologna et al. 2022). The newly produced data sets of each systematic group will be accompanied by data papers published by the journal *Biogeographia*, explaining the work behind the checklists and commenting their results (Bologna et al. 2022).

Ants are a large family of aculeate hymenopterans comprising around 14,000 species (Bolton 2021). Their monophyly is well-demonstrated and their taxonomic treatment as a single family appears solid (Ward 2014, Borowiec et al. 2019). Thanks to their widespread success in terrestrial ecosystems of the globe, ants play important ecological roles, often interacting directly with human activities, and are considered a model for biodiversity monitoring (Hölldobler & Wilson 1990, Lach et al. 2010, Parker & Kronauer 2021).

During the last few decades, ant taxonomy has made important advancements at the global and European level (Ward 2007, Seifert 2018). For instance, it has largely improved its methodologies in order to uncover cryptic species (i.e. species only diagnosable by quantitative morphology or genetics, and not by qualitative morphology) and find stable and objective criteria for their delimitation through integrative approaches (e.g. Csósz et al. 2015, Seifert et al. 2017, Wagner et al. 2017, Steiner et al. 2018, Schifani et al. 2021a). Moreover, phylogenetics and phylogenomics have allowed to clarify evolutionary relationships between taxa at an unprecedented level (e.g. Ward et al. 2015, 2016). New online resources have greatly facilitated taxonomic and faunistic investigations (see AntCat, AntMaps and AntWebl, see Janicki et al. 2016, Guénard et al. 2017, Bolton et al. 2022). Nonetheless, the Mediterranean ant fauna, which is especially rich and diverse due to a highly complex paleogeographic history and generally favourable climatic conditions, still needs significant efforts to become well-described.

The first Italian checklist of ants, published in a monograph by Emery (1916), included 186 taxa (including ‘varieties’) illustrated through dichotomous keys. A second checklist was then published by Baroni Urba-

ni (1971), who listed 225 taxa and provided a meticulous compendium of all published literature and of all distribution data available at that time. Both Emery (1916) and Baroni Urbani (1971) based their checklist on a territory extending beyond the political borders of Italy, adducing biogeographical reasons. At last, a third checklist of 252 taxa was published by Poldi et al. (1995) within the ‘Checklist della Fauna d’Italia’ framework, this time considering only the Italian territory. As part of the project for a new ‘Checklist of the Italian Fauna’ (Bologna et al. 2022), I present in this paper a new checklist of the Italian ants, containing significant updates to the previous work by Poldi et al. (1995). The data presented in this paper refer to the state of art at November 2020, while data on the online version of the checklist (LifeWatch) will witness regular updates.

MATERIALS AND METHODS

Study area

Like the previous one (Poldi et al. 1995), the new checklist is based on the Italian territory with the only addition of the microstates of San Marino and Vatican City. The study area is divided into four regions to illustrate species distribution (Fig. 1): ‘N’ corresponds to continental Italy and is the sum of eight administrative regions of northern Italy (Aosta Valley, Emilia-Romagna, Friuli-Venezia Giulia, Liguria, Lombardy, Piedmont, Veneto, Trentino-Alto Adige); ‘S’ corresponds to peninsular Italy and contains the remaining ten peninsular regions from central and southern Italy (Abruzzo, Apulia, Calabria, Campania, Basilicata, Marche, Molise, Lazio, Tuscany, Umbria) plus San Marino and the Vatican City; ‘Sa’ corresponds to the region of Sardinia; ‘Si’ corresponds to the region of Sicily. It is worth noting that the administrative regions of Sardinia and Sicily include not only the two homonymous very large islands, but also all the small islands that surround them. This subdivision mostly overlaps with the biogeographic patterns of the Italian ants (Wang et al. 2022).




Figure 1. Study area: The Italian territory is divided into four sub-regions following Poldi et al. (1995) and the general framework of the project for a new Checklist of the Italian Fauna: N (Continental Italy), S (Peninsular Italy), Si (Sicily), Sa (Sardinia).

Literature data

The current version is based on data published until November 2020. Literature was searched through the search engines Google Scholar and Scopus, as well as through the AntMaps database on geographic distribution (www.antmaps.org; see Janicki et al. 2016, Guénard et al. 2017) and on the AntCat taxonomic catalogue (www.antcat.org; see Bolton 2021). Distribution records or nomenclatural changes were verified directly by examining and evaluating the information within each paper. Novel species records originally published by Radchenko et al. (2004) within the Fauna Europaea framework (and sometimes reported by following authors, e.g. Borowiec 2014) were not considered: they are clearly the result of an error, since they introduce several distribution novelties, some of which are rather incredible, in a checklist context in which distribution novelties were not expected in the first place (e.g. Schifani & Alicata 2018).




Figure 2. Worker specimens of three Italian endemic ant species described after the checklist published by Poldi et al. (1995); images from AntWeb (www.antweb.org): a) *Myrmecina melonii* Rigato, 1999 from Sardinia (CASENT0904563), b) *Temnothorax alienus* Schulz, Heinze & Pusch, 2007 from the Italian peninsula (ANTWEB1041296), c) *Aphaenogaster trinacriæ* Alicata & Schifani, 2019 from Sicily (ANTWEB1041230). Photos by Will Ericson, Roland Schultz and Enrico Schifani. Scale: 1 mm.

Data set organization

The data set is available on-line version from the LifeWatch website. A simplified version is given in the Supplementary file S1.

The ant checklist contains 22 of the 46 columns that are common to all animal groups. The first columns are dedicated to the systematic classification of each species, from phylum rank to family, genus, species and

subspecies, including the descriptors of the former three. They are followed by an additional column presenting the name of the species/subspecies in the Fauna Europaea database (www.faunaeur.com) whenever possible, and two columns describing its status as either endemic (e), subendemic (s), introduced (a) or possibly introduced (?). Finally, two columns contain references for the nomenclatural or distribution changes compared to the checklist by Poldi et al. (1995).

The Supplementary file S2 contains a list of taxa that were present in the previous edition (Poldi et al. 1995) but were excluded from the update checklist for one of the following three reasons according to the relevant literature: *i*) they were considered dubious names; *ii*) they were considered absent from Italy; *iii*) after careful examination, their inclusion by Poldi et al. (1995) was found to be unsupported by any published paper.

Data set information

Object name: Checklist of the Italian Fauna: Formicidae

Characters encoding: Unicode (UTF-8)

Data set citation:

Schifani E., 2021. Formicidae. In: Bologna M.A., Zapparoli M., Oliverio M., Minelli A., Bonato L., Cianferoni F., Stoch F. (eds.), Checklist of the Italian fauna. Version 1.0. Last update: 2021-05-31.

Format name: xml, Extensible Markup Language, for the online version at LifeWatch Italy.

Format version: 1.0

Distribution: <https://dataportal.lifewatchitaly.eu/view/urn%3Auuid%3Ac1f2ab37-61e4-48e9-b3a9-15bdbf002f9d>

Date of creation: November 2nd, 2020

Date of last revision: November 28th, 2021

Date of publication: July 23rd, 2021

Language: English

License of use: Creative Commons Attribution 4.0 International License

Metadata language: English

Metadata managers: Marco Bologna, Lucio Bonato, Fabio Cianferoni, Alessandro Minelli, Marco Oliverio, Fabio Stoch, Marzio Zapparoli & LifeWatch Italy

Management details

Project title: The new Checklist of the Italian Fauna: Formicidae

Database manager: Enrico Schifani; Marco Bologna, Lucio Bonato, Fabio Cianferoni, Alessandro Minelli, Marco Oliverio, Fabio Stoch, Marzio Zapparoli & LifeWatch

Temporal coverage: Anything published until November 2020.

Record basis: Published records in the scientific and grey literature.

Funding grants: funding for the general project of the checklist of the Italian fauna was obtained from LifeWatch Italy; no funding was specifically available for the update of Formicidae.

Geographic information

General description: The dataset includes records from the national territory of Italy, including the two major islands Sardinia and Sicily, together with archipelagos and minor islands politically under the Italian legislation.

Geographic units: The geographical units within the Italian national territory for terrestrial records refer to the administrative boundaries of the Italian regions, in addition to San Marino Republic and Vatican City. Continental Italy was also divided in only two units, namely North (Friuli - Venezia Giulia, Veneto, Trentino - Alto Adige, Lombardia, Valle d'Aosta, Piemonte, Liguria, Emilia Romagna) and South (Toscana, Marche, Umbria, Lazio, Abruzzo, Molise, Campania, Puglia, Basilicata, Calabria).

Bounding box: All areas falling under Italian administration (in addition to San Marino and Vatican City) from 35° 25' to 47° 06' N and from 6° 35' to 19° 20' E (WGS84 reference system) were included.

Sampling design: We did not perform any additional sampling to collect records, but we used only published data.

Habitat type: NA.

Biogeographic region: Palearctic.

Countries: Italy, San Marino, Vatican City.

Quality control for geographic data: We checked that the published localities matched the geographical units used for the checklist at the level of administrative regions.

Literature records

General description: Only published records are included.

Literature search methods: Google Scholar, Scopus.

Literature list: Müller (1921), Finzi (1924), Menozzi (1925), Bernard (1967), Baroni Urbani (1971, 1977), Kutter (1977), Süss (1979), Agosti & Collingwood (1987), Ortiz & Tinaut (1987), Buschinger et al. (1988), Casevitz-Weulersse (1990), Seifert (1992a, b, 1994), Bolton (1995), Mei (1995), Cagniant (1996), Carniel (1998), Buschinger et al. (1999), Rigato (1999, 2011), Sanetra et al. (1999), Seifert (2000, 2002, 2005, 2012a, b, 2018, 2020), Bolton (2003), Csősz & Seifert (2003), Glaser (2003, 2004), Limonta & Colombo (2003), Radchenko & Elmes (2003, 2010), Bonelli et al. (2004), Bernasconi et al. (2006), Radchenko et al. (2006), Scupola (2006, 2017), Csősz et al. (2007, 2014, 2015), Baroni Urbani & De Andrade (2007), Schultz et al. (2007), Verdinelli et al. (2007), Wetterer et al. (2007), Boer (2008, 2013), Elmes et al. (2008), Jucker et al. (2008), Seifert et al. (2009, 2014, 2017), Glaser et al. (2011, 2012), Steiner et al. (2010, 2018), Bolton & Fisher (2011), Rigato & Toni (2011), Yoshimura & Fisher (2012), Loi (2013), Brácko et al. (2014), Borowiec et al. (2015), Salata & Borowiec (2015, 2018a), Sarnat et al. (2015), Seifert & Csősz (2015), Ward et al. (2015), Brácko (2017), Centorame et al. (2017), Galkowski & Cagniant (2017), Schifani (2017), Wagner et al. (2017), Salata et al. (2018, 2021), Schifani & Alicata (2018, 2019), Ali-

cata & Schifani (2019), Galkowski et al. (2019), Giannetti et al. (2019), Masoni et al. (2019), Schär et al. (2020). Quality control for literature data: Additional references were searched through the literature, scanning the cited references of each paper.

Taxonomic information

General description: Only records reporting species or subspecies were included, disregarding records identified only at higher levels like genus, family, etc.

Taxonomic coverage: family Formicidae

Taxonomic rank: Only species and subspecies are reported; the dataset reports higher taxa for each species, including Phylum, Class, Order, Family, Genus.

Taxon specialists: Enrico Schifani.

Nomenclature: The nomenclature followed the most recently updated taxonomic framework.

Taxonomic remarks: All taxonomic changes that occurred since the publication of the previous checklists (1995) until November 2020. Species that were included in the previous checklist and are now considered not valid or not present in Italian waters, are excluded from the dataset.

Quality control for taxonomic data: Taxonomic data were checked and updated to include revision of names, synonyms, delimitation of genera and higher taxa.

RESULTS

Summary statistics

The checklist contains 267 species and subspecies belonging to 42 genera and 7 subfamilies: Amblyoponinae (2), Dolichoderinae (13), Formicinae (80), Leptanillinae (3), Myrmicinae (156), Ponerinae (8), and Proceratiinae (4). The distribution of these taxa in the four sub-regions is the following: N: 191; S: 160; Sa: 74; Si: 132.

A total of 33 taxa is currently considered endemic to the Italian territory, while 12 are considered subendemic (Figure 2). Endemic taxa are concentrated in the S (58%) and Si (45%), while only a few occur in N (12%) and Sa (9%). Most of the endemic taxa belong to the subfamily Myrmicinae (78%). On the other hand, there are 13 alien species and the status of 2 species as either native or introduced is dubious. Most introduced species are known to inhabit N (70%) or Si (60%), followed by S (40%) and Sa (20%).

The global IUCN red list includes 24 species occurring in Italy: 5 *Formica* species from the *rufa* group are considered Near Threatened (NT), and 19 inquiline social parasites are considered Vulnerable (VU) (they belong to the genera *Camponotus*, *Formicoxenus*, *Harpagoxenus*, *Lasius*, *Leptothorax*, *Myrmica*, *Plagiolepis*, *Strongylognathus*, *Temnothorax*, and *Tetramorium*) (Social Insect Specialists group 1996).

Changes in comparison to the previous checklist by Poldi et al. (1995) resulted from information published in 86 literature sources (articles or books). Some of them were actually published earlier than the last checklist, and were either missed by Poldi et al. (1995) or published at a stage when their work was already in the process of being published. They are 17 references in total: Müller (1921), Finzi (1924), Menozzi (1925), Bernard (1967), Baroni Urbani (1971, 1977), Kutter (1977), Süss (1979), Agosti & Collingwood (1987), Ortiz & Tinaut (1987), Buschinger et al. (1988), Casevitz-Weulersse (1990), Seifert (1992a-b, 1994), Bolton (1995), Mei (1995). In addition, novelties came from the following 69 references (summarized in Table 1) that were published after 1995: Cagniant (1996), Carniel (1998), Buschinger et al. (1999), Rigato (1999, 2011), Sanetra et al. (1999), Seifert (2000, 2002, 2005, 2012a, b, 2018, 2020), Bolton (2003), Csósz & Seifert (2003), Glaser (2003, 2004), Limonta & Colombo (2003), Radchenko & Elmes (2003, 2010), Bonelli et al. (2004), Bernasconi et al. (2006), Radchenko et al. (2006), Scupola (2006, 2017), Csósz et al. (2007, 2014, 2015), Baroni Urbani & De Andrade (2007), Schultz

et al. (2007), Verdinelli et al. (2007), Wetterer et al. (2007), Boer (2008, 2013), Elmes et al. (2008), Jucker et al. (2008), Seifert et al. (2009, 2014, 2017), Glaser et al. (2011, 2012), Steiner et al. (2010, 2018), Bolton & Fisher (2011), Rigato & Toni (2011), Yoshimura & Fisher (2012), Loi (2013), Bračko et al. (2014), Borowiec et al. (2015), Salata & Borowiec (2015, 2018a), Sarnat et al. (2015), Seifert & Csósz (2015), Ward et al. (2015), Bračko (2017), Centorame et al. (2017), Galkowski & Cagniant (2017), Schifani (2017), Wagner et al. (2017), Salata et al. (2018, 2021), Schifani & Alicata (2018, 2019), Alicata & Schifani (2019), Galkowski et al. (2019), Giannetti et al. (2019), Masoni et al. (2019), Schär et al. (2020).

Recent changes to be included in future updates

A total of 10 papers were published after this dataset was uploaded on the LifeWatch platform (2021-2022), and will be included the first update of the checklist in the near future.

They imply the addition of 8 species and 1 genus new to Italy, including 4 species of new description and 3 new endemics, the removal of 1 species, 6 nomenclatural changes and 47 distribution novelties. Boudinot et al. (2021) demonstrated the inconsistency of the subgeneric classification of *Lasius* and synonymized five subgenera including Italian species. *Aphaenogaster strioloides* Forel, 1890 was recorded in Italy for the first time and discovered to replace *A. crocea* André, 1881 in Si, implying the absence of *A. crocea* in Italy (Schifani et al. 2021a). The presence of further 9 species in Si was confirmed or recorded for the first time while other 19 were considered absent (Schifani et al. 2021b). *Solenopsis lusitanica* Emery, 1915 was recorded in Italy for the first time, in both Sa and S regions, while other 4 species were added to Sa and 2 were removed (Schifani et al. 2021c,d). *Aphaenogaster ovaticeps* (Emery, 1898), previously considered an Italian endemic, was synonymized with *A. muelleriana* Wolf, 1915 (Salata et al. 2021). Another two new distribution records were published for S and Si (Schifani et al. 2022a). The alien *Brachypon-*

era chinensis (Emery, 1895) was added to the Italian fauna based on a discovery in S (Menchetti et al. 2022). The newly described *Colobopsis imitans* Schifani et al. 2021 was discovered to replace *C. truncata* (Spinola, 1808) in Si (Schifani et al. 2022b). *Acropyga palearctica* Menozzi, 1936 was recorded in

Italy for the first time, in the S region (Scupola et al. 2022). Three new *Temnothorax* species endemic or subendemic to Si were described, while *T. lagrecai* (Baroni Urbani, 1964) was discovered to be subendemic and not endemic due to its presence the Maltese islands (Schifani et al. 2022c).

Table 1. Temporal distribution of novelties produced since the checklist by Poldi et al. (1995) was published. The ‘new species described’ column refers to descriptions of entirely new taxa, while redescriptions and resurrections also had an important role. The ‘nomenclatural changes’ are intended as the number of new synonymies, changes in taxonomic rank or taxonomic recombinations occurred. The ‘distribution novelties’ column only refers to novelties changing species distribution in relation to the subdivision of Italy in four sub-regions used in this and in the previous checklist.

Years	New papers	New species added	New endemic species described	Nomenclatural changes	All distribution novelties
1994-2000	9	5	3	33	11
2001-2005	8	5	0	9	5
2006-2010	16	12	4	9	17
2011-2015	17	7	1	17	20
2016-2020	19	17	1	0	35
Total	69	50	9	68	88

DISCUSSION

Our understanding of the Italian ant fauna has made very important steps forward since the first list was published more than a century ago (Emery 1916), and changes were also significant during the last 25 years (Poldi et al. 1995). Recently, even a first regional handbook was published (Scupola 2018). At present, Italy appears to be the third most species-rich country in Europe, right after Greece and Spain (Salata & Borowiec 2018b, Tinaut & Ruano 2021), and significantly distances all the other countries from the continent (Csósz et al. 2021). In the Mediterranean region, it falls to the fourth place, as it is preceded by Turkey as well (Kiran & Karaman 2020).

Despite significant improvements during the last decades, there are still many open questions and unsolved issues. For instance, the actual figure of endemic taxa has to be revised as

there are still many taxa whose current valid status is unsupported by any evidence, with at least a few which are most-likely synonyms of species with a broader distribution. This has already happened for subspecific taxa such as *Tetramorium semilaeve siciliense* Santschi, 1927 and *T. s. liparaeum* Santschi, 1927 (see Sanetra et al. 1999). More in general, all subspecific taxa reported in the list need to be taxonomically revised. Furthermore, knowledge over the geographic distribution of several species is extensively (sometimes almost exclusively) based on records that were produced before their taxonomic boundaries were significantly reworked by modern revisions. Obvious examples are those of *Lasius alienus* (Foerster, 1850), *Solenopsis fugax* (Latreille, 1798), or *Tapinoma erraticum* (Latreille, 1798): these species were considered extremely widespread in Europe and Italy due to confusion with a variety of morphologically similar taxa, and their presence in most

or even all the Italian territory is currently uncertain (e.g. Seifert 2012, 2020, Seifert et al. 2017, Schifani et al. 2021b,c). On the other hand, distribution data on those species with cryptic habits, either inquiline social parasites (some Formicinae and Myrmicinae) or endogean taxa (mostly Amblyoponinae, Leptanillinae, Proceratiinae and some Ponerinae), are often scarce. In particular, the ancestral genus *Leptanilla* suffers from a combination of both insufficient taxonomic understanding and very limited distribution knowledge (Scupola & Ballarin 2009, Pérez-González et al. 2020).

No conservation priorities have yet been defined specifically for the Italian ant fauna, while the conservation status of ants in Europe is also rarely well assessed. Among the ‘Vulnerable’ social parasite species of Italy, some appear to be truly localized (Schifani et al. 2021e), while others are probably widespread (Schifani et al. 2021b). All are certainly difficult to sample (Espadaler & López-Soria 1991, Schifani 2017), making it difficult to have a clear understanding of their status. In the IUCN list, the status of these species is usually accompanied by a “needs updates” note (Social Insect Specialists group 1996). The conservation status of narrowly-distributed Italian endemics is rarely documented (Schifani et al. 2020, 2022c). On the other hand, new exotic species becoming established are increasing across the Mediterranean, especially in urban areas (Schifani 2019, Menchetti et al. 2022). The establishment of part of the species most recently recorded from Italy and provisionally included in the list is still uncertain and requires further investigation (see *Camponotus atriceps* (Smith, 1858), *Technomyrmex pallipes* (Smith, 1876), and *Tetramorium lucayanum* Wheeler, 1905 reported by Jucker et al. 2008).

A significant number of nomenclatural changes, distribution novelties and species descriptions, were already published during 2021–2022, outdated the version of the checklist currently available on LifeWatch (and in the supplementary file of this paper) and testifying a very dynamic state of art. A significantly more accurate and stable picture will hopefully be achieved in the near future, allowing to better-

define biogeographic patterns and conservation priorities.

ACKNOWLEDGEMENTS

I am thankful for the invitation I received to curate the Formicidae checklist within the new Checklist of the Italian Fauna. For this opportunity, I wish to thank the Scientific Committee of the project for this opportunity, as well as Fabrizio Rigato, co-author of the previous 1995 checklist edition.

REFERENCES

- Agosti, D. & Collingwood, C.A. (1987) A provisional list of the Balkan ants (Hym. Formicidae) with a key to the worker caste. II. Key to the worker caste, including the European species without the Iberian. Mitteilungen der Schweizerischen Entomologischen Gesellschaft, 60, 261-293.
- Alicata, A. & Schifani, E. (2019) Three endemic *Aphaenogaster* from the Siculo-Maltese archipelago and the Italian Peninsula: part of a hitherto unrecognized species group from the Maghreb? (Hymenoptera: Formicidae: Myrmicinae). Acta Entomologica Musei Nationalis Prae-gae, 59, 1-16. DOI: 10.2478/aemnp-2019-0001
- Baroni Urbani, C. (1971) Catalogo delle specie di Formicidae d'Italia (Studi sulla mirmecofauna d'Italia X). Memorie della Società Entomologica Italiana, 50, 5-287.
- Baroni Urbani, C. (1977) Materiali per una revisione della sottofamiglia Leptanillinae Emery (Hymenoptera: Formicidae). Entomologica Basiliensis, 2, 427-488.
- Baroni Urbani, C. & de Andrade, M.L. (2003) The ant genus *Proceratium* in the extant and fossil record (Hymenoptera: Formicidae). Museo Regionale di Scienze Naturali, Torino, 36, 1-492.
- Bernard, F. (1967) Faune de l'Europe et du Bassin Méditerranéen. 3. Les fourmis (Hymenoptera Formicidae) d'Europe occidentale et septentrionale. Paris, Masson, 411 pp.

- Bernasconi, C., Maeder, A., Freitag A., Cherix, D. (2006) *Formica paralugubris* (Hymenoptera, Formicidae) in the Italian Alps from new data and old data revisited. *Myrmecological News*, 8, 251-256.
- Boer, P. (2008) *Plagiolepis obscuriscapa* Santschi, 1923, a junior synonym of *Plagiolepis pygmaea* (Latrelle, 1798) (Hymenoptera: Formicidae: Formicinae) and the use of pigmentation as discriminating character in ant taxonomy. *Zoologische Mededelingen*, 82, 485-488.
- Boer, P. (2013) Revision of the European ants of the *Aphaenogaster testaceopilosa*-group (Hymenoptera: Formicidae). *Tijdschrift voor Entomologie*, 156, 57-93.
- Bologna, M.A., Bonato, L., Cianferoni, F., Minelli, A., Oliverio, M., Stoch, F. & Zapparoli, M. (2022) Towards the new Checklist of the Italian fauna. *Biogeographia – The Journal of Integrative Biogeography*, 37, ucl001. DOI: 10.21426/B637156271
- Bolton, B. (1995b) A new general catalogue of the ants of the world. Cambridge, Mass.: Harvard University Press, 504 pp.
- Bolton, B. (2003) Synopsis and Classification of Formicidae. Memoirs of the American Entomological Society, 71, 370 pp.
- Bolton, B. & Fisher, B.L. (2011) Taxonomy of Afrotropical and West Palaearctic ants of the ponerine genus *Hypoponera* Santschi (Hymenoptera: Formicidae). *Zootaxa*, 2843, 1-118. DOI: 10.11646/zootaxa.2843.1.1
- Bolton, B. (2021) An online catalog of the ants of the world. Available from <https://antcat.org>. (accessed [15.12.2021])
- Bonelli, S., Barbero, F., Chiantelassa, M. & Balletto, E. (2004) Prima segnalazione di *Myrmica tulinae* Elmes et al. 2002 (Hymenoptera: Formicidae) in Italia. In: 65° Congresso Nazionale dell'Unione Zoologica Italiana, pp. 97-97.
- Borowiec, L. (2014) Catalogue of ants of Europe, the Mediterranean Basin and adjacent regions (Hymenoptera: Formicidae). *Genus*, 25, 1-340.
- Borowiec L., Galkowski, C. & Salata, S. (2015) What is *Tetramorium semilaeve* André, 1883? (Hymenoptera, Formicidae). *ZooKeys*, 512, 39-62. DOI: 10.3897/zookeys.512.10006
- Borowiec, M.L., Rabeling, C., Brady, S.G., Fisher, B.L., Schultz, T.R. & Ward, P.S. (2019) Compositional heterogeneity and outgroup choice influence the internal phylogeny of the ants. *Molecular Phylogenetics and Evolution*, 134, 111-121. DOI: 10.1016/j.ympev.2019.01.024
- Boudinot, B.E., Borowiec, M.L. & Prebus, M.M. (2021) Phylogeny, evolution, and classification of the ant genus *Lasius*, the tribe Lasiini, and the subfamily Formicinae (Hymenoptera: Formicidae). *Systematic Entomology*, 47, 113-151. DOI: 10.1111/syen.1252210
- Bračko, G., Wagner, H.C., Schulz, A., Gioahin, E., Matičič, J. & Tratnik, A. (2014) New investigation and a revised checklist of the ants (Hymenoptera: Formicidae) of the Republic of Macedonia. *North-Western Journal of Zoology*, 10, 13207.
- Bračko, G. (2017) First discoveries of colonies of the rare ant species *Camponotus tergestinus* Müller, 1921 (Hymenoptera: Formicidae) in situ. *Natura Sloveniae*, 19, 5-14.
- Buschinger, A., Ehrhardt, W., Fischer, K. & Ofer, J. (1988) The slave-making ant genus *Chalepoxenus* (Hymenoptera, Formicidae). I. Review of literature, range, slave species. *Zoologische Jahrbücher*, 115, 383-401.
- Buschinger A. (1999) Bemerkenswerte Ameisenfunde aus Südtirol (Hymenoptera: Formicidae). *Myrmecologische Nachrichten*, 3, 1-8.
- Cagniant, H. (1996) Les *Camponotus* du Maroc (Hymenoptera: Formicidae): clé et catalogue des espèces. *Annales de la Société entomologique de France* (N.S.), 32, 87-100.
- Carniel, A. (1998) Ricerche sulla mirmecofauna delle Prealpi Orobiche (Lombardia) (Insecta, Hymenoptera, Formicidae). *Atti del Museo Civico di Storia Naturale di Morbegno*, 9, 29-39.
- Casevitz-Weulersse, J. (1990) Étude systématique de la myrmécofaune corse (Hymenoptera, Formicidae) (Première partie). *Bulletin du Muséum national d'histoire naturelle, Section A, Zoologie, biologie et écologie animales*, 12, 135-163.

Centorame, M., Lancia, A., Mori, E., D'Eustacchio, D. & Fanfani, A. (2017) Could *Linepithema humile* (Hymenoptera Formicidae) influence ant community composition? A preliminary study in a natural area in Italy. *Redia*, 100, 89-94. DOI: 10.19263/REDIA-100.17.11

Csósz, S. & Seifert, B. (2003) *Ponera testacea* Emery, 1895 stat. n. - A sister species of *P. coarctata* (Latreille, 1802) (Hymenoptera: Formicidae). *Acta Zoologica Academiae Scientiarum Hungaricae*, 49, 201-214.

Csósz, S., Radchenko, A. & Schulz, A. (2007) Taxonomic revision of the Palaearctic *Tetramorium chefketi* species complex (Hymenoptera: Formicidae). *Zootaxa*, 1405, 1-38. DOI: 10.11646/zootaxa.1405.1.1

Csósz, S., Wagner, H.C., Bozsó, M., Seifert, B., Arthofer, W., Schlick-Steiner, B.C., Steiner, F.M. & Pénzes, Z. (2014) *Tetramorium indocile* Santschi, 1927 stat. rev. is the proposed scientific name for *Tetramorium* sp. C sensu Schlick-Steiner et al. (2006) based on combined molecular and morphological evidence (Hymenoptera: Formicidae). *Zoologischer Anzeiger*, 253, 469-481. DOI: 10.1016/j.jcz.2014.06.002

Csósz, S., Heinze, J. & Mikó, I. (2015) Taxonomic synopsis of the Ponto-Mediterranean ants of *Temnothorax nylanderi* species-group. *PLoS One*, 10, e0140000. DOI: 10.1371/journal.pone.0140000

Csósz, S., Báthori, F., Gallé, L., Lörinczi, G., Maák, I., Tartally, A., Kovács, E., Somogyi, A.A. & Markó, B. (2021) The Myrmecofauna (Hymenoptera: Formicidae) of Hungary: Survey of Ant Species with an Annotated Synonymic Inventory. *Insects*, 12, 78. DOI: 10.3390/insects12010078

Elmes, G. W., Radchenko, A., Balletto, E. & Barbero, F. (2008) *Myrmica obscura* Finzi—a good ant species (Hymenoptera: Formicidae) endemic to Italy. *Fragmenta Faunistica*, 51, 91-100.

Emery, C. (1916) Fauna entomologica italiana. I. Hymenoptera.-Formicidae. *Bollettino della Società Entomologica Italiana*, 47, 79-275.

Espadaler, X. & López-Soria, L. (1991) Rareness of certain Mediterranean ant species: fact or arti-

fact?. *Insectes Sociaux*, 38, 365-377. DOI: 10.1007/BF01241872

Finzi, B. (1924) Secondo contributo alla conoscenza della fauna mirmecologica della Venezia Giulia. *Bollettino della Società Entomologica Italiana*, 56, 120-123.

Galkowski, C. & Cagniant, H. (2017) Contribution à la connaissance des fourmis du groupe angustulus dans le genre *Temnothorax* (Hymenoptera, Formicidae). *Revue de l'Association Roussillonaise d'Entomologie*, 26, 180-191.

Galkowski, C., Aubert, C. & Blatrix, R. (2019) *Aphaenogaster ichnusa* Santschi, 1925, bona species, and redescription of *Aphaenogaster subterranea* (Latreille, 1798) (Hymenoptera, Formicidae). *Sociobiology*, 66, 420-425. DOI: 10.13102/sociobiology.v66i3.3660

Giannetti, D., Castracani, C., Spotti, F. A., Mori, A. & Grasso, D. A. (2019) Gall-Colonizing Ants and Their Role as Plant Defenders: From 'Bad Job' to 'Useful Service'. *Insects*, 10, 392. DOI: 10.3390/insects10110392

Glaser, F. (2003) Die Ameisenfauna (Hymenoptera, Formicidae) des Vinschgaus (Südtirol, Italien)—eine vorläufige Artenliste. *Gredleriana*, 3, 209-230.

Glaser, F. (2004) Verbreitung und Gefährdung von Ameisen (Hymenoptera, Formicidae) in Auen- und Uferlebensräumen der Etsch (Südtirol, Italien). *Gredleriana*, 4, 203-246.

Glaser, F., Michael, J. & Seifert, B. (2011) Rediscovered after 140 years at two localities: *Myrmica myrmicoxena* Forel, 1895 (Hymenoptera: Formicidae). *Myrmecological News*, 14, 107-111.

Glaser, F., Freitag, A. & Martz, H. (2012) Ants (Hymenoptera: Formicidae) in the Münstertal (Val Müstair)—a hot spot of regional species richness between Italy and Switzerland. *Gredleriana*, 12, 273-284.

Guénard, B., Weiser, M., Gomez, K., Narula, N., Economo, E.P. (2017) The Global Ant Biodiversity Informatics (GABI) database: a synthesis of ant species geographic distributions. *Myrmecological News*, 24, 83-89.

- Hölldobler, B., & Wilson, E. O. (1990). The ants. Harvard University Press, USA.
- Janicki, J., Narula, N., Ziegler, M., Guénard, B. & Economo, E.P. (2016) Visualizing and interacting with large-volume biodiversity data using client-server web-mapping applications: The design and implementation of antmaps.org. Ecological Informatics, 32, 185-193. DOI: 10.1016/j.ecoinf.2016.02.006
- Jucker, C., Rigato, F. & Regalin, R. (2008) Exotic ant records from Italy (Hymenoptera, Formicidae). Bollettino di Zoologia agraria e di Bachicoltura, 40, 99-107.
- Kiran, K. & Karaman, C. (2020) Additions to the ant fauna of Turkey (Hymenoptera, Formicidae). Zoosystema, 42, 285-329. DOI: 10.5252/zosystema2020v42a18
- Kutter, H. (1977) Hymenoptera, Formicidae. Insecta Helvetica. Fauna, 6, 1-298.
- Lach, L., Parr, C. & Abbott, K. (eds.) (2010) Ant ecology. Oxford University Press, UK. DOI: 10.1093/acprof:oso/9780199544639.001.0001
- Limonta, L. & Colombo, M. (2003) Record of *Pheidole megacephala* (F.), *Pheidole nodus* Smith and *Tetramorium bicarinatum* Nylander (Hymenoptera Formicidae), tropical species, in nursey imported plants. Bollettino di Zoologia Agraria e di Bachicoltura, 35, 287-289.
- Loi, A. (2013) Osservazioni sulla mirmecofauna e l'afidofauna della quercia da sughero in Sardegna. Doctoral Thesis at the University of Sassari, Italy.
- Masoni, A., Frizzi, F., Natali, C., Bernasconi, C., Ciofi, C. & Santini, G. (2019) Molecular identification of imported red wood ant populations in the Campigna biogenetic nature Reserve (Foreste Casentinesi National Park, Italy). Conservation Genetics Resources, 11, 231-236. DOI: 10.1007/s12686-018-0982-7
- Mei, M. (1995) Arthropoda di Lampedusa, Linosa e Pantelleria (Canale di Sicilia, Mar Mediterraneo). Hymenoptera Formicidae (con diagnosi di due nuove specie). Il Naturalista Siciliano, 19, 753-772.
- Menchetti, M., Schifani, E., Gentile, V. & Vila, R. (2022). The worrying arrival of the invasive Asian needle ant *Brachyponera chinensis* in Europe (Hymenoptera: Formicidae). Zootaxa, 5115, 146-150. DOI: 10.11646/zootaxa.5115.1.10
- Menozzi, C. (1925) Res mutinenses. Formicidae (Hymenoptera). Atti della Società Naturalisti e Matematici di Modena, 55, 22-47.
- Minelli, A., Ruffo, S. & La Posta, S. (eds.) (1993-1995) Checklist delle specie della Fauna d'Italia. Ministero dell'Ambiente, Comitato Scientifico per la Fauna d'Italia. Calderini Editore.
- Müller, G. (1921) Due nuove formiche della regione Adriatica. Bollettino della Società adriatica di scienze naturali in Trieste, 27, 46-49.
- Ortiz, F.J. & Tinaut, A. (1987) Citas nuevas o interesantes de formícidos (Hym. Formicidae) para Andalucía. Boletín de la Asociación española de Entomología, 11, 31-34.
- Parker, J. & Kronauer, D.J. (2021) How ants shape biodiversity. Current Biology, 31, R1208-R1214. DOI: 10.1016/j.cub.2021.08.015
- Pérez-González, S., Gómez-Durán, J.M. & Martínez-Ibáñez, M.D. (2020) Highlighting the Elusive: New Findings and a Redescription of the Rare Ant *Leptanilla plutonia* López, Martínez Et Barandica, 1994, Presenting Morphological Novelties for the Genus. Annales Zoologici, 70, 289-304. DOI: 10.3161/00034541ANZ2020.70.2.009
- Poldi, B., Mei, M. & Rigato, F. (1995) Hymenoptera Formicidae. In: Minelli, A., Ruffo, S., La Posta, S. (eds). Checklist delle Specie della Fauna Italiana, Calderini, Bologna, 1-10.
- Radchenko, A. & Elmes, G.W. (2003) A taxonomic revision of the socially parasitic *Myrmica* ants (Hymenoptera: Formicidae) of the Palaearctic region. Annales Zoologici, 53, 217-243.
- Radchenko, A. (2004) Formicidae. - Fauna Europaea version 1.1: www.faunaeur.org. (accessed [15.12.2021]).
- Radchenko, A., Elmes, G. W. & Alicata, A. (2006) Taxonomic revision of the *schencki*-group of the ant genus *Myrmica* Latreille (Hymenoptera:

- Formicidae) from the Palaearctic region. *Annales Zoologici*, 56, 499-538.
- Radchenko, A. & Elmes, G.W. (2010) *Myrmica* ants (Hymenoptera: Formicidae) of the old world. Warszawa: Natura optima dux Foundation, Poland.
- Rigato, F. (1999) *Myrmecina melonii* n. sp., a new ant from Sardinia, with a review of the West Palaearctic *Myrmecina* (Hymenoptera Formicidae). *Bollettino della Società Entomologica Italiana*, 131, 83-92.
- Rigato, F. (2011) Contributions to the taxonomy of West European and North African *Stenamma* of the *westwoodii* species-group. (Hymenoptera Formicidae). *Memorie della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano*, 37, 1-56.
- Rigato, S. & Toni, I. (2011) Short notes 21. Hymenoptera, Formicidae. In: Nardi, G., Whitmore, D., Bardiani, M., Birtele, D., Mason, F., Spada, L. & Cerretti, P. (eds.) Biodiversity of Marganai and Montimannu (Sardinia). Research in the framework of the ICP Forests network. *Conservazione Habitat Invertebrati*, 5. Sommacampagna, Verona: Cierre Edizioni, Italy.
- Salata, S. & Borowiec, L. (2015) A taxonomic revision of the genus *Oxyopomyrmex* André, 1881 (Hymenoptera: Formicidae). *Zootaxa*, 4025, 1-66. DOI: 10.11646/zootaxa.4025.1.1
- Salata, S. & Borowiec, L. (2018a) Redescription of *Aphaenogaster muschtaidica* Emery, 1908 with a key to *gibbosa* species group. *Asian Myrmecology*, 10, e010002. DOI: 10.20362/am.010002
- Salata, S. & Borowiec, L. (2018b). Taxonomic and faunistic notes on Greek ants (Hymenoptera: Formicidae). *Annals of the Upper Silesian Museum in Bytom Entomology*, 27, 1-51. DOI: 10.5281/zenodo.2199191
- Salata, S., Borowiec, L. & Trichas, A. (2018) Taxonomic revision of the Cretan fauna of the genus *Temnothorax* Mayr, 1861 (Hymenoptera: Formicidae), with notes on the endemism of ant fauna of Crete. *Annales Zoologici*, 68, 769-808. DOI: 10.3161/00034541ANZ2018.68.4.004
- Salata, S., Karaman, C., Kiran, K. & Borowiec, L. (2021). Review of the *Aphaenogaster splendida* species-group (Hymenoptera: Formicidae). *Annales Zoologici*, 71, 297-343. DOI: 10.3161/00034541ANZ2021.71.2.008
- Sanetra, M., Güsten, R. & Schulz, A. (1999) On the taxonomy and distribution of Italian *Tetramorium* species and their social parasites (Hymenoptera Formicidae). *Memorie della Società Entomologica Italiana*, 77, 317-357.
- Sarnat, E.M., Fischer, G., Guénard, B. & Economo, E.P. (2015) Introduced *Pheidole* of the world: taxonomy, biology and distribution. *ZooKeys*, 543, 1-109. DOI: 10.3897/zookeys.543.6050
- Schär, S., Menchetti, M., Schifani, E., Hinojosa, J. C., Platania, L., Dapporto, L. & Vila, R. (2020) Integrative biodiversity inventory of ants from a Sicilian archipelago reveals high diversity on young volcanic islands (Hymenoptera: Formicidae). *Organisms Diversity and Evolution*, 20, 405-416. DOI: 10.1007/s13127-020-00442-3
- Schifani, E. (2017) First record of the vulnerable social parasite ant *Plagiolepis grassei* in Italy (Hymenoptera: Formicidae). *Fragmenta entomologica*, 49, 61-64. DOI: 10.4081/fe.2017.231
- Schifani, E. (2019) Exotic ants (Hymenoptera, Formicidae) invading Mediterranean Europe: a brief summary over about 200 years of documented introductions. *Sociobiology*, 66, 198-208. DOI: 10.13102/sociobiology.v66i2.4331
- Schifani, E. & Alicata, A. (2018) Exploring the myrmecofauna of Sicily: thirty-two new ant species recorded, including six new to Italy and many new aliens (Hymenoptera, Formicidae). *Polish Journal of Entomology*, 87, 323-348. DOI: 10.2478/pjen-2018-0023
- Schifani, E. & Alicata, A. (2019) *Aphaenogaster finzii* Müller, 1921, a trans-Ionian species new to Italy (Hymenoptera, Formicidae). *Biogeographia*, 34, 51-57. DOI: 10.21426/B634043636
- Schifani, E., Scupola, A. & Alicata, A. (2020) Morphology, ecology and biogeography of *Myrmecina sicula* André, 1882, rediscovered after 140 years (Hymenoptera, Formicidae). *Biogeographia*, 35, 105-116. DOI: 10.21426/B635048444

- Schifani, E., Costa, S., Mei, M., & Alicata, A. (2021a). A new species for the Italian fauna: *Aphaenogaster strioloides*, not *A. crocea*, inhabits Pantelleria Island (Hymenoptera: Formicidae). *Fragmenta Entomologica*, 53, 21-23. DOI: 10.13133/2284-4880/482
- Schifani, E., Csósz, S., Viviano, R. & Alicata, A. (2021b). Ant diversity on the largest Mediterranean islands: on the presence or absence of 28 species in Sicily (Hymenoptera, Formicidae). *Natural History Sciences*, 8, 55-70. DOI: 10.4081/nhs.2021.532
- Schifani, E., Nalini, E., Gentile, V., Alamanni, F., Ancona, C., Caria, M., Cillo, D. & Bazzato, E. (2021c). Ants of Sardinia: an updated checklist based on new faunistic, morphological and biogeographical notes. *Redia*, 104, 21-35. DOI: 10.19263/REDIA-104.21.03
- Schifani, E., Scupola, A., Menchetti, M., Bazzato, E. & Espadaler, X. (2021d). Morphology and Phenology of Sexually and New Distribution Data on the Blind Mediterranean Ant *Hypoponera abeillei* (Hymenoptera, Formicidae). *Sociobiology*, 68, e7261. DOI: 10.13102/sociobiology.v68i4.7261
- Schifani, E., Castracani, C., Spotti, F.A., Giannetti, D., Ghizzoni, M., Gobbi, M., Lencioni, V., Pedrotti, L., Grasso, D.A. & Mori, A. (2021e). Social Parasite Ants in the Alps: a New Site of the Vulnerable *Myrmica myrmicoxena* and New Uppermost Altitudinal Limit for *M. microrubra*. *Sociobiology*, 68, e7276. DOI: 10.13102/sociobiology.v68i4.7276
- Schifani, E., Alicata, A. & Menchetti, M. (2022a). Following the Apennines: updating the distribution of *Formica clara* and *F. rufibarbis* in Italy (Hymenoptera, Formicidae). *Biogeographia*, 37, a014. DOI: 10.21426/B637155107
- Schifani, E., Giannetti, D., Csósz, S., Castellucci, F., Luchetti, A., Castracani, C., Spotti, F.A., Mori, A. & Grasso, D.A. (2022b). Is mimicry a diversification-driver in ants? Biogeography, ecology, ethology, genetics and morphology define a second West-Palaearctic *Colobopsis* species (Hymenoptera: Formicidae). *Zoological Journal of the Linnean Society*, zlab035. DOI: 10.1093/zoolinnean/zlab035
- Schifani, E., Prebus, M.M. & Alicata, A. (2022c). Integrating morphology with phylogenomics to describe four island endemic *Temnothorax* species from Sicily and Malta (Hymenoptera, Formicidae). *European Journal of Taxonomy*, in press.
- Schulz, A., Heinze, J. & Pusch, K. (2007). Description of two new *Temnothorax* species (Hymenoptera: Formicidae) from Italy. *Zootaxa*, 1471, 1-14. DOI: 10.11646/zootaxa.1471.1.1
- Scupola, A. (2006) *Ponera coarctata* var. *crassissquama* Emery, 1916 a new synonym of *P. testacea* Emery, 1895 (Hymenoptera, Formicidae). *Bullettino del Museo Civico di Storia Naturale di Verona*, 30, 161-164.
- Scupola, A. & Ballarin, R. (2009) The genus *Leptanilla* Emery, 1870 in Sicily (Hymenoptera: Formicidae). *Myrmecological News*, 12, 129-132.
- Scupola, A. (2017) *Aphaenogaster muelleriana* Wolf, 1915 (Hymenoptera Formicidae) in Salento (South East Italy). *Biodiversity Journal*, 8, 3-8.
- Scupola, A. (2018) Le formiche del Veneto. WBA Handbooks, Italy.
- Scupola, A., Durante, A., Giannuzzi, F. & Pellizzari, G. (2022) The coccid-tending ant genus *Acropyga* Roger and its obligate associated myrmecophilous scale insect genus *Eumyrmococcus* Silvestri new to Italy (Hymenoptera: Formicidae; Hemiptera: Xenococcidae). *Fragmenta Entomologica*, 54, 89-94. DOI: 10.13133/2284-4880/722
- Seifert, B. (1992a) *Formica nigricans* Emery, 1909 - an ecomorph of *Formica pratensis* Retzius, 1783 (Hymenoptera, Formicidae). *Entomologica Fennica*, 2, 217-226.
- Seifert, B. (1992b) A taxonomic revision of the Palaearctic members of the ant subgenus *Lasius* s.str. (Hymenoptera: Formicidae). *Abhandlungen und Berichte des Naturkundemuseums Görlitz*, 66, 1-67.
- Seifert, B. (1994) Die freilebenden Ameisenarten Deutschlands (Hymenoptera: Formicidae) und Angaben zu deren Taxonomie und Verbreitung. *Abhandlungen und Berichte des Naturkundemuseums Görlitz*, 67, 1-44.

- Seifert, B. (2000) A taxonomic revision of the ant subgenus *Coptoformica* Mueller, 1923 (Hymenoptera: Formicidae). *Zoosystema*, 22, 517-568.
- Seifert, B. (2002) A taxonomic revision of the *Formica cinerea* group (Hymenoptera: Formicidae). *Abhandlungen und Berichte des Naturkundemuseums Görlitz*, 74, 245-272.
- Seifert, B. (2005) Rank elevation in two European ant species: *Myrmica lobulicornis* Nylander, 1857, stat. n. and *Myrmica spinosior* Santschi, 1931, stat. n. (Hymenoptera: Formicidae). *Myrmecologische Nachrichten*, 7, 1-7.
- Seifert, B., Schlick-Steiner, B.C. & Steiner, F.M. (2009) *Myrmica constricta* Karavajev, 1934 – a cryptic sister species of *Myrmica hellenica* Finzi, 1926. *Soil Organisms*, 81, 53-76.
- Seifert, B. (2012a) Clarifying naming and identification of the outdoor species of the ant genus *Tapinoma* Förster, 1850 in Europe north of the Mediterranean region with description of a new species. *Myrmecological News*, 16, 139-147.
- Seifert, B. (2012b) A review of the West Palaearctic species of the ant genus *Bothriomyrmex* Emery, 1869. *Myrmecological News*, 17, 91-104.
- Seifert, B., Csösz, S. & Schulz, A. (2014) NC-Clustering demonstrates heterospecificity of the cryptic ant species *Temnothorax luteus* (FOREL, 1874) and *T. racovitzai* (Bondroit, 1918) (Hymenoptera: Formicidae). *Beiträge zur Entomologie*, 64, 47-57. DOI: 10.21248/contrib.entomol.6.4.147-57
- Seifert, B. & Csösz, S. (2015) *Temnothorax crasecundus* sp. n.—a cryptic Eurocaucasian ant species (Hymenoptera, Formicidae) discovered by Nest Centroid Clustering. *ZooKeys*, 479, 37-64. DOI: 10.3897/zookeys.479.8510
- Seifert, B., D'Eustacchio, D., Kaufmann, B., Centorame, M., Lorite, P. & Modica, M. (2017) Four species within the supercolonial ants of the *Tapinoma nigerrimum* complex revealed by integrative taxonomy (Hymenoptera: Formicidae). *Myrmecological News*, 24, 123-144.
- Seifert, B. (2018) Ants of Northern and Central Europe. Lutra Verlags- und Vertriebsgesellschaft, Germany.
- Seifert, B. (2020) A taxonomic revision of the Palaearctic members of the subgenus *Lasius* s. str. (Hymenoptera, Formicidae). *Soil Organisms*, 92, 15-86. DOI: 10.25674/so92iss1pp15
- Social Insect Specialists Group (1996). The IUCN Red List of Threatened Species 1996: e.T17464A7087398.
- Steiner, F.M., Seifert, B., Moder, K. & Schlick-Steiner, B.C. (2010) A multisource solution for a complex problem in biodiversity research: Description of the cryptic ant species *Tetramorium alpestre* sp. n. (Hymenoptera: Formicidae). *Zoologischer Anzeiger*, 249, 223-254. DOI: 10.1016/j.jcz.2010.09.003
- Steiner, F. M., Csösz, S., Markó, B., Gamisch, A., Rinnhofer, L., Folterbauer, C., Hammerle, S., Stauffer, C., Arthofer, W. & Schlick-Steiner, B.C. (2018) Turning one into five: Integrative taxonomy uncovers complex evolution of cryptic species in the harvester ant *Messor "struttator"*. *Molecular Phylogenetics and Evolution*, 127, 387-404.
- Süss, L. (1979) Formiche infestanti l'industria alimentare: esperienze di difesa antiparassitaria. In Domenichini, G. (ed.) Atti 2° Simposio “La difesa antiparassitaria nelle industrie alimentari e la protezione degli alimenti”, Tip. Le Co., Italy.
- Tinaut, A. & Ruano, F. (2021) Biogeography of Iberian Ants (Hymenoptera: Formicidae). *Diversity*, 13, 88. DOI: 10.3390/d13020088
- Verdinelli, M., Sassu, A., Molinu, A. & Fois, X. (2007) An updated list of Sardinia's ants (Hymenoptera Formicidae). *Redia*, 90, 61-66.
- Wagner, H.C., Arthofer, W., Seifert, B., Muster, C., Steiner, F.M. & Schlick-Steiner, B.C. (2017) Light at the end of the tunnel: Integrative taxonomy delimits cryptic species in the *Tetramorium caespitum* complex. *Myrmecological News*, 25, 95-129.
- Wang, R., Kass, J.M., Galkowski, C., Garcia, F., Hamer, M.T., Radchenko, A., Salata, S., Schifani, E., Yusupov, Z.M., Economo, E.P. & Guénard, B. (2022). New distribution data and phylogenetic approach reveal bioregionalization of the western Palearctic ants. *bioRxiv*. DOI: 10.1101/2022.03.10.483749

- Ward, P.S. (2007). Phylogeny, classification, and species-level taxonomy of ants (Hymenoptera: Formicidae). *Zootaxa*, 1668, 549-563. DOI: 10.11646/zootaxa.1668.1.26
- Ward, P.S. (2014) The phylogeny and evolution of ants. *Annual Review of Ecology, Evolution, and Systematics*, 45, 23-43. DOI: 10.1146/annurev-ecolsys-120213-091824
- Ward, P.S., Brady, S.G., Fisher, B.L. & Schultz, T.R. (2015) The evolution of myrmicine ants: phylogeny and biogeography of a hyperdiverse ant clade (Hymenoptera: Formicidae). *Systematic Entomology*, 40, 61-81. DOI: 10.1111/syen.12090
- Ward, P.S., Blaimer, B.B. & Fisher, B.L. (2016) A revised phylogenetic classification of the ant subfamily Formicinae (Hymenoptera: Formicidae), with resurrection of the genera *Colobopsis* and *Dinomyrmex*. *Zootaxa*, 4072, 343-357. DOI: 10.11646/zootaxa.4072.3.4
- Wetterer, J.K., Espadaler, X., Wetterer, A.L., Aguin-Pombo, D. & Franquinho-Aguiar, A.M. (2007). Ants (Hymenoptera: Formicidae) of the Madeiran Archipelago. *Sociobiology*, 49, 265-297.
- Yoshimura, M. & Fisher, B.L. (2012) A revision of male ants of the Malagasy Amblyoponinae (Hymenoptera: Formicidae) with resurrections of the genera *Stigmatomma* and *Xymmer*. *PLoS One*, 7, e33325. DOI: 10.1371/journal.pone.0033325.

Submitted: 26 December 2021

First decision: 20 April 2022

Accepted: 25 May 2022

*the Guest editors for the special section
“The new Checklist of the Italian Fauna”:*

*Marco A. Bologna, Lucio Bonato,
Fabio Cianferoni, Alessandro Minelli,
Marco Oliverio, Fabio Stoch, Marzio Zapparoli*

Datapaper citation:

Schifani E., 2022. The new Checklist of the Italian Fauna: Formicidae. Biogeographia, 37, ucl006

Supplementary file S1

List of all species with occurrence in the geographical areas of the new Checklist of the Italian Fauna

Legend: a, alien; e, endemic; s, subendemic

Citation of the contribution in the Checklist online:

Schifani E., 2021. Formicidae. In: Bologna M.A., Zapparoli M., Oliverio M., Minelli A., Bonato L., Cianferoni F., Stoch F. (eds.), Checklist of the Italian fauna. Version 1.0. Last update: 2021-05-31.

	endemic	alien	Continental Italy	Peninsular Italy	Sicilia	Sardegna	taxonomic remarks		occurrence remarks
Amblyoponinae									
<i>Stigmatomma denticulatum</i> Roger, 1859	-	-	+	+	+	-	Yoshimura & Fisher 2012 (PLoS One, 7: e33325): combination in <i>Stigmatomma</i>		Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348): Si
<i>Stigmatomma impressifrons</i> Emery, 1869	-	-	+	+	+	-	Yoshimura & Fisher 2012 (PLoS One, 7: e33325): combination in <i>Stigmatomma</i>		
Dolichoderinae									
<i>Bothriomyrmex communis</i> Santschi, 1919	-	-	+	+	+	-	Seifert 2012 (Myrmecological News, 17: 91-104): senior synonym of <i>B. adriacus</i>		
<i>Bothriomyrmex corsicus</i> Santschi, 1923	-	-	+	+	+	-	Seifert 2012 (Myrmecological News, 17: 91-104): senior synonym of <i>B. gibbus</i> , <i>B. liguricus</i> , <i>B. menozzii</i>	Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348): Si	
<i>Dolichoderus quadripunctatus</i> (Linnaeus, 1771)	-	-	+	+	+	-			Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348): Si
<i>Linepithema humile</i> (Mayr, 1868)	-	a	+	+	+	+			
<i>Liometopum microcephalum</i> (Panzer, 1798)	-	-	+	+	+	-			
<i>Tapinoma darioi</i> Seifert, D'Eustacchio, Kaufmann, Centorame & Modica, 2017	-	-	+	+	-	-			Seifert et al. 2017 (Myrmecological News, 24: 123-144): N, S (description)
<i>Tapinoma erraticum</i> (Latreille, 1798)	-	-	+	+	+	+			
<i>Tapinoma madeirense</i> Forel, 1895	-	-	-	-	+	+	Wetterer et al. 2007 (Sociobiology, 49: 265-297): senior synonym of <i>T. ambiguum</i>	Rigato & Toni 2011 (Research in the framework of the ICP Forests network. Conservazione Habitat Invertebrati, 5: 873-882): Sa; Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348): Si	
<i>Tapinoma magnum</i> Mayr, 1861	-	-	+	+	+	+			Seifert et al. 2017 (Myrmecological News, 24: 123-144): N, S, Sa, Si
<i>Tapinoma pygmaeum</i> (Dufour, 1857)	-	-	+	-	-	-			
<i>Tapinoma simrothi</i> Krausse, 1911	-	-	-	-	+	+			
<i>Tapinoma subboreale</i> Seifert, 2012	-	-	+	+	-	-			Glaser et al. 2012 (Gredleriana, 12: 273-284): N; Seifert et al. 2017 (Myrmecological News, 24: 123-144): S
<i>Technomyrmex pallipes</i> (Smith, 1876)	-	a	+	-	-	-			Jucker et al. 2008 (Bollettino di Zoologia agraria e di Bachicoltura, 40: 99-107): N

	endemic	alien	Continental Italy	Peninsular Italy	Sicilia	Sardegna	taxonomic remarks	occurrence remarks
Formicinae								
<i>Camponotus aethiops</i> (Latreille, 1798)	-	-	+	+	+	+		
<i>Camponotus atriceps</i> (Smith, 1858)	-	a	+	-	-	-		Jucker et al. 2008 (Bollettino di Zoologia agraria e di Bachicoltura, 40: 99-107): N
<i>Camponotus barbaricus</i> Emery, 1905	-	-	-	-	+	-		
<i>Camponotus cruentatus</i> (Latreille, 1802)	-	-	+	-	-	-		
<i>Camponotus dalmaticus</i> (Nylander, 1849)	-	-	+	+	-	-		
<i>Camponotus fallax</i> (Nylander, 1856)	-	-	+	+	-	-		
<i>Camponotus gestroi</i> Emery, 1878	-	-	-	-	+	+		
<i>Camponotus herculeanus</i> (Linnaeus, 1758)	-	-	+	-	-	-		
<i>Camponotus lateralis</i> (Olivier, 1792)	-	-	+	+	+	+		
<i>Camponotus ligniperda</i> (Latreille, 1802)	-	-	+	+	+	-		
<i>Camponotus micans</i> (Nylander, 1856)	-	-	-	+	+	-		
<i>Camponotus nylanderi</i> Emery, 1921	e	-	-	+	+	-		
<i>Camponotus piceus</i> (Leach, 1825)	-	-	+	+	+	+		
<i>Camponotus pilicornis siculo</i> Grandi, 1935	e	-	-	-	+	-		
<i>Camponotus ruber</i> Emery, 1925	-	-	-	-	+	-	Cagniant 1996 (Annales de la Société Entomologique de France, 32: 87-100): species status	
<i>Camponotus spissinodis</i> Forel, 1909	-	-	-	-	+	-		
<i>Camponotus sylvaticus</i> (Olivier, 1792)	-	-	+	-	-	-		
<i>Camponotus tergestinus</i> Müller, 1921	-	-	+	+	+	-		Braško 2017 (Natura Sloveniae, 19: 5-14): S; Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348): Si
<i>Camponotus universitatis</i> Forel, 1890	-	-	-	+	+	+		Kutter 1977 (Insecta Helvetica: Fauna, 6: 1-298): Si; Rigato & Toni 2011 (Research in the framework of the ICP Forests network. Conservazione Habitat Invertebrati, 5: 873-882) (Research in the framework of the ICP Forests network. Conservazione Habitat Invertebrati, 5: 873-882): Sa
<i>Camponotus vagus</i> (Scopoli, 1763)	-	-	+	+	+	+		
<i>Cataglyphis italicica</i> (Emery, 1906)	e	-	-	+	-	-		
<i>Cataglyphis viatica</i> (Fabricius, 1787)	-	-	-	-	+	-		Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348): Si
<i>Colobopsis truncata</i> (Spinola, 1808)	-	-	+	+	-	+		
<i>Formica aquilonia</i> Yarrow, 1955	-	-	+	-	-	-		
<i>Formica bruni</i> Kutter, 1967	-	-	+	-	-	-		Seifert 2000 (Zoosistema, 22: 517-568): N
<i>Formica cinerea</i> Mayr, 1853	-	-	+	+	-	-	Agosti & Collingwood 1987 (Mitteilungen der Schweizerischen Entomologischen Gesellschaft, 60: 261-293): senior synonym of <i>F. subrufofides</i> ; Seifert 2002 (Abhandlungen und Berichte des Naturkundemuseums Görlitz, 74: 245-272): senior synonym of <i>F. cinerea italicica</i>	

	endemic	alien	Continental Italy	Peninsular Italy	Sicilia	Sardegna	taxonomic remarks	occurrence remarks
<i>Formica clara</i> Forel, 1886	-	-	+	+	-	+		Rigato & Toni 2011 (Research in the framework of the ICP Forests network. Conservazione Habitat Invertebrati, 5: 873-882): Sa; Glaser et al. 2012 (Gredieriana, 12: 273-284): N; Scupola 2016: S
<i>Formica cunicularia</i> Latreille, 1798	-	-	+	+	+	+		
<i>Formica exsecta</i> Nylander, 1846	-	-	+	+	-	-	Bernard 1967 (Masson, Paris, 411 pp.): senior synonym of <i>F. exsecta etrusca</i>	
<i>Formica foreli</i> Bondroit, 1918	-	-	+	-	-	-		
<i>Formica fusca</i> Linnaeus, 1758	-	-	+	+	-	-		Verdinelli et al. 2007 (Redia): absence from Sa
<i>Formica fuscocinerea</i> Forel, 1874	-	-	+	-	-	-	Seifert 2002 (Abhandlungen und Berichte des Naturkundemuseums Görlitz, 74: 245-272): species status	
<i>Formica gagates</i> Latreille, 1798	-	-	+	+	-	-		
<i>Formica lemani</i> Bondroit, 1917	-	-	+	+	-	-		
<i>Formica lugubris</i> Zetterstedt, 1838	-	-	+	+	+	+		
<i>Formica paralugubris</i> Seifert, 1996	s	-	+	+	-	-		Bernasconi et al. 2006 (Myrmecologische Nachrichten, 8: 251-256): N. Masoni et al. 2019 (Conservation Genetics Resources, 11: 231-236): S
<i>Formica picea</i> Nylander, 1846	-	-	+	-	-	-		Menozzi 1925 (Atti della Società dei Naturalisti e Matematici di Modena, 55: 22-4): N
<i>Formica polyctena</i> Foerster, 1850	-	-	+	-	-	-		
<i>Formica pratensis</i> Retzius, 1783	-	-	+	+	-	-	Seifert 1992 (Abhandlungen und Berichte des naturkundemuseums Görlitz, 66: 1-67) (Entomologica Fennica, 2: 217-226): senior synonym of <i>F. nigricans</i>	
<i>Formica pressilabris</i> Nylander, 1846	-	-	+	-	-	-		
<i>Formica rufa</i> Linnaeus, 1761	-	-	+	-	-	-		
<i>Formica rufibarbis</i> Fabricius, 1793	-	-	+	+	-	-	Bernard 1967 (Masson, Paris, 411 pp.): senior synonym of <i>F. cinereorufibarbis</i>	Rigato & Toni 2011 (Research in the framework of the ICP Forests network. Conservazione Habitat Invertebrati, 5: 873-882): Sa
<i>Formica sanguinea</i> Latreille, 1798	-	-	+	+	+	-		
<i>Formica selysi</i> Bondroit, 1918	-	-	+	-	-	-		
<i>Formica truncorum</i> Fabricius, 1804	-	-	+	-	-	-		
<i>Lasius alienus</i> (Foerster, 1850)	-	-	+	+	+	+		
<i>Lasius bicornis</i> (Foerster, 1850)	-	-	+	+	+	+		Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348): Si
<i>Lasius bombycinus</i> Seifert & Galkowski, 2016	-	-	+	-	-	-		Seifert 2020 (Soil organisms, 2020: 15-86): N
<i>Lasius brunneus</i> (Latreille, 1798)	-	-	+	+	+	+		
<i>Lasius carniolicus</i> Mayr, 1861	-	-	+	+	-	-		
<i>Lasius casevitzi</i> Seifert & Galkowski, 2016	s	-	+	-	+	-		Seifert 2020 (Soil organisms, 2020: 15-86): N; Schär et al. 2020 (Organisms Diversity and Evolution, 20: 405-416): Si;
<i>Lasius citrinus</i> Emery, 1922	-	-	+	+	-	-		Verdinelli et al. 2007 (Redia, 90: 61-66): absence in Sa

	endemic	alien	Continental Italy	Peninsular Italy	Sicilia	Sardegna	taxonomic remarks	occurrence remarks
<i>Lasius distinguendus</i> (Emery, 1916)	-	-	+	+	-	-		
<i>Lasius emarginatus</i> (Olivier, 1792)	-	-	+	+	+	+		
<i>Lasius flavus</i> (Fabricius, 1782)	-	-	+	+	+	+		
<i>Lasius fuliginosus</i> (Latreille, 1798)	-	-	+	+	+	-		
<i>Lasius lasioides</i> (Emery, 1869)	-	-	+	+	+	+		Rigato & Toni 2011 (Research in the framework of the ICP Forests network. Conservazione Habitat Invertebrati, 5: 873-882); Sa; Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348); Si; Seifert 2020 (Soil organisms, 2020: 15-86); N, S
<i>Lasius meridionalis</i> (Bondroit, 1920)	-	-	+	-	-	-		
<i>Lasius mixtus</i> (Nylander, 1846)	-	-	+	+	-	-		
<i>Lasius myops</i> Forel, 1894	-	-	+	+	+	-		
<i>Lasius neglectus</i> Van Loon, Boomsma & Andrasfalvy, 1990	-	a	+	+	-	-		Seifert 2000 (Deutsche Entomologische Zeitschrift, 47: 173-179): replaces <i>L. turcicus</i> in this region Seifert 2020 (Soil organisms, 2020: 15-86); S;
<i>Lasius niger</i> (Linnaeus, 1758)	-	-	+	+	+	+		
<i>Lasius nitidigaster</i> Seifert, 1996	-	-	+	-	-	-		Carniel 1998 (Atti del Museo Civico di Storia Naturale di Morbegno 9: 29-39); N
<i>Lasius paralienus</i> Seifert, 1992	-	-	+	+	-	+		Seifert 1992 (Abhandlungen und Berichte des naturkundemuseums Görlitz, 66: 1-67); N, S; Seifert & Galkowski 2016 (Zootaxa, 4132: 44-58); Sa
<i>Lasius platytorax</i> Seifert, 1991	-	-	+	+	-	+		Rigato & Toni 2011 (Research in the framework of the ICP Forests network. Conservazione Habitat Invertebrati, 5: 873-882); Sa; Seifert 2020 (Soil organisms, 2020: 15-86); N, S
<i>Lasius psammophilus</i> Seifert, 1992	-	-	+	+	+	-		Glaser 2004 (Gredleriana, 4: 203-246); N; Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348); Si; Seifert 2020 (Soil organisms, 2020: 15-86); S
<i>Lasius reginæ</i> Faber, 1967	-	-	+	-	-	-		Glaser et al. 2012 (Gredleriana, 12: 273-284); N
<i>Lasius umbratus</i> (Nylander, 1846)	-	-	+	+	+	-		Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348); Si
<i>Lepisiota frauenfeldi</i> (Mayr, 1855)	-	-	-	+	+	-		
<i>Lepisiota nigra</i> (Dalla Torre, 1893)	-	-	-	+	+	-		
<i>Nylanderia jaegerskioeldi</i> (Mayr, 1904)	-	a	-	-	+	-		Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348); Si
<i>Paratrechina longicornis</i> (Latreille, 1802)	-	a	+	-	+	-		Süss 1979 (Tip.Le.Co., Piacenza, 359-366); N; Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348); Si
<i>Plagiolepis ampeloni</i> (Faber, 1969)	-	-	+	-	-	-		Buschinger et al. 1999; N
<i>Plagiolepis grassei</i> Le Masne, 1956	-	-	-	-	+	-		Schifani 2017; Si
<i>Plagiolepis pallescens</i> Forel, 1889	-	-	+	+	+	-	Salata et al. 2018 (Zookeys, 543: 1-19): senior synonym of <i>P. vindobonensis</i>	Schär et al. 2020 (Organisms Diversity and Evolution, 20: 405-416); Si
<i>Plagiolepis pygmaea</i> (Latreille, 1798)	-	-	+	+	+	+	Boer 2008 (Zoologische Mededelingen, 82: 485-488): senior synonym of <i>P. obscuriscapa</i>	
<i>Plagiolepis schmitzii</i> Forel, 1895	-	-	-	-	+	-		
<i>Plagiolepis xene</i> Stärcke, 1936	-	-	+	+	+	+		Loi 2013 (Tesi di Dottorato, Università degli studi di Sassari); Sa

	endemic	alien	Continental Italy	Peninsular Italy	Sicilia	Sardegna	taxonomic remarks	occurrence remarks
<i>Polyergus rufescens</i> (Latreille, 1798)	-	-	+	+	-	-		
<i>Prenolepis nitens</i> (Mayr, 1853)	-	-	+	-	-	-		
Leptanillinae								
<i>Leptanilla doderoi</i> Emery, 1915	e	-	-	-	-	+		
<i>Leptanilla poggi</i> Mei, 1995	e	-	-	-	+	-		Mei 1995 (Il Naturalista Siciliano, 19: 753-772): Si
<i>Leptanilla revelieri</i> Emery, 1870	-	-	-	-	-	+	Baroni Urbani 1977 (Entomologica Basiliensis, 2: 427-488): senior synonym of <i>L. revelierii sardoa</i>	
Myrmicinae								
<i>Aphaenogaster campana</i> Emery, 1878	e	-	-	+	-	-		
<i>Aphaenogaster crocea</i> André, 1881	-	-	-	-	+	-		
<i>Aphaenogaster dulcinea</i> Emery, 1924	-	-	+	-	-	-		
<i>Aphaenogaster epirotes</i> (Emery, 1895)	-	-	+	-	-	-		
<i>Aphaenogaster finzii</i> Müller, 1921	-	-	-	+	-	-		Schifani & Alicata 2019 (Biogeographia, 34): S
<i>Aphaenogaster fiorii</i> Emery, 1915	s	-	-	-	+	-		
<i>Aphaenogaster gibbosa</i> (Latreille, 1798)	-	-	+	-	-	-	Alicata & Schifani 2019 (Acta Entomologica Musei Nationalis Pragae, 59: 1-16): species status	
<i>Aphaenogaster ichnusa</i> Santschi, 1925	-	-	-	+	+	+	Galkowski et al. 2018 (Sociobiology, 66: 420-425): absence in Sa	Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348): S, Si
<i>Aphaenogaster inermita</i> Bolton, 1995	s	-	-	+	-	-		
<i>Aphaenogaster italicica</i> Bondroit, 1918	s	-	+	+	+	-		Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348): Si
<i>Aphaenogaster muelleriana</i> Wolf, 1915	-	-	-	+	-	-		Scupola 2017 (Biodiversity Journal, 8: 3-8): S
<i>Aphaenogaster ovaticeps</i> (Emery, 1898)	e	-	+	-	-	-		
<i>Aphaenogaster pallida</i> (Nylander, 1849)	-	-	-	+	+	-		
<i>Aphaenogaster picena</i> Baroni Urbani, 1971	e	-	-	+	-	-		
<i>Aphaenogaster sardoa</i> Mayr, 1853	-	-	-	-	+	+		
<i>Aphaenogaster semipolita</i> (Nylander, 1856)	e	-	-	+	+	-		Boer 2013 (Tijdschrift voor entomologie, 156: 57-93): absence of <i>A. ionia</i> , junior synonym of <i>A. balcanica</i>
<i>Aphaenogaster senilis</i> Mayr, 1853	-	-	-	-	-	+		
<i>Aphaenogaster sicula</i> Emery, 1808	e	-	-	+	+	-		Alicata & Schifani 2019 (Acta Entomologica Musei Nationalis Pragae, 59: 1-16): S
<i>Aphaenogaster spinosa</i> Emery, 1878	-	-	+	+	-	+	Baroni Urbani 1971 (Memorie della Società Entomologica Italiana, 50: 5-287): senior synonym of <i>A. spinosa nitida</i> ; Boer 2013 (Tijdschrift voor entomologie, 156: 57-93): senior synonym of <i>A. spinosa etrusca</i>	
<i>Aphaenogaster splendida</i> (Roger, 1859)	-	-	+	+	+	-		
<i>Aphaenogaster subterranea</i> (Latreille, 1798)	-	-	+	+	+	-		Galkowski et al. 2018 (Sociobiology, 66: 420-425): absence in Sa
<i>Aphaenogaster trinacriae</i> Alicata & Schifani,	e	-	-	-	+	-		Alicata & Schifani 2019 (Acta Entomologica Musei Nationalis Pragae, 59: 1-16): S

	endemic	alien	Continental Italy	Peninsular Italy	Sicilia	Sardegna	taxonomic remarks	occurrence remarks
2019								(description)
<i>Cardiocondyla elegans</i> Emery, 1869	-	-	+	+	-	-		
<i>Cardiocondyla mauritanica</i> Forel, 1890	-	?	+	+	+	+	Ortiz & Tinaut 1987: species status	
<i>Crematogaster laestrygon</i> Emery, 1869	-	-	-	-	+	-		
<i>Crematogaster schmidti</i> (Mayr, 1853)	-	-	+	-	-	-		
<i>Crematogaster scutellaris nigra</i> Krausse, 1912	-	-	-	-	-	+		
<i>Crematogaster scutellaris scutellaris</i> (Olivier, 1792)	-	-	+	+	+	+		
<i>Crematogaster sordidula</i> (Nylander, 1849)	-	-	+	+	+	-	Bolton 1995 (Cambridge, Mass.: Harvard University Press, 504 pp.): senior synonym of <i>C. mayri</i>	
<i>Formicoxenus nitidulus</i> (Nylander, 1846)	-	-	+	-	-	-		
<i>Harpagoxenus sublaevis</i> (Nylander, 1849)	-	-	+	-	-	-		
<i>Leptothorax acervorum</i> (Fabricius, 1793)	-	-	+	+	-	-		
<i>Leptothorax gredleri</i> Mayr, 1855	-	-	+	-	-	-		
<i>Leptothorax kutteri</i> Buschinger, 1966	-	-	+	-	-	-	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Leptothorax</i>	
<i>Leptothorax muscorum</i> (Nylander, 1846)	-	-	+	+	-	-		
<i>Leptothorax pacis</i> (Kutter, 1945)	-	-	+	-	-	-	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Leptothorax</i>	
<i>Manica rubida</i> (Latreille, 1802)	-	-	+	-	-	-		
<i>Messor barbarus</i> (Linnaeus, 1767)	-	-	+	-	-	-		Mei 1995 (Il Naturalista Siciliano, 19: 753-772): absence in Si
<i>Messor bouvieri</i> Bondroit, 1918	-	-	+	-	+	-		
<i>Messor capitatus</i> (Latreille, 1798)	-	-	+	+	+	+		
<i>Messor ibericus</i> Santschi, 1931	-	-	+	+	+	+	Steiner et al. 2018 (Molecular phylogenetics and evolution, 127: 387-404): replaces <i>M. structor</i> in the Western-Mediterranean basin	Steiner et al. 2018 (Molecular phylogenetics and evolution, 127: 387-404): N; Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348): Si
<i>Messor minor calabricus</i> Santschi, 1927	e	-	-	+	-	-		
<i>Messor minor capreensis</i> Santschi, 1927	e	-	-	+	-	-		
<i>Messor minor minor</i> (André, 1883)	-	-	-	+	+	+		
<i>Messor sanctus</i> Emery, 1921	-	-	-	-	+	-		
<i>Messor wasmanni</i> Krausse, 1910	-	-	-	+	-	+	Casevitz-Weulersse 1990 (Revue d'écologie et de biologie du sol, 27: 29-59): raised to species	
<i>Monomorium monomorium</i> Bolton, 1987	-	-	+	+	+	-		
<i>Monomorium pharaonis</i> (Linnaeus, 1758)	a	+	+	-	-	-		
<i>Monomorium sommieri</i> Emery, 1916	-	-	-	-	+	-		
<i>Monomorium subopacum</i> (Smith, 1858)	-	-	-	+	+	-		

	endemic	alien	Continental Italy	Peninsular Italy	Sicilia	Sardegna	taxonomic remarks	occurrence remarks
<i>Myrmecina graminicola</i> (Latreille, 1802)	-	-	+	+	+	+		
<i>Myrmecina melonii</i> Rigato, 1999	e	-	-	-	-	+		
<i>Myrmecina sicula</i> André, 1882	e	-	-	-	+	-		
<i>Myrmica constricta</i> Karavaiev, 1934	-	-	+	-	-	-	Seifert et al. 2009 (Soil Organisms, 81: 53-76): species status	
<i>Myrmica hellenica</i> Finzi, 1926	-	-	+	-	-	-		
<i>Myrmica karavajevi</i> (Arnoldi, 1930)	-	-	+	-	-	-	Radchenko & Elmes 2010 (Warszawa: Natura optima dux Foundation): N	
<i>Myrmica laurae</i> (Emery, 1907)	e	-	-	+	-	-	Radchenko & Elmes 2003 (Annales Zoologici, Vol. 53: 217-243): senior synonym of <i>M. laurae</i>	
<i>Myrmica lobicornis</i> Nylander, 1846	-	-	+	+	-	-		
<i>Myrmica lobulicornis</i> Nylander, 1857	-	-	+	+	+	-	Seifert 2005 (Myrmecologische Nachrichten, 7: 1-7): species level	Radchenko & Elmes 2010 (Warszawa: Natura optima dux Foundation): N; Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348): Si
<i>Myrmica lonae</i> Finzi, 1926	-	-	+	-	-	-	Seifert 1994: species level	
<i>Myrmica myrmicoxena</i> Forel, 1895	s	-	+	-	-	-		Glaser et al. 2010 (Myrmecological News, 14: 107-111): N
<i>Myrmica obscura</i> Finzi, 1926	e	-	-	+	-	-		
<i>Myrmica rubra</i> (Linnaeus, 1758)	-	-	+	+	-	-		
<i>Myrmica ruginodis</i> Nylander, 1846	-	-	+	-	-	-		
<i>Myrmica rugulosa</i> Nylander, 1849	-	-	+	-	-	-		
<i>Myrmica sabuleti</i> Meinert, 1861	-	-	+	+	+	-		Verdinelli et al. 2007 (Redia, 90: 61-66): absence in Sa
<i>Myrmica scabrinodis</i> Nylander, 1846	-	-	+	+	+	-		
<i>Myrmica schencki</i> Viereck, 1903	-	-	+	+	-	-		
<i>Myrmica siciliana</i> Radchenko, Elmes & Alicata, 2006	e	-	-	-	+	-		Radchenko et al. 2006 (Annales Zoologici, 56:499-538): Si (description)
<i>Myrmica speciooides</i> Bondroit, 1918	-	-	+	+	-	-	Radchenko & Elmes 2010 (Warszawa: Natura optima dux Foundation): senior synonym of <i>M. striata</i>	
<i>Myrmica spinosior</i> Santschi, 1931	-	-	+	+	-	+	Seifert 2005 (Myrmecologische Nachrichten, 7: 1-7): species level	Verdinelli et al. 2007 (Redia, 90: 61-66): Sa
<i>Myrmica sulcinodis</i> Nylander, 1846	-	-	+	+	-	-		
<i>Myrmica tulinae</i> Elmes, Radchenko & Aktac, 2002	-	-	+	-	-	-	Seifert 2018 (Lutra Verlags- und Vertriebsgesellschaft: Tauer, Germany): possibly an occasional hybrid form of <i>M. sabuleti</i> and <i>M. scabrinodis</i> and not a species	Bonelli et al. 2004 (Congresso Nazionale dell'Unione Zoologica Italiana): N
<i>Myrmica vandeli</i> Bondroit, 1920	-	-	+	-	-	-		Seifert 2018 (Lutra Verlags- und Vertriebsgesellschaft: Tauer, Germany): N
<i>Oxyopomyrmex saulcyi</i> Emery, 1889	-	-	-	+	+	-	Salata & Borowiec 2015 (Zootaxa, 4225: 1-66): senior synonym of <i>Oxyopomyrmex santschii</i>	
<i>Pheidole indica</i> Mayr, 1879	-	a	-	-	+	-	Sarnat et al. 2015: senior synonym of <i>P. teneriffana</i>	Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348): Si
<i>Pheidole pallidula</i> (Nylander, 1849)	-	-	+	+	+	+		
<i>Solenopsis fugax</i> (Latreille, 1798)	-	-	+	+	-	-		
<i>Solenopsis latro sicula</i> Emery, 1915	e	-	-	-	+	-		

	endemic	alien	Continental Italy	Peninsular Italy	Sicilia	Sardegna	taxonomic remarks	occurrence remarks
<i>Solenopsis lusitanica</i> Emery, 1915	-	-	-	-	-	+		
<i>Solenopsis orbula</i> Emery, 1875	-	-	-	+	-	+		
<i>Stenamma debile</i> (Foerster, 1850)	-	-	+	+	+	+		Rigato 2011 (Memorie della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano, 37: 1-56); Rigato & Toni 2011 (Research in the framework of the ICP Forests network. Conservazione Habitat Invertebrati, 5: 873-882): Sa
<i>Stenamma petiolatum</i> Emery, 1897	e	-	+	+	+	-		Rigato 2011 (Memorie della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano, 37: 1-56): endemic to Italy, since records from other regions are dubious. Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348): Si
<i>Stenamma sardoum</i> Emery, 1915	-	-	-	-	-	+		
<i>Stenamma siculum</i> Rigato, 2011	e	-	-	-	+	-		Rigato 2011 (Memorie della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano, 37: 1-56): Si (description)
<i>Stenamma striatulum</i> Emery, 1895	-	-	+	+	-	+		
<i>Stenamma zanoni</i> Rigato, 2011	s	-	+	-	-	-		Rigato 2011 (Memorie della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano, 37: 1-56): N (description)
<i>Strongylognathus alboini</i> Finzi, 1924	s	-	+	-	-	-		
<i>Strongylognathus alpinus</i> Wheeler, 1909	s	-	+	+	+	-		Sanetra et al. 1999 (Memorie della Società Entomologica Italiana, 77: 317-357): Si
<i>Strongylognathus destefanii</i> Emery, 1915	e	-	-	+	+	-	Sanetra et al. 1999 (Memorie della Società Entomologica Italiana, 77: 317-357): senior synonym of <i>S. cecconii</i> and <i>S. emeryi</i>	Sanetra et al. 1999 (Memorie della Società Entomologica Italiana, 77: 317-357): S
<i>Strongylognathus huberi</i> Forel, 1874	-	-	+	-	-	-		
<i>Strongylognathus italicus</i> Finzi, 1924	s	-	-	+	-	-		
<i>Strongylognathus pisarskii</i> Poldi, 1994	e	-	-	+	-	-		
<i>Strongylognathus testaceus</i> (Schenck, 1852)	-	-	+	+	+	+		Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348): Si
<i>Strumigenys argiola</i> (Emery, 1869)	-	-	+	+	+	+	Baroni Urbani & De Andrade 2007 (Annali del Museo Civico di Storia Naturale "G. Doria", 99: 1-191): combination in <i>Strumigenys</i>	
<i>Strumigenys baudueri</i> (Emery, 1875)	-	-	+	+	+	+	Baroni Urbani & De Andrade 2007 (Annali del Museo Civico di Storia Naturale "G. Doria", 99: 1-191): combination in <i>Strumigenys</i>	
<i>Strumigenys membranifera</i> Emery, 1869	-	a	-	+	+	+	Baroni Urbani & De Andrade 2007 (Annali del Museo Civico di Storia Naturale "G. Doria", 99: 1-191): combination in <i>Strumigenys</i>	
<i>Strumigenys tenuipilis</i> Emery, 1915	-	-	+	+	+	+	Baroni Urbani & De Andrade 2007 (Annali del Museo Civico di Storia Naturale "G. Doria", 99: 1-191): combination in <i>Strumigenys</i>	Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348): Si
<i>Temnothorax affinis</i> Mayr, 1855	-	-	+	+	+	+		Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348): Si
<i>Temnothorax albipennis</i> (Curtis, 1854)	-	-	+	+	-	-	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i>	Glaser et al. 2012 (Gredleriana, 12: 273-284): N; Giannetti et al. 2019 (Insects, 10: 392): S.
<i>Temnothorax alienus</i> Schultz, Heinze & Pusch, 2007	e	-	-	+	-	-		Schultz et al. 2007 (Zootaxa, 1471: 1-14): S (description)
<i>Temnothorax angustulus</i> (Nylander, 1856)	-	-	+	+	-	-	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i>	Galkowski & Cagniant 2017 (Revue de l'Association Roussillonnaise

	endemic	alien	Continental Italy	Peninsular Italy	Sicilia	Sardegna	taxonomic remarks	occurrence remarks
								d'Entomologie, 26: 180-191): absence from Si
<i>Temnothorax aveli</i> (Bondroit, 1918)	-	-	+	+	-	-	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i>	Menozzi 1925 (Atti della Società dei Naturalisti e Matematici di Modena, 55: 22-4): N; Centorame et al. 2017 (Redia, 100: 89-94): S
<i>Temnothorax clypeatus</i> (Mayr, 1853)	-	-	+	+	-	-	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i>	
<i>Temnothorax corsicus</i> (Emery, 1895)	-	-	-	+	-	-	Ward et al. 2015 (Systematic Entomology, 40: 61-81): combination in <i>Temnothorax</i>	
<i>Temnothorax corticalis</i> (Schenck, 1852)	-	-	+	-	-	-	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i>	
<i>Temnothorax crassispinus</i> (Karavaiev, 1926)	-	-	+	-	-	-		Seifert & Csösz 2015 (Zookeys, 479: 37-64): N
<i>Temnothorax exilis</i> Emery, 1869	-	-	+	+	+	+	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i> ; Salata et al. 2019: senior synonym of <i>T. specularis</i>	
<i>Temnothorax finzii</i> Menozzi, 1924	-	-	+	-	-	-	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i>	Bračko et al. 2014 (Entomologica romanica, 16: 54): not an Italian endemic, also present in the Balkans
<i>Temnothorax flavicornis</i> Emery, 1870	-	-	+	+	+	-	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i>	Buschinger et al. 1988 (Zoologische Jahrbücher, 115: 383-401): Si
<i>Temnothorax gordigiani</i> (Ruzsky, 1902)	-	-	+	-	-	-		Finzi 1924 (Bollettino della Società Entomologica Italiana, 56: 120-123): N
<i>Temnothorax graecus</i> (Forel, 1911)	-	-	+	-	-	-		Müller 1921 (Bollettino della Società adriatica di scienze naturali in Trieste, 27: 46-49): N
<i>Temnothorax interruptus</i> (Schenck, 1852)	-	-	+	-	-	-	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i>	The previous edition of Faunatalia (Poldi et al. 1995) considered this species to be present in Si but no data has ever been published to support this claim
<i>Temnothorax kraussei</i> Emery, 1915	-	-	+	+	-	+	Ward et al. 2015 (Systematic Entomology, 40: 61-81): combination in <i>Temnothorax</i>	
<i>Temnothorax laestrygon</i> Santschi, 1931	e	-	-	-	+	-	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i>	
<i>Temnothorax lagrecai</i> Baroni Urbani, 1964	e	-	-	-	+	-	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i>	
<i>Temnothorax leviceps</i> (Emery, 1898)	-	-	-	+	-	-	Bolton 1995 (Cambridge, Mass.: Harvard University Press, 504 pp.): species status; Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i>	
<i>Temnothorax lichtensteini</i> Bondroit, 1918	-	-	+	+	+	+	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i>	Rigato & Toni 2011 (Research in the framework of the ICP Forests network. Conservazione Habitat Invertebrati, 5: 873-882): Sa; Schär et al. 2020 (Organisms Diversity and Evolution, 20: 405-416): Si
<i>Temnothorax luteus</i> Forel, 1874	-	-	+	+	+	-	Seifert et al. 2014 (Beiträge zur Entomologie, 64: 47-57): senior synonym of <i>T. tristis</i>	
<i>Temnothorax mediterraneus</i> Ward, Brady, Fisher & Schultz, 2015	s	-	-	-	+	+	Ward et al. 2015 (Systematic Entomology, 40: 61-81): replacement name for <i>T. kraussei</i>	
<i>Temnothorax minozzii</i> Santschi, 1932	e	-	+	-	-	-	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i>	
<i>Temnothorax muellerianus</i> Kutter, 1950	-	-	+	+	+	-	Buschinger et al. 1988 (Zoologische Jahrbücher, 115: 383-401): senior synonym of <i>T. insubricus</i> , <i>T. siciliensis</i> ; Ward et al. 2015 (Systematic Entomology, 40: 61-81): combination in <i>Temnothorax</i>	
<i>Temnothorax niger</i> Forel, 1894	-	-	+	-	-	-	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i>	
<i>Temnothorax nigriceps</i> (Mayr, 1855)	-	-	+	+	-	-		
<i>Temnothorax nylanderi</i> (Foerster, 1850)	-	-	+	+	-	+	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i>	
<i>Temnothorax parvulus</i> (Schenck, 1852)	-	-	+	+	-	+	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i>	

	endemic	alien	Continental Italy	Peninsular Italy	Sicilia	Sardegna	taxonomic remarks	occurrence remarks
<i>Temnothorax racovitzai</i> (Bondroit, 1918)	-	-	+	+	-	-	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i> ; Casevitz-Weulersse & Galkowski, 2009: senior synonym of <i>T. massiliensis</i>	
<i>Temnothorax ravouxi</i> (André, 1861)	-	-	+	+	+	+	Ward et al. 2015 (Systematic Entomology, 40: 61-81): combination in <i>Temnothorax</i>	
<i>Temnothorax recedens</i> (Nylander, 1856)	-	-	+	+	+	+		
<i>Temnothorax rogeri</i> Emery, 1869	-	-	+	-	-	-		
<i>Temnothorax rottenbergii</i> (Emery, 1870)	-	-	+	+	+	-	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i>	
<i>Temnothorax sardous</i> Santschi, 1909	e	-	-	-	-	+	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i>	
<i>Temnothorax saxatilis</i> Schultz, Heinze & Pusch, 2007	e	-	-	+	-	-		Schultz et al. 2007 (Zootaxa, 1471: 1-14): S (description)
<i>Temnothorax sordidulus</i> (Müller, 1923)	-	-	+	-	-	-	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i> ; Schulz, 1991 (Zeitschrift der Arbeitsgemeinschaft Österreichischer Entomologen, 43:120-122): senior synonym of <i>T. carinthiacus</i>	
<i>Temnothorax stumpeli</i> Kutter, 1950	-	-	+	-	-	-	Ward et al. 2015 (Systematic Entomology, 40: 61-81): combination in <i>Temnothorax</i>	
<i>Temnothorax tergestinus</i> (Finzi, 1928)	-	-	+	-	-	-	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i> ; Csósz et al. 2015 (PloS One: 10: e0140000) : species status	
<i>Temnothorax tuberum</i> (Fabricius, 1775)	-	-	+	+	+	+	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i>	
<i>Temnothorax unifasciatus</i> (Latreille, 1798)	-	-	+	+	+	+	Bolton 2003 (Memoirs of the American Entomological Institute, 71: 1-370): combination in <i>Temnothorax</i>	Buschinger et al. 1988 (Zoologische Jahrbücher, 115: 383-401): Si
<i>Tetramorium alpestre</i> Steiner, Schlick-Steiner & Seifert, 2010	-	-	+	+	+	-		Steiner et al. 2010 (Zoologischer Anzeiger, 249: 223-254): N (description); Wagner et al. 2017 (Myrmecological News, 25: 95-129): S, Si
<i>Tetramorium alternans</i> Santschi, 1929	-	-	-	-	+	-	Csósz et al. 2007: senior synonym of <i>T. biskrense kahenae</i>	
<i>Tetramorium atratum</i> (Schenck, 1852)	-	-	+	-	+	-	Ward et al. 2015 (Systematic Entomology, 40: 61-81): combination in <i>Tetramorium</i>	Sanetra et al. 1999 (Memorie della Società Entomologica Italiana, 77: 317-357): S. Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348): Si
<i>Tetramorium bicarinatum</i> (Nylander, 1846)	-	a	+	+	+	-		Limonta & Colombo 2003 (Bollettino di Zoologia agraria e di Bachicoltura, 35: 287-289): N; Borowiec & Salata 2015 (Sociobiology, 62: 181-186): S; Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348): Si
<i>Tetramorium brevicorne</i> Bondroit, 1918	-	-	-	-	-	+		
<i>Tetramorium caespitum</i> (Linnaeus, 1758)	-	-	+	+	-	+		Wagner et al. 2017 (Myrmecological News, 25: 95-129): N, S, Sa
<i>Tetramorium diomedea</i> Emery, 1908	-	-	-	+	+	-		
<i>Tetramorium fusciclava</i> Consani & Zangheri, 1952	e	-	+	+	-	-	Wagner et al. 2017 (Myrmecological News, 25: 95-129): species level	Wagner et al. 2017 (Myrmecological News, 25: 95-129): N, S
<i>Tetramorium immigrans</i> Santschi, 1927	-	a	+	+	+	+	Wagner et al. 2017 (Myrmecological News, 25: 95-129): species level	Wagner et al. 2017 (Myrmecological News, 25: 95-129): N, S, Sa; Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348): Si
<i>Tetramorium impurum</i> (Foerster, 1850)	-	-	+	+	-	-		Wagner et al. 2017 (Myrmecological News, 25: 95-129): N, S
<i>Tetramorium indocile</i> Santschi, 1927	-	-	+	+	-	-	Csósz et al. 2014 (Zoologischer Anzeiger, 253: 469-481): species level	Wagner et al. 2017 (Myrmecological News, 25: 95-129): N, S;
<i>Tetramorium lanuginosum</i> Mayr, 1870	-	a	-	-	+	-		Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348): Si
<i>Tetramorium lucayanum</i> Wheeler, 1905	-	a	+	-	-	-		Jucker et al. 2008 (Bollettino di Zoologia agraria e di Bachicoltura, 40: 99-107): N

	endemic	alien	Continental Italy	Peninsular Italy	Sicilia	Sardegna	taxonomic remarks	occurrence remarks
<i>Tetramorium meridionale</i> Emery, 1870	-	-	+	+	-	+		
<i>Tetramorium moravicum</i> Novák & Sadil, 1941	-	-	+	+	-	-	Csősz et al. 2007 (Zootaxa, 1405: 1-38): N, S	
<i>Tetramorium punctatum</i> Santschi, 1927	-	-	-	+	+	-	Sanetra et al. 1999 (Memorie della Società Entomologica Italiana, 77: 317-357): species level	
<i>Tetramorium sanetrai</i> Csősz et al. 2007	e	-	-	+	-	-		Csősz et al. 2007 (Zootaxa, 1405: 1-38): S (description)
<i>Tetramorium semilaeve italicum</i> Menozzi, 1932	e	-	-	+	-	-	Borowiec et al. 2015 (Zookeys, 512: 39-62): dubious status	
<i>Tetramorium semilaeve semilaeve</i> André, 1883	-	-	+	+	+	+	Sanetra et al. 1999 (Memorie della Società Entomologica Italiana, 77: 317-357): senior synonym of <i>T. semilaeve liparaeum</i> , <i>T. semilaeve siciliense</i>	
Ponerinae								
<i>Cryptopone ochracea ochracea</i> (Mayr, 1855)	-	-	+	+	+	-		
<i>Cryptopone ochracea sicula</i> (Emery, 1909)	e	-	-	-	+	-		
<i>Hypoponera abeillei</i> (André, 1881)	-	-	+	+	+	-		Mei 1995 (Il Naturalista Siciliano, 19: 753-772): Si
<i>Hypoponera eduardi</i> (Forel, 1894)	-	-	+	+	+	+		Rigato & Toni 2011 (Research in the framework of the ICP Forests network. Conservazione Habitat Invertebrati, 5: 873-882): Sa
<i>Hypoponera punctatissima</i> (Roger, 1859)	-	a	+	-	+	+	Bolton & Fisher 2011 (Zootaxa, 2843: 1-118): senior synonym of <i>Ponera sulcitana</i>	Bolton & Fisher 2011 (Zootaxa, 2843: 1-118): Si
<i>Hypoponera ragusai</i> (Emery, 1894)	-	-	-	+	+	-		Bolton & Fisher 2011 (Zootaxa, 2843: 1-118): S
<i>Ponera coarctata</i> (Latreille, 1802)	-	-	+	+	+	+		
<i>Ponera testacea</i> Emery, 1895	-	-	+	+	+	+	Csősz & Seifert 2003 (Acta Zoologica Academiae Scientiarum Hungaricae, 49: 201-214): species status	
Proceratiinae								
<i>Proceratium algiricum</i> Forel, 1899	-	-	-	+	+	-		
<i>Proceratium melinum</i> (Roger, 1860)	-	-	+	-	-	-		
<i>Proceratium melitense</i> de Andrade, 2003	s	-	-	-	+	-		Baroni Urbani & de Andrade 2003 (Museo Regionale di Scienze Naturali Monografie (Turin), 36: 1-492): Si (description)
<i>Proceratium numidicum</i> Santschi, 1912	-	-	-	-	+	-		Schifani & Alicata 2018 (Polish Journal of Entomology, 87: 323-348): Si

SUPPLEMENTARY FILE S2

Schifani E., 2021. Formicidae. In: Bologna M.A., Zapparoli M., Oliverio M., Minelli A., Bonato L., Cianferoni F., Stoch F. (eds.), Checklist of the Italian fauna. Version 1.0. Last update: 2021-05-31

Subfamily	Genus	Author and year of the genus	Subgenus	Author and year of the subgenus	Species	Author and year of the species	Subspecies	Author and year of the subspecies	Reason for the exclusion
Formicinae	<i>Formica</i>	Linnaeus, 1758	Serviformica	Forel, 1913	<i>pyrenaea</i>	Bondroit, 1918			The previous edition of Faunalia (Poldi et al. 1995) considered this species to be present in Italy (N) but no data has ever been published to sustain this claim
Formicinae	<i>Formica</i>	Linnaeus, 1758	Serviformica	Forel, 1913	<i>transcaucasica</i>	Nassonow, 1899			Seifert 2004 (Myrmecologische Nachrichten, 6: 29-38): <i>incertae sedis</i>
Myrmicinae	<i>Messor</i>	Forel, 1890			<i>sanctus</i>	Emery, 1921	<i>obscuriventris</i>	Karawajew, 1912	Bolton 1995 (Cambridge, Mass.: Harvard University Press, 504 pp.): unavailable name
Myrmicinae	<i>Solenopsis</i>	Westwood, 1840			<i>orbula</i>	Emery, 1875	<i>terniensis</i>	Forel, 1905	Mei 1995 (Il Naturalista Siciliano, 19: 753-772) considers the historic records of this taxon doubtful
Myrmicinae	<i>Cardiocondyla</i>	Emery, 1869			<i>nuda</i>	(Mayr, 1866)			Seifert et al. 2017 (Zootaxa, 4290: 324-356): absent from the Mediterranean region and Europe
Myrmicinae	<i>Pheidole</i>	Westwood, 1839			<i>sinaitica</i>	Mayr, 1862			Mei 1995 (Il Naturalista Siciliano, 19: 753-772): absence from Lampedusa where it had been previously recorded
Dolichoderinae	<i>Bothriomyrmex</i>	Emery, 1869			<i>siculus</i>	Emery, 1925			Shattuck 1994 (University of California Press): unavailable name
Dolichoderinae	<i>Bothriomyrmex</i>	Emery, 1869			<i>costae</i>	Emery, 1869			Seifert 2012 (Myrmecological News, 17: 91-104): <i>incertae sedis</i>
Dolichoderinae	<i>Tapinoma</i>	Foerster, 1850			<i>nigerrimum</i>	Nylander, 1856			Seifert et al. 2017 (Myrmecological News, 24: 123-144): absent from Italy