

Mitochondrial DNA Part B Resources

ISSN: (Print) (Online) Journal homepage: <https://www.tandfonline.com/loi/tmdn20>

Complete mitochondrial genome of the H3 haplotype Argentine ant *Linepithema humile* (Mayr, 1868) (Formicidae; Hymenoptera)

Jonghyun Park, Chan-Ho Park & Jongsun Park

To cite this article: Jonghyun Park, Chan-Ho Park & Jongsun Park (2021) Complete mitochondrial genome of the H3 haplotype Argentine ant *Linepithema humile* (Mayr, 1868) (Formicidae; Hymenoptera), Mitochondrial DNA Part B, 6:3, 786-788, DOI: [10.1080/23802359.2021.1882900](https://doi.org/10.1080/23802359.2021.1882900)

To link to this article: <https://doi.org/10.1080/23802359.2021.1882900>



© 2021 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 11 Mar 2021.



Submit your article to this journal [↗](#)



Article views: 22



View related articles [↗](#)



View Crossmark data [↗](#)

MITOGENOME ANNOUNCEMENT



Complete mitochondrial genome of the H3 haplotype Argentine ant *Linepithema humile* (Mayr, 1868) (Formicidae; Hymenoptera)

Jonghyun Park^{a,b}, Chan-Ho Park^c  and Jongsun Park^{a,b} 

^aInfoBoss Inc, Seoul, Republic of Korea; ^bInfoBoss Research Center, Seoul, Republic of Korea; ^cNational Institute of Biological Resources, Incheon, Korea

ABSTRACT

The Argentine ant, *Linepithema humile* (Mayr, 1867), is an invasive ant species that has spread across the world. We have determined the mitochondrial genome of *L. humile* collected in South Korea, which is 15,934 bp containing 10 SNPs and 5 INDELs compared to the previous mitogenome. Most SNPs were found in *cox3*, followed by *cytb*. From SNPs our mitogenome was identified as a H3 haplotype, which was previously recorded in Japan and the U.S. while the previous mitogenome was H1 haplotype. Phylogenetic analysis was congruent to previous study within the tribe Leptomyrmecini but not between other tribes of subfamily Dolichoderinae.

ARTICLE HISTORY

Received 19 October 2020
Accepted 24 January 2021

KEYWORDS

Linepithema humile;
mitochondrial genome; H3
haplotype;
Formicidae; Korea

Linepithema humile, belonging to Formicidae family, more often called 'Argentine ants' is an infamous tramp ant species named after its neotropical origin. They have a unique feature of 'unicolonality' where multiple colonies cooperate forming so called supercolonies (Sunamura et al. 2009). Mating occurs within colonies and new colonies are made when small fractions of the mother colony bud out forming smaller colonies (Keller and Passera 1992). While these supercolonies remained insignificant in native ranges, genetic bottleneck of initial introduction and following 'genetic cleansing' presumably by annual queen executions (Keller et al. 1989; Inoue et al. 2015) caused extreme relatedness within introduced populations which allowed massive supercolonies distributed over multiple continents (Tsutsui et al. 2003). In Korea, Argentine ants have just arrived in southern ports of Busan and Kwangyang (Lee et al. 2020). Securing the mitogenome of the newly introduced populations will not only increase our understandings of its genetic background but aid the identification and managements of invasive species (Park et al. 2020).

Worker ants of *L. humile* were collected in Busan station, Busan, Republic of Korea (35°07'25.8"N 129°02'55.0"E). The ants were yellowish brown, about 2.5 mm from tip to tip, and were covered in light pubescence. They were easily identified as *L. humile* since no other dolichoderine ants in Korea had such large eyes located close to the anterior margin, elongated body structure with the propodeum much lower compared to the pronotum and mesonotum (Dong 2017). Total DNA was extracted using DNeasy Blood & Tissue Kit (QIAGEN, Hilden, Germany). Raw sequences obtained from Illumina NovaSeq6000 (350-bp paired-end library) at Macrogen Inc.,

Korea, were filtered by Trimmomatic v0.33 (Bolger et al. 2014), *de novo* assembled and confirmed by Velvet v1.2.10 (Zerbino and Birney 2008), SOAPGapCloser v1.12 (Zhao QY et al. 2011), BWA 0.7.17 (Li et al. 2009), and SAMtools v1.9 (Song and Liang 2013) under the environment of Genome Information System (GeIS; <http://geis.infoboss.co.kr>). Geneious R11 v11.1.5 (Biomatters Ltd, Auckland, New Zealand) was used for annotations based on the mitogenome (NC_045057; Zhao E et al. 2017). DNA sample and specimen (95% ethanol) were deposited in InfoBoss Cyber Herbarium (IN; <http://herbarium.infoboss.co.kr>; Contact: Suhyeon Park, shpark817@infoboss.co.kr; KFDS00420).

The new mitogenome of *L. humile* (GenBank accession is MT890564) is 15,934 bp long, 5 bp longer than the previous mitogenome due to 5 insertion and deletions (INDELs): three in control region and one each in *tRNA-Ile* and *tRNA-Cys*. Gene components and order of 37 genes, 13 protein-coding genes (PCGs), 2 rRNAs, and 22 tRNAs, were conserved compared to other dolichoderine mitogenomes. Ten single nucleotide polymorphisms (SNPs) were identified between the two mitogenomes, all in PCGs: five in *cox3*, 2 in *cytb*, and 1 in *cox1*, *cox2*, and *nad4*.

To investigate the genetic properties of Argentine ants supercolonies, fractions of *cox1*, *cox2*, and *cytb* genes were developed as mitochondrial markers suggesting 19 different haplotypes in both native and introduced ranges (Vogel et al. 2009, 2010; Inoue et al. 2013). Our mitogenome was identical to the H3 haplotype while the previous mitogenome was H1 haplotype (LH3 and LH1 in Inoue et al. 2013). H1 was the intercontinental supercolony's haplotype while H3 was a minority found in Bermuda, Chile, Ecuador, North Carolina

CONTACT Jongsun Park  starflr@infoboss.co.kr  InfoBoss Inc, Seoul, Republic of Korea

© 2021 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. (A) Photo of *Linepithema humile* workers foraging in the collected site of Busan, South Korea. (B) Lateral view of *Linepithema humile* collected in Busan, South Korea. (C) Bayesian inference (1,100,000 generations) phylogenetic tree of 13 ant mitochondrial genomes. The numbers above branches indicate the posterior probabilities of the Bayesian inference tree.

and the lesser supercolonies of California and Kobe (Vogel et al. 2010; Inoue et al. 2013). The last two countries, Japan and the U.S., could be suspected as the origin of Korean population as they are the third and second largest countries in imports of Korea (Choi and Choi 2017).

Thirteen PCGs from 13 ants consisting of 7 dolichoderine ants, representative ants to each subfamily, and an outgroup species, *Apis mellifera ligustica*, were aligned independently using MAFFT v7.450 (Katoh and Standley 2013) and were concatenated for phylogenetic purposes. Bayesian inference tree was generated using MrBayes v3.2.6 (Ronquist et al. 2012). In the phylogenetic tree, two *L. humile* was grouped with *Ochetellus glaber* and two remaining leptomyrmecine species were also grouped (Figure 1), congruent to the previous phylogeny of the tribe (Ward et al. 2010). Relationship between the tribes, however, was incongruent to the previous study as tribe Leptomyrmecini was grouped with tribe Tapinomini not Dolichoderini based on nuclear genes (Ward et al. 2010). In addition, the two nodes showing low support values (Figure 1) are also congruent to the previous studies (Park et al. 2020c, 2020a, 2020b), requiring additional mitogenomes to improve them.

Our findings show only two of ten SNPs were used as haplotype markers, suggesting further utilization on other regions of

mitogenome will be useful. The highly variable *cox3* could be utilized for further dividing haplotypes, which could separate supercolonies distinguishable in behavioral test but not in haplotypes (e.g., the minor supercolonies of California). It also shows that the upstream region amplified by LCO1490-HCO2198 universal marker (Folmer et al. 1994) can be considered additional marker region as the only SNP was found on *cox1*.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

This work was supported by both InfoBoss Research Grant (IBG-0017) and Cooperative Research Program for Agriculture Science & Technology Development [Project No. PJ013389052020] Rural Development Administration, Republic of Korea.

ORCID

Chan-Ho Park  <http://orcid.org/0000-0003-4047-1918>
Jongsun Park  <http://orcid.org/0000-0003-0786-4701>

Data availability statement

Mitochondrial genome sequence can be accessed via accession number MT890564 in GenBank of NCBI at <https://www.ncbi.nlm.nih.gov>. The associated BioProject, SRA, and Bio-Sample numbers are PRJNA669488, SAMN16446430, and SRR12834976, respectively.

References

- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*. 30(15):2114–2120.
- Choi S-H, Choi J-I. 2017. Analysis of volatility and directionality of Korean imports and exports: focused on USA, Japan, China, UK. *J Digit Convergence*. 15(10):113–121.
- Dong M. 2017. The ants of Korea. Seoul: Nature and Ecology.
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R. 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol Mar Biol Biotechnol*. 3(5): 294–299.
- Inoue MN, Ito F, Goka K. 2015. Queen execution increases relatedness among workers of the invasive Argentine ant, *Linepithema humile*. *Ecol Evol*. 5(18):4098–4107.
- Inoue MN, Sunamura E, Suhr EL, Ito F, Tatsuki S, Goka K. 2013. Recent range expansion of the Argentine ant in Japan. *Diversity Distrib*. 19(1): 29–37.
- Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol*. 30(4):772–780.
- Keller L, Passera L. 1992. Mating system, optimal number of matings, and sperm transfer in the Argentine ant *Iridomyrmex humilis*. *Behav Ecol Sociobiol*. 31(5):359–366.
- Keller L, Passera L, Suzzoni JP. 1989. Queen execution in the Argentine ant, *Iridomyrmex humilis*. *Physiol Entomol*. 14(2):157–163.
- Lee HS, Kim DE, Lyu DP. 2020. Discovery of the invasive Argentine ant, *Linepithema humile* (Mayr)(Hymenoptera: Formicidae: Dolichoderinae) in Korea. *Kor J Appl Entomol*. 59(1):71–72.
- Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R. 2009. The sequence alignment/map format and SAMtools. *Bioinformatics*. 25(16):2078–2079.
- Park J, Xi H, Kim D, Park J, Lee W, Lee K. 2020. Current status of mitochondrial gene sequences of invasive and exotic pests: directions for managing the integrated platform for invasive pests (IPIP). 2020 Spring International Conference of KSAE:74.
- Park J, Xi H, Park J. 2020a. The complete mitochondrial genome of *Aphaenogaster famelica* (Smith, 1874) (Hymenoptera: Formicidae). *Mitochondrial DNA Part B*. 5(1):492–494.
- Park J, Xi H, Park J. 2020b. The complete mitochondrial genome of *Nylanderia flavipes* (Smith, 1874) (Hymenoptera: Formicidae). *Mitochondrial DNA Part B*. 5(1):420–421.
- Park J, Xi H, Park J. 2020c. The complete mitochondrial genome of *Ochetellus glaber* (Mayr, 1862) (Hymenoptera: Formicidae). *Mitochondrial DNA Part B*. 5(1):147–149.
- Ronquist F, Teslenko M, Van Der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP. 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Syst Biol*. 61(3):539–542.
- Song N, Liang A-P. 2013. A preliminary molecular phylogeny of planthoppers (Hemiptera: Fulgoroidea) based on nuclear and mitochondrial DNA sequences. *PLOS One*. 8(3):e58400
- Sunamura E, Espadaler X, Sakamoto H, Suzuki S, Terayama M, Tatsuki S. 2009. Intercontinental union of Argentine ants: behavioral relationships among introduced populations in Europe, North America, and Asia. *Insect Soc*. 56(2):143–147.
- Tsutsui ND, Suarez AV, Grosberg RK. 2003. Genetic diversity, asymmetrical aggression, and recognition in a widespread invasive species. *Proc Natl Acad Sci*. 100(3):1078–1083.
- Vogel V, Pedersen JS, d'Ettorre P, Lehmann L, Keller L. 2009. Dynamics and genetic structure of Argentine ant supercolonies in their native range. *Evolution*. 63(6):1627–1639.
- Vogel V, Pedersen JS, Giraud T, Krieger MJ, Keller L. 2010. The worldwide expansion of the Argentine ant. *Divers Distrib*. 16(1):170–186.
- Ward PS, Brady SG, Fisher BL, Schultz TR. 2010. Phylogeny and biogeography of dolichoderine ants: effects of data partitioning and relict taxa on historical inference. *Syst Biol*. 59(3):342–362.
- Zerbino DR, Birney E. 2008. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. *Genome Res*. 18(5):821–829.
- Zhao E, Bi G, Yang J, Zhang Z, Liu G, Du Q, Shang E. 2017. Complete mitochondrial genome of the Argentine ant, *Linepithema humile* (Hymenoptera: Formicidae). *Mitochondrial DNA Part A*. 28(2):210–211.
- Zhao QY, Wang Y, Kong YM, Luo D, Li X, Hao P. 2011. Optimizing *de novo* transcriptome assembly from short-read RNA-Seq data: a comparative study. *BMC Bioinf*. 12(Suppl 14):S2. 2011: BioMed Central.