

Biogeography and ecology of the ant fauna of Madagascar (Hymenoptera: Formicidae)

B. L. FISHER

*Department of Entomology, University of California, Davis,
CA 95616, USA*

(Accepted 1 May 1996)

In the Malagasy region, 52 ant genera have been recorded, of which 48 are estimated to be indigenous. On the island of Madagascar, 41 out of 45 recorded genera are considered to be indigenous. Four genera are endemic to Madagascar and one to Mauritius. Currently, there are 383 names of described species-group taxa from the Malagasy region, and 320 from Madagascar. The level of endemism is high at the species level, >90%. The ant fauna of the Malagasy region, however, is poorly understood; an estimated two thirds of the ant species are undescribed. The diversity of ant genera on the island is comparable to that of other biogeographic regions. On the basis of generic and species level comparisons, the Malagasy fauna shows greater affinities to the Afrotropical region than to peninsular India and the Oriental region. The distinct gaps in the taxonomic and ecological composition of the fauna of Madagascar are evaluated in the context of island radiations and relict taxa. The absence of driver ants in Madagascar may have spurred the diversification of Cerapachyinae and may have permitted the persistence of other relict taxa such as the Amblyoponini.

KEYWORDS: Formicidae, biogeography, Madagascar, systematics, Africa, ecology.

Introduction

For the Malagasy region, ant taxonomy had a prolific beginning in the late 1800's and early 1900's. In 1891, and with an additional supplement in 1892, Forel provided the first and only thorough treatment of the ants of the Malagasy region. On the basis of this work, Emery (1920c) evaluated the similarities of the ant genera of Madagascar with Asia and Africa. A few years later, Wheeler (1922c) provided an updated list of the ant fauna of the region. Except for Donisthorpe's work based on the collections of Mamet and Vinson in the Mascarenes (summarized in Mamet 1954), almost 50 years passed without additional work on the ants of Madagascar (Fig. 1). One reason for the lack of taxonomic interest in Madagascar was the paucity of new material since the late 19th and early 20th century collections of Camboué, Sikora, Mocquerys, Voeltzkow, Perrot, Alluaud, and Hildebrandt.

In the late 1960's, Brown provided greatly needed new collections from Madagascar, and returned our attention to the importance of the Malagasy region in understanding ant phylogeny (Brown 1975, 1978a; and see Bolton 1979, 1981). Over the last five years, collections and field observations from Madagascar have greatly increased, improving our understanding of the evolution, taxonomy, and ecology of




FIG. 1. Date of publication of all named ants from the Malagasy region. The number of described taxa are grouped by five year intervals.

ants of the island, and most of all, have demonstrated that the fauna is incompletely known (Alpert 1992, Ward 1994, Fisher in press, Olson and Ward in press).

The time is right to reassess what we know of the ants of Madagascar, to provide baseline taxonomic data for the resurgence of interest in the region's ant fauna, and to provide a summary of the biogeographic patterns ascertainable from available data. To this end, I add to the early work of Forel, Emery, and Wheeler, and provide an updated species list for the Malagasy region and discuss the generic affinities of the fauna to the Afrotropical and Oriental regions. I propose possible origins for the Malagasy fauna and evaluate the striking gaps in the taxonomic composition of the fauna in the context of island radiations.

Results and discussion

Ant species of the Malagasy region

Data on the richness and endemism of Malagasy ants are summarized in Appendix 1 which contains a list of all valid species-group names. In all discussions and analyses, Madagascar includes the coastal islands (e.g., Nossi Bé, St. Marie), and the Malagasy Region refers to Madagascar and neighbouring islands of the Indian Ocean: Mauritius, Réunion, Seychelles, Rodrigues, Aldabra, Farquhar, Chagos, and the Comoro Is.

In 1893, Dalla Torre listed 119 Malagasy species (excluding subspecies), and

Table 1. The number of valid specific and subspecific names, and the number of introduced and endemic taxa. Introduced and endemic taxa contain both species and subspecies. Data summarized from the list of ant species-group names of the Malagasy Region in Appendix 1.

	Madagascar	Malagasy Region
Valid specific and subspecific names	320	383
Valid subspecific names	76	93
Introduced taxa	19	33
Endemic taxa	271	331
Per cent endemism	90%	95%

Wheeler (1922c) enumerated 237 species (excluding subspecies) for the Malagasy Region. The present total of 290 species (383 including subspecies) for the Malagasy region is a small increase in described taxa over the last 75 years (Appendix 1, Fig. 1). Recent collections by the author, G. D. Alpert *et al.*, D. M. Olson, and P. S. Ward suggest that there may be over 1000 species in the Malagasy region. The recently collected, undescribed material, indicates that the level of endemism for Madagascar and the Malagasy region is greater than the figures of 90% and 95% presented in Table 1. New and more intense inventory techniques are collecting an amazing number of new species that are cryptic and locally distributed in Madagascar. A recent ant survey of a wet tropical closed forest site in the RNI d'Andringitra found an estimated 100% of the 134 ant species in this mountainous region to be endemic to Madagascar (Fisher, in press).

Biogeographical affinities

Griswold (1991) described two main approaches to understanding historical biogeographic patterns, the phenetic and cladistic methods. The cladistic approach determines the relationship among areas based on sister-group relationships among taxa endemic to those areas. The phenetic approach does not use a phylogeny to determine relationships among areas, but evaluates relationships based on proportion of shared taxa. Though Griswold clearly demonstrates that the cladistic method is the preferred approach, most invertebrate groups lack the necessary phylogenetic information to construct the area cladograms necessary to implement these methods. In these cases, the phenetic approach is appropriate, but with the understanding that historical patterns may need future testing with cladistic methods. For the Malagasy ant fauna, the phenetic comparison must be done at the generic level, since the species level taxonomy is inadequate.

In this paper, the origins and affinities of the ants of Madagascar are investigated by comparing the number of genera in common with neighbouring land masses of India and Africa which were all once united in Gondwana (Briggs, 1987). The ant genera present in Madagascar and other biogeographic regions are presented in Appendix 2. In addition to described taxa and literature records, the list contains data from undescribed material in museum collections.

To further understand affinities with the African ant fauna, I have compared the genera of Madagascar with geographically distinct regions within Africa: West Africa (W.A.), East Africa (E.A.), and southern Africa (S.A.) (see Appendix 2 for area descriptions). Ideally these regions would be equal in size to Madagascar, and contain equal representation of the habitats found on the island. Unfortunately, we lack detailed information on the distribution of the ant genera in Africa. For example, there are insufficient collections records to develop a list of the ant genera of Mozambique, the land mass closest to Madagascar. I am not proposing that these areas of Africa are biogeographically distinct. The regions differ in the proximity to Madagascar with South and East Africa providing closer grounds for dispersal to Madagascar. This division also separates the mostly mesic tropical forest of West Africa from the more xeric habitats of East and southern Africa.

There are 52 ant genera, 48 of which are estimated to be indigenous, in the Malagasy region. The classification of a genus as indigenous or introduced is based on a subjective evaluation of historical distribution records and the biology of the species. There are five endemic genera in the region: *Adetomyrma* [Mad.], *Eutetramorium* [Mad.], *Ireneopone* [Mau.], *Pilotrochus* [Mad.], and an undescribed genus [Mad.].

Is the richness of 48 genera in Madagascar depauperate compared with other




FIG. 2. The log number of genera for each Gondwanaland biogeographic region plotted with respect to the log area of the region. The data fit the equation: $f(x) = 0.215(x) + 0.52$ with a $R^2 = 0.60$. If the subregions EA, SA, WA and IND are removed $R^2 = 0.553$. Biogeographic regions: Neotropical (NEO), Afrotropical (AF), Malagasy (MAL), Palaeartic (PAL), Oriental (ORI), Indo-Australian (INA), Australian (AUS), India (IND), West Africa (WA), East Africa, (EA), southern Africa (SA). Regions are defined in Appendix 2.

tropical regions? Ideally, a comparison of diversity would be made at the species level, but these data do not exist. In Fig. 2, I present a genera-area curve for the Gondwanaland regions and subregions. The log of the total number of genera present in each of the regions is plotted in relation to the log of the area of each region. The slope of the log-log relationship has a z-value of 0.215 ($R^2 = 0.60$). The log number of genera in the Malagasy region falls very close to this line. Thus, the ant fauna, at least at the generic level, is not depauperate. The Indo-Australian region is the only visible outlier, showing a highly diverse fauna in a small area. This could be the result of an actual higher generic diversity, or an artifact of a poorly defined biogeographic region. As defined here in Bolton (1995b), the Indo-Australian region includes islands on both sides of the Wallace line. Future work should be directed at clarifying the boundaries of this region.

These genera are shared between the Malagasy region and India but are not found in Africa: *Aphaenogaster*, *Kyidris*, and *Vollenhovia*. These three genera are found in other biogeographic regions and are not uniquely shared between the Malagasy region and India (Appendix 2).

Nine genera are shared between the Malagasy region and Africa and are not recorded from India: *Anoplolepis* [All Africa], *Discothyrea* [S.A., W.A.], *Glomyrmex* [W.A., E.A.], *Melissotarsus* [All Africa], *Mystrium* [W.A.], *Prionopelta* [S.A., W.A.], *Pristomyrmex* [All Africa], *Simopone* [S.A., W.A.], and *Terataner* [All Africa]. *Melissotarsus* and *Terataner* are the only genera uniquely shared between the Malagasy region and Africa. *Simopone*, *Mystrium*, *Prionopelta*, *Discothyrea*, and *Pristomyrmex* are also




FIG. 3. Dendrogram resulting from a single linkage cluster analysis of the genera present in each biogeographic region: Neotropical (NEO), Nearctic (NEA), Afrotropical (AFR), Malagasy (MAL), Palaeartic (PAL), Oriental (ORI), Indo-Australian (INA), and Australian (AUS). Data shown in Appendix 2.

Table 2. Number of genera shared and the Jaccard Index of similarity between biogeographic regions presented in Appendix 2. Higher values represent greater similarity. (Mal. = Malagasy region; Ori. = Oriental Region; Afr. = Afrotropical region; E.A. = East Africa; W.A. = West Africa; S.A. = southern Africa; see Appendix 2 for further explanation of regions.)

	Mal.-Ori.	Mal.-India	Mal.-Afr.	Mal.-W.A.	Mal.-E.A.	Mal.-S.A.
Genera shared	37	33	40	39	36	37
Unique to Mal.	11	15	8	11	15	11
Unique to other	60	41	49	42	19	23
Jaccard index	0.343	0.371	0.412	0.424	0.514	0.521

found in the Oriental biogeographic region and it is possible that these genera may have become locally extinct in India.

The number of uniquely shared genera is very low, but it does suggest that the ant fauna of Madagascar has closer affinities to Africa than to India and the Orient (two genera uniquely shared with Africa, zero with India and the Oriental region). This type of analysis would be more appropriate at the species level, but the poor state of the taxonomy of these regions does not permit it.

A cluster analysis (single linkage; SAS, 1994) of the genera present in each biogeographic region groups the Malagasy and African Regions as most similar (Fig. 3). An analysis of faunal similarity using the Jaccard Index (Magurran, 1988) also indicates that the Malagasy fauna is most similar to the African, and more precisely, to that of the southern and East Africa regions (Table 2; Jaccard Index: $C_j = j / (a + b - j)$ where j = number of genera shared by both regions, a = number of genera in Madagascar, and b = number of genera in region B).

An examination of the only recent world revision of a genus that included Madagascar (Bolton, 1979) indicates that the Madagascar-Africa relationship is also true at the level of species-groups and species. Of the 36 described species of *Tetramorium* in the Malagasy region, 29 are endemic, four are shared with Africa, and three are pantropical tramp species. The 36 described species represent eight species groups of which three are endemic to Madagascar, two are shared with Africa and three are widespread. In addition, three indigenous species from the Malagasy region are shared with Africa, while none are shared with the Oriental or Indo-Australian regions. Closer affinities with the Oriental and Indo-Australian regions, however, may exist in other pantropical genera, such as *Strumigenys* and *Pheidole* which have numerous undescribed species in collections awaiting revision.

Nineteen genera are found in both Africa and the Oriental region, but not in Madagascar: *Aenictus*, *Calyptomyrmex*, *Carebara*, *Cataglyphis*, *Centromyrmex*, *Cryptopone*, *Dorylus*, *Epitritus*, *Leptanilla*, *Messor*, *Myrmicaria*, *Oecophylla*, *Paedalgus*, *Pheidologeton*, *Polyrhachis*, *Probolomyrmex*, *Rhoptromyrmex*, *Pseudolasius*, and *Sphinctomyrmex*.

Nesting behaviour may help explain which ant genera successfully colonized the island. The following discussion assumes that the ant genera found on Madagascar arrived primarily by dispersal (see discussion of the origin of genera below). The nineteen general listed above nest either in the ground, with termite, in the leaf litter or in rotten wood, or in self constructed nests. *Polyrhachis* is the only genus that has some species that nest in plant cavities. Brown (1973) suggested that those ants that nest in performed plant cavities are best suited for surviving long distance dispersal by rafting across an ocean. Not surprisingly, however, there are other genera that successfully colonized Madagascar which do not nest in plant cavities. In addition to not nesting in plant cavities, the arrival of the driver ants (*Aenictus* and *Dorylus*) and *Leptanilla* to Madagascar has been hindered by the fact that queens are wingless. The genus, *Aenictus*, however, has managed to establish on the islands beyond the Asian mainland, crossing even the Wallace Line (Gotwald 1995).

The striking gaps in the taxonomic composition of the fauna of Madagascar suggest a disharmonic or unbalanced condition. The driver ants which dominate the forest floor, and the weaver ants (*Oecophylla*) which dominate the forest canopy in Afrotropical and Oriental tropical regions are absent from the Malagasy region. Weaver ants and especially driver ants are important predators of other ants and have been shown to influence ant population structure and the diversity of ant communities (Hölldobler and Wilson, 1990; Gotwald, 1995). Because of the absence of driver ants and weaver ants from Madagascar, the ant population dynamics of Malagasy ant communities may differ greatly from other Old World ant communities. During a raid, driver ants provide a natural disturbance in the ant community. They create a gap, like a treefall for plants, that provides opportunities for successful colonization by foundress queens. Driver ants have also been shown to be the most important predator of termites in Africa (Gotwald, 1995). Even the long legs and leaping ability of certain African shrews are suggested to be adaptations for escape during driver ant attack (Brosset, 1988). Driver ants, because of their affect on ant populations and other insects, have been labelled 'keystone species' (Gotwald, 1995). An investigation of African and Malagasy ant community ecology provides a natural experiment for evaluating the effect of these dominant African ants on ant communities.

Possible consequences of the imbalance in the fauna in Madagascar may include unique island radiations. The lack of driver ants in Madagascar may have spurred the diversification of the tribe Cerapachyini (*Cerapachys* and *Simopone*) on the island (Hölldobler and Wilson, 1990; Olson and Ward, in press). Recent faunal surveys of the ants of Madagascar have revealed an unprecedented morphologically diverse and species rich assemblage of cerapachyine species. Certain *Cerapachys* are morphologically convergent with the driver ant genus *Aenictus* found in Africa. In recent phylogenetic analyses, the cerapachyines appear to be a sister group to the army ants (Cerapachyinae + (Ecitoninae + Dorylinae + Aenictogitoninae + Aenictinae); Bolton, 1990; Baroni Urbani *et al.*, 1992) and have convergently evolved army ant habits of raiding the nests of other ants (Brown, 1975; personal observation). We do not know whether the lack of driver ants in Madagascar has led to the diversification of *Cerapachys* in Madagascar, or has permitted their persistence. *Cerapachys* may have

been free to develop and capitalize on their army ant predatory behaviour or they may have been able to survive despite their less successful behaviour in comparison to driver ants.

Some species of *Cerapachys* and *Simopone* have queens that are winged and both genera are known to nest in preformed plant cavities. It is this combination of nesting in plant cavities and having winged queens that has most likely resulted in their successful colonization of Madagascar. This assumes, that these genera were not present on Madagascar before it separated from Africa (see below).

Mystrium, in the morphologically primitive tribe Amblyoponini, represents another genus that may have been able to persist in the absence of competition from more recently evolved genera that failed to colonize Madagascar. Currently there are six described species of *Mystrium* from Madagascar, while only one species from west Africa and one species from the Oriental, Indo-Australian, and Australian regions still exists. *Mystrium*, along with the other genera in the tribe, are specialized arthropod predators and possess many anatomically and behaviourally primitive characters. For example, personal observations of *Mystrium* and parallel observations of the genus *Amblyopone* suggest they lack social food transfer and the queens are exclusively dependent on the hemolymph of their own larvae (Wheeler and Wheeler, 1988).

Origin of the ant genera of Madagascar

Where did the ant fauna of Madagascar come from? Which taxa were present when Madagascar separated from Africa? We have no fossil records of any ants from Madagascar that could help us understand the Mesozoic or Tertiary ant fauna of Madagascar. The origin of the ant fauna of Madagascar must be inferred from our understanding of the movements of continents and our scant knowledge of the ancestry of ants. The high number of extant ant genera found in Baltic amber fossils (Oligocene) suggests that many extant genera may be at least 35 million years old (Hölldobler and Wilson, 1990). Since only one of 10 genera of ants found in Cretaceous and Eocene deposits are extant genera, most modern taxa may not have been abundant or even extant before the Oligocene. If this is true, then all modern genera probably evolved after the proposed separation of Madagascar from Africa (around 120 million years ago (Ma); Rabinowitz *et al.*, 1983) and the separation of India from Madagascar > 80 Ma; Storey *et al.*, 1995). By the early Cretaceous, Madagascar had reached its present position (Storey *et al.*, 1995).

The tribe Amblyoponini is diverse in Madagascar at the genus and species level. They may represent relic taxa still present in Madagascar that once had a wider distribution. Brown (1960) proposed that the extent Amblyoponini represent surviving species of a much more varied group of ants that date back to the Cretaceous. Even if this ancestral group of ants diversified as early as the Cretaceous, then the amblyoponini genera *Adetomyrma*, *Amblyopone*, *Mystrium* and *Prionopelta* could have arrived only onto Madagascar by queen dispersal of rafting. Dispersal during the Cretaceous, however, may have been easier than during recent times. Briggs (1989) suggested that during the Cretaceous, India was in direct contact with Northern Africa, and the Seychelles had already broken off from India as it moved northward. The proximity of the Seychelles to Madagascar and India could have provided a means of faunal exchange with India and Africa during the Cretaceous. Recent faunal inventories in Madagascar have shown that the most species-rich and abundant genera of Madagascar are *Pheidole*, *Crematogaster*, *Tetramorium* and

Camponotus. Because of their virtual absence in pre-Miocene deposits, Brown (1973) proposed that these genera experienced an explosive radiation since the Miocene, and possibly slightly earlier for *Camponotus*. Assuming this is correct, their arrival in Madagascar must be from long distance dispersal. Phylogenetic studies will be needed to show whether the species in Madagascar are derived from an early stock of these genera, or represent species derived from a series of colonizations by these genera. Evidence from *Tetramorium* suggests that a majority of species may have arisen from an old evolutionary stock. Bolton (1976, 1979) observed that *Tetramorium* with 11-segmented antennae dominate in Australia and Madagascar, where they outnumber the 12-segmented forms which dominate elsewhere around the world. Bolton postulated that the 11-merous forms are remnants of an earlier radiation that has been replaced elsewhere by later-evolving 12-segmented forms. As with the genus *Myrmica*, evidence suggests that Madagascar contains taxa with numerous plesiomorphic characters.

The above discussion assumes that the modern ant genera are not older than the Cretaceous. An alternative scenario, though not supported by the fossil record, is to hypothesize that some extant genera are much older and may date back to around the break up of Gondwana, around 100 million years ago (Ma). Bolton (1990) and Baroni Urbani *et al.* (1992) offer some indirect evidence to support this. They hypothesized that the Old and New World army ants are sister groups, and therefore form a monophyletic clade and share a common ancestor. It is most parsimonious to assume that the ancestor of this clade had wingless queens as do all the modern taxa. The present distribution of Ecitoninae in the New World and Dorylinae in the Old World, may be explained by a common ancestor present before the complete breakup of South America and Africa which is estimated to have occurred between 84 Ma and 106 Ma (Pitman *et al.*, 1993). Since the Dorylinae are absent from Madagascar, this common ancestor may have evolved after the separation of Madagascar from Africa around 120 Ma (Rabinowitz *et al.*, 1983). Other genera, such as in the tribe Amblyoponini which have a Gondwanan distribution, may actually be much older than the Cretaceous.

One of the most perplexing finds in Madagascar was the 1993 discovery of the new genus *Adetomyrma* (Ward 1994). This genus was not only found to belong to a group that has retained many primitive characters, but it also created a seemingly insoluble problem in the reconstruction of the internal phylogeny of the ants. *Adetomyrma* lacks two derived characters, previously thought to be unique and irreversible, of the subfamily Ponerinae in which it is placed (tergosternal fusion of abdominal segments 3 and 4). Despite lacking these two apomorphic characters, it does possess all the other apomorphies of the tribe Amblyoponini. Faunal inventories and the greatly needed revisionary work on the ants of Madagascar, hold great promise to provide new insight and clarification of the higher phylogeny of ants.

Exotic ants

There are almost 20 ant species that are believed to have been introduced into Madagascar (Table 1). Some of these species were present when Forel (1891), completed the first treatment of the Malagasy ant fauna. Open and disturbed habitats in Madagascar and other islands are the most susceptible to invasion by the exotic species. Some species, such as *Technomyrmex albipes*, have begun to invade undisturbed forest sites near rice fields and villages in the northeast of Madagascar. There is some danger that even though small remnant patches of forest may be

preserved in the region, invasion by aggressive, exotic ants may drive native ants locally extinct. On smaller islands such as Mauritius, this phenomenon is already in its late stages. Exotic ants have followed the advance of invasive weeds and have left the island with little chance for preserving most of the approximately 10 endemic species, including the endemic genus *Ireneopone* (Ward, 1990).

Conclusions

Our understanding of the ant fauna of Madagascar is far from complete. Almost every one of the 48 indigenous genera is in need of revision. Nonetheless, evidence does suggest that the Malagasy ant fauna shows closer ties with Africa than with the Orient or Indo-Australian regions. The absence of dominant genera found in Africa, such as *Oecophylla*, *Aenictus* and *Dorylus* had posed unique opportunities for the ants of Madagascar, possibly allowing some to persist and others to radiate. The completion of taxonomic revisions of the most important genera of the region, coupled with phylogenetic analysis, will help clarify these hypotheses.

Acknowledgements

I thank G. D. Alpert, B. Bolton, W. L. Brown, H. Robertson, and P. S. Ward for sharing their vast field experience and collections in compiling the distribution of the world ant genera and the species list for the Malagasy region. I am grateful to B. Bolton and P. S. Ward for comments on the manuscript and for their constant enthusiasm and discussion on all aspects of ant systematics. Field work that provided the basis for this work could not have been completed without the gracious support of the Malagasy people and the funding by World Wide Fund for Nature—Madagascar, National Geographic Society (no. 5152-93), and the National Science Foundation (INT 9319515).

References

- ALPERT, G. D., 1992, Observations on the genus *Terataner* in Madagascar (Hymenoptera: Formicidae), *Psyche* (Camb.), **99**, 117–127.
- ALPERT, G. D. and RABESON, P. R., in press, Notes on *Eutetramorium*, a rare ant genus endemic to Madagascar, with a description of a singular new species (Hymenoptera: Formicidae), *Psyche* (Camb.).
- ANDRÉ E., 1887, Description de quelques fourmis nouvelles ou imparfaitement connues, *Revue d'Entomologie* (Caen), **6**, 280–298.
- ANDRÉ E., 1890, Matériaux pour servir à la faune myrmécologique de Sierra Leone (Afrique occidentale), *Revue d'Entomologie*, **9**, 311–327.
- BARONI URBANI, C., 1972, Studi sui *Camponotus* (Hymenoptera: Formicidae), *Verhandlungen der Naturforschenden Gesellschaft in Basel*, **82**, 122–135.
- BARONI URBANI, C., 1973, Die Gattung *Xenometa*, ein objektives Synonym (Hymenoptera, Formicidae), *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, **46**, 199–201.
- BARONI URBANI, C., 1978, Contributo alla conoscenza del genere *Amblyopone* Erichson (Hymenoptera:Formicidae), *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, **51**, 39–51.
- BARONI URBANI, C., BOLTON, B. and WARD, P. S., 1992, The internal phylogeny of ants (Hymenoptera:Formicidae), *Systematic Entomology*, **17**, 301–329.
- BERNARD, F., 1953, La réserve naturelle intégrale du Mt Nimba. 11. Hyménoptères Formicidae, *Mémoires de l'Institut Français d'Afrique Noire*, **19**, 165–270.
- BOLTON, B., 1974, A revision of the Palaeotropical arboreal ant genus *Cataulacus* F. Smith (Hymenoptera: Formicidae), *Bulletin of the British Museum (Natural History)*, *Entomology*, **30**, 1–105.

- BOLTON, B., 1975, A revision of the ant genus *Leptogenys* Roger (Hymenoptera:Formicidae) in the Ethiopian region with a review of the Malagasy species, *Bulletin of the British Museum (Natural History)*, **Entomology**, **31**, 235–305.
- BOLTON, B., 1976, The ant tribe Tetramoriini (Hymenoptera:Formicidae). Constituent genera, review of smaller genera and revision of Triglyphothrix Forel, *Bulletin of the British Museum (Natural History)*, **Entomology**, **34**, 281–379.
- BOLTON, B., 1977, The ant tribe Tetramoriini (Hymenoptera:Formicidae). The genus *Tetramorium* Mayr in the Oriental and Indo-Australian regions, and in Australia, *Bulletin of the British Museum (Natural History)*, **Entomology**, **36**, 67–151.
- BOLTON, B., 1979, The ant tribe Tetramoriini (Hymenoptera:Formicidae). The genus *Tetramorium* Mayr in the Malagasy region and in the New World, *Bulletin of the British Museum (Natural History)*, **Entomology**, **38**, 129–181.
- BOLTON, B., 1980, The ant tribe Tetramoriini (Hymenoptera:Formicidae). The genus *Tetramorium* Mayr in the Ethiopian zoogeographical region, *Bulletin of the British Museum (Natural History)*, **Entomology**, **40**, 193–384.
- BOLTON, B., 1981, A revision of six minor genera of Myrmicinae (Hymenoptera:Formicidae) in the Ethiopian zoogeographical region, *Bulletin of the British Museum (Natural History)*, **Entomology**, **43**, 245–307.
- BOLTON, B., 1982, Afrotropical species of the myrmicine ant genera *Cardiocondyla*, *Leptothorax*, *Melissotarsus*, *Messor* and *Cataulacus* (Formicidae), *Bulletin of the British Museum (Natural History)*, **Entomology**, **45**, 307–370.
- BOLTON, B., 1983, The Afrotropical dacetine ants (Formicidae), *Bulletin of the British Museum (Natural History)*, **Entomology**, **46**, 267–416.
- BOLTON, B., 1987, A review of the *Solenopsis* genus-group and revision of Afrotropical *Monomorium* Mayr (Hymenoptera:Formicidae), *Bulletin of the British Museum (Natural History)*, **Entomology**, **54**, 263–452.
- BOLTON, B., 1990, Army ants reassessed: the phylogeny and classification of the doryline section, *Journal of Natural History*, **24**, 1339–1364.
- BOLTON, B., 1995a, A taxonomic and zoogeographical census of the extant ant taxa (Hymenoptera:Formicidae), *Journal of Natural History*, **29**, 1037–1056.
- BOLTON, B., 1995b, *A new general catalogue of the ants of the world* (Cambridge, MA: Harvard University Press), 512 pp.
- BRIGGS, J. C., 1987, *Biogeography and plate tectonics* (Amsterdam: Elsevier), x + 204 pp.
- BRIGGS, J. C., 1989, The historic biogeography of India: isolation or contact? *Systematic Zoology*, **38**, 322–332.
- BROSSET, A., 1988, Le peuplement de mammifères insectivores des forêts du nord-est du Gabon, *Revue d'Écologie la Terre et la Vie*, **43**, 23–46.
- BROWN, W. L. Jr., 1960, Contributions toward reclassification of the Formicidae. 3. Tribe Amblyponini, *Bulletin of the Museum of Comparative Zoology at Harvard College*, **122**, 145–230.
- BROWN, W. L. Jr., 1971, Characters and synonymies among the genera of ants. Part IV. Some genera of subfamily Myrmicinae (Hymenoptera:Formicidae), *Breviora*, **365**, 1–5.
- BROWN, W. L. Jr., 1973, A comparison of the Hylean and Congo-West African rain forest ant faunas, in B. J. Meggers, E. S. Ayensu, W. D. Duckworth (eds) *Tropical forest ecosystems in Africa and South America: a comparative review* (Washington, D.C.: Smithsonian Inst. Press) pp. 161–185.
- BROWN, W. L. Jr., 1974, A remarkable new island isolate in the genus *Proceratium* (Hymenoptera:Formicidae), *Psyche* (Camb.), **81**, 70–83.
- BROWN, W. L. Jr., 1975, Contributions toward a reclassification of the Formicidae. V. Ponerinae, tribes Platythyreini, Cerapachyini, Cylindromyrmecini, Acanthostichini, and Aenictogitini, *Search. Agriculture* (Ithaca, NY), **5**(1), 1–115.
- BROWN, W. L. Jr., 1976, Contributions toward a reclassification of the Formicidae. Part VI. Ponerinae, tribe Ponerini, subtribe Odontomachiti. Section A. Introduction, subtribal characters. Genus *Odontomachus*, *Studia Entomologica*, **19**, 67–171.
- BROWN, W. L. Jr., 1978a (1977), An aberrant new genus of myrmicine ant from Madagascar, *Psyche* (Camb.), **84**, 218–224.
- BROWN, W. L. Jr., 1978b, Contributions toward a reclassification of the Formicidae. Part VI. Ponerinae, tribe Ponerini, subtribe Odontomachiti. Section B. Genus *Anochetus* and bibliography, *Studia Entomologica*, **20**, 549–638.

- BROWN, W. L. Jr., 1980 (1979), A remarkable new species of *Proceratium*, with dietary and other notes on the genus (Hymenoptera:Formicidae), *Psyche* (Camb.) **86**, 337–346.
- CRAWLEY, W. C., 1921, New and little-known species of ants from various localities, *Annals and Magazine of Natural History*, (9)7, 87–97.
- DALLA TORRE, K. W. von, 1893, *Catalogus Hymenopterorum hucusque descriptorum systematicus et synonymicus. Vol. 7. Formicidae (Heterogyna)* (Leipzig: W. Engelmann), 289 pp.
- DONISTHORPE, H., 1942, The Formicidae (Hym.) of the Armstrong College Expedition to the Siwa Oasis, *Annals and Magazine of Natural History*, (11)9, 26–33.
- DONISTHORPE, H., 1946a (1945), New species of ants (Hym., Formicidae) from the Island of Mauritius, *Annals and Magazine of Natural History*, (11)12, 776–782.
- DONISTHORPE, H., 1946b, *Ireneopone gibber* (Hym., Formicidae), a new genus and species of myrmicine ant from Mauritius *Entomologist's Monthly Magazine*, **82**, 242–243.
- DONISTHORPE, H., 1946c, The ants (Hym. Formicidae) of Mauritius *Annals and Magazine of Natural History*, (11)13, 25–35.
- DONISTHORPE, H., 1946d, A new genus and species of Formicidae (Hym.) from Mauritius, *Proceedings of the Royal Entomological Society of London. Series B*, **15**, 145–147.
- DONISTHORPE, H., 1947 (1946), New species of ants from China and Mauritius, *Annals and Magazine of Natural History*, (11)13, 283–286.
- DONISTHORPE, H., 1949, A new *Camponotus* from Madagascar and a small collection of ants from Mauritius, *Annals and Magazine of Natural History*, (12)2, 271–275.
- DUFOUR, L., 1864, Note sur une nouvelle espèce de fourmi (Formica Vinsonnella), *Annales de la Société Entomologique de France*, (4)4, 210.
- EMERY, C., 1877, Catalogo delle formiche esistenti nelle collezioni del Museo Civico di Genova. Parte prima. Formiche provenienti dal Viaggio dei signori Antinori, Beccari e Issel nel Mar Rosso e nel paese dei Bogos, [concl] *Annali del Museo Civico di Storia Naturale*, **9**, 369–381.
- EMERY, C., 1886, Alcune formiche africane, *Bullettino della Società Entomologica Italiana*, **18**, 355–366.
- EMERY, C., 1890, Studii sulle formiche della fauna neotropica, *Bullettino della Società Entomologica Italiana*, **22**, 38–80.
- EMERY, C., 1893, Voyage de M. E. Simon à l'île de Ceylan (janvier-février 1892). Formicides, *Annales de la Société Entomologique de France*, **62**, 239–258.
- EMERY, C., 1894, Mission scientifique de M. Ch. Alluaud aux îles Séchelles (mars, avril, mai 1892). 2e mémoire. Formicides, *Annales de la Société Entomologique de France*, **63**, 67–72.
- EMERY, C., 1894c, Descrizione di una nuova formica di Sicilia, *Naturalista Siciliano. Giornale di Scienze Naturali*, **14**, 28.
- EMERY, C., 1895a, Mission scientifique de M. Ch. Alluaud dans le territoire de Diego-Suarez (Madagascar-nord) (Avril-août 1893). Formicides, *Annales de la Société Entomologique de Belgique*, **39**, 336–345.
- EMERY, C., 1895b, Voyage de M. E. Simon dans l'Afrique australe (janvier-avril 1893). 3e mémoire. Formicides, *Annales de la Société Entomologique de France*, **64**, 15–56.
- EMERY, C., 1899, Formiche di Madagascar raccolte dal Sig. A. Mocquers nei pressi della Baia di Antongil (1897–1898), *Bullettino della Società Entomologica Italiana*, **31**, 263–290.
- EMERY, C., 1905, Le forme paleartiche del *Camponotus maculatus* F., *Rendiconti delle Sessioni della Reale Accademia delle Scienze dell'Istituto di Bologna*, (n.s.) **9**, 27–44.
- EMERY, C., 1912, Études sur les Myrmicinae. [I-IV.] *Annales de la Société Entomologique de Belgique*, **56**, 94–105.
- EMERY, C., 1915, Les *Pheidole* du groupe *megacephala* (Formicidae), *Revue Zoologique Africaine* (Brussels), **4**, 223–250.
- EMERY, C., 1917, Questions de nomenclature et synonymies relatives à quelques genres et espèces de Formicides (Hym.), *Bulletin de la Société Entomologique de France*, **1917**, 94–97.
- EMERY, C., 1920a, Le genre *Camponotus* Mayr. Nouvel essai de la subdivision en sous-genres, *Revue Zoologique Africaine* (Brussels), **8**, 229–260.
- EMERY, C., 1920b, Studi sui *Camponotus*, *Bullettino della Società Entomologica Italiana*, **52**, 3–48.
- EMERY, C., 1920c, La distribuzione geografica attuale delle formiche. Tentativo di spiegarne la genesi col soccorso di ipotesi filogenetiche e paleogeografiche, *Atti della Reale*

- Accademia dei Lincei. Memorie. Classe di Scienze Fisiche, Matematiche e Naturali*, **13**(5), 357–450.
- EMERY, C., 1922, Hymenoptera. Fam. Formicidae. Subfam. Myrmicinae [part], *Genera Insectorum*, **174B**, 95–206.
- EMERY, C., 1925, Hymenoptera. Fam. Formicidae. Subfam. Formicinae, *Genera Insectorum*, **183**, 1–302.
- FABRICIUS, J. C., 1782 (1781), *Species insectorum exhibentes eorum differentias specificas, synonymia, auctorum loca natalia, metamorphosin adiectis observationibus, descriptionibus*. Tome I (Hamburg and Kiel: C. E. Bohn), 552 pp.
- FABRICIUS, J. C., 1793, *Entomologia systematica emendata et aucta. Secundum classes, ordines, genera, species, adjectis synonymis, locis observationibus, descriptionibus*. Tome 2 (Copenhagen [= Hafniae]: C. G. Proft), 519 pp.
- FABRICIUS, J. C., 1798, *Supplementum entomologiae systematicae* (Copenhagen [= Hafniae]: Proft and Storch), 572 pp.
- FABRICIUS, J. C., 1804, *Systema piezatorum secundum ordines, genera, species, adjectis synonymis, locis, observationibus, descriptionibus* (Brunswick: C. Reichard), xiv + 15–439 + 30 pp.
- FISHER, B. L., in press, Ant diversity patterns along an elevational gradient in the Réserve Naturelle Intégrale d'Andringitra, *Fieldiana*.
- FOREL, A., 1881, Die Ameisen der Antille St. Thomas, *Mitteilungen der Münchener Entomologischen Verein*, **5**, 1–16.
- FOREL, A., 1886a, Diagnoses provisoires de quelques espèces nouvelles de fourmis de Madagascar, récoltées par M. Grandidier, *Annales de la Société Entomologique de Belgique*, **30**, ci–cvii.
- FOREL, A., 1886b, Études myrmécologiques en 1886, *Annales de la Société Entomologique de Belgique*, **30**, 131–215.
- FOREL, A., 1887, Fourmis récoltées à Madagascar par le Dr. Conrad Keller, *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, **7**, 381–389.
- FOREL, A., 1891, Les formicides. [part], in A. Grandidier, *Histoire physique, naturelle, et politique de Madagascar*. Volume XX. *Histoire naturelle des Hyménoptères*. Deuxième partie (28e fascicule) (Paris: Hachette et Cie), v + 237 pp.
- FOREL, A., 1892a, Attini and Cryptocerini. Zwei neue Apterostigma-Arten, *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, **8**, 344–349.
- FOREL, A., 1892b, Nouvelles espèces de Formicides de Madagascar (récoltées par M. Sikora), Première série. *Annales de la Société Entomologique de Belgique*, **36**, 516–535.
- FOREL, A., 1892c, Les formicides. [concl.] in A. Grandidier, *Histoire physique, naturelle, et politique de Madagascar*. Volume XX. *Histoire naturelle des Hyménoptères*. Deuxième partie. Supplément au 28e fascicule, pp. 229–280, (Paris: Hachette et Cie).
- FOREL, A., 1892d, Liste der aus dem Somaliland von Hrn. Prof. Dr. Conr. Keller aus der Expedition des Prinzen Ruspoli im August und September 1891 zurückgebrachten Ameisen, *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, **8**, 349–354.
- FOREL, A., 1893, Formicides de l'Antille St. Vincent, récoltées par Mons. H. H. Smith, *Transactions of the Entomological Society of London*, **1893**, 333–418.
- FOREL, A., 1894a, Abessinische und andere afrikanische Ameisen, gesammelt von Herrn Ingenieur Alfred Ilg, von Herrn Dr. Liengme, von Herrn Pfarrer Missionar P. Berthoud, Herrn Dr. Arth. Müller etc., *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, **9**, 64–100.
- FOREL, A., 1894b, Les Formicides de l'Empire des Indes et de Ceylon. Part IV, *Journal of the Bombay Natural History Society*, **8**, 396–420.
- FOREL, A., 1894c, Quelques fourmis de Madagascar (récoltées par M. le Dr. Voltzkow); de Nouvelle Zélande (récoltées par M. W. W. Smith); de Nouvelle Calédonie (récoltées par M. Sommer); de Queensland (Australie) (récoltées par M. Wiederkehr); et de Perth (Australie occidentale) (récoltées par M. Chase), *Annales de la Société Entomologique de Belgique*, **38**, 226–237.
- FOREL, A., 1895a, Nouvelles fourmis de diverses provenances, surtout d'Australie, *Annales de la Société Entomologique de Belgique*, **39**, 41–49.
- FOREL, A., 1895b, Nouvelles fourmis de l'Imerina oriental (Moramanga etc.), *Annales de la Société Entomologique de Belgique*, **39**, 243–251.

- FOREL, A., 1895c, Quelques fourmis du centre de Madagascar, *Annales de la Société Entomologique de Belgique*, **39**, 485–488.
- FOREL, A., 1897, Ameisen aus Nossi-Bé, Majunga, Juan de Nova (Madagaskar), den Aldabra-Inseln und Sansibar, gesammelt von Herrn Dr. A. Voeltzkow aus Berlin. Mit einem Anhang über die von Herrn Privatdocenten Dr. A. Brauer in Marburg auf den Seychellen und von Herrn, *Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft*, **21**, 185–208.
- FOREL, A., 1899, Trois notices myrmécologiques, *Annales de la Société Entomologique de Belgique*, **43**, 303–310.
- FOREL, A., 1900a, Ponerinae et Dorylinae d'Australie récoltés par MM. Turner, Froggatt, Nugent, Chase, Rothney, J.-J. Walker, etc., *Annales de la Société Entomologique de Belgique*, **44**, 54–or.
- FOREL, A., 1900b, Les Formicides de l'Empire des Indes et de Ceylan. Part VII. *Journal of the Bombay Natural History Society*, **13**, 303–332.
- FOREL, A., 1902, Myrmicinae nouveaux de l'Inde et de Ceylan, *Revue Suisse de Zoologie*, **10**, 165–249.
- FOREL, A., 1904 (1903), Note sur les fourmis du Musée Zoologique de l'Académie Impériale des Sciences à St. Pétersbourg, *Ezhgodnik Zoologicheskago Muzeya*, **8**, 368–388.
- FOREL, A., 1905, Miscellanea myrmécologiques II (1905), *Annales de la Société Entomologique de Belgique*, **49**, 155–185.
- FOREL, A., 1907a, The Percy Sladen Trust Expedition to the Indian Ocean in 1905, under the leadership of Mr. J. Stanley Gardiner, M.A. VI. Fourmis des Seychelles, Amirantes, Farquhar et Chagos, *Transactions of the Linnean Society of London. Zoology*, (2)**12**, 91–94.
- FOREL, A., 1907b, Ameisen von Madagaskar, den Comoren und Ostafrika, *Wissenschaftliche Ergebnisse. Reise in Ostafrika*, **2**, 75–92.
- FOREL, A., 1908, Fourmis de Costa-Rica récoltés par M. Paul Biolley, *Bulletin de la Société Vaudoise des Sciences Naturelles*, **44**, 35–72.
- FOREL, A., 1909, Fourmis du Musée de Bruxelles. Fourmis de Benguela récoltées par M. Creighton Wellman, et fourmis du Congo récoltées par MM. Luja, Kohl et Laurent, *Annales de la Société Entomologique de Belgique*, **53**, 51–73.
- FOREL, A., 1910a, Glanures myrmécologiques, *Annales de la Société Entomologique de Belgique*, **54**, 6–32.
- FOREL, A., 1910b, Formicides australiens reçus de MM. Froggatt et Rowland Turner, *Revue Suisse de Zoologie*, **18**, 1–94.
- FOREL, A., 1910c, Note sur quelques fourmis d'Afrique, *Annales de la Société Entomologique de Belgique*, **54**, 421–458.
- FOREL, A., 1912, The Percy Sladen Trust Expedition to the Indian Ocean in 1905, under the leadership of Mr J. Stanley Gardiner, M.A. Volume 4. No. XI. Fourmis des Seychelles et des Aldabras, reçues de M. Hugh Scott, *Transactions of the Linnean Society of London. Zoology*, **15**(2), 159–167.
- FOREL, A., 1913a, Ameisen aus Rhodesia, Kapland usw (Hym.) gesammelt von Herrn G. Arnold, Dr. H. Brauns und Anderen, *Deutsche Entomologische Zeitschrift*, **1913** (Suppl.), 203–225.
- FOREL, A., 1913b, Wissenschaftliche Ergebnisse einer Forschungsreise nach Ostindien ausgeführt im Auftrage der Kgl. Preuss. Akademie der Wissenschaften zu Berlin von H. v. Buttel-Reepen. II. Ameisen aus Sumatra, Java, Malacca und Ceylon. Gesammelt von Herrn Prof. Dr. v. Buttel-Reepen in den Jahren 1911–1912, *Zoologische Jahrbucher. Abteilung für Systematik, Geographie und Biologie der Tiere*, **36**, 1–148.
- FOREL, A., 1916, Fourmis du Congo et d'autres provenances récoltées par MM. Hermann Kohl, Luja, Mayné, etc., *Revue Suisse de Zoologie*, **24**, 397–460.
- FOREL, A., 1918, Quelques fourmis de Madagascar récoltées par le Dr. Friederichs et quelques remarques sur d'autres fourmis, *Bulletin de la Société Vaudoise des Sciences Naturelles*, **52**, 151–156.
- FOREL, A., 1922, Glanures myrmécologiques en 1922, *Revue Suisse de Zoologie*, **30**, 87–102.
- GOTWALD, W. H., Jr., 1995, *Army ants: the biology of social predation* (Ithaca: Cornell University Press), xviii + 302 pp.
- GREGG, R. E., 1958, Two new species of *Metapone* from Madagascar (Hymenoptera: Formicidae), *Proceedings of the Entomological Society of Washington*, **60**, 111–121.

- GRISWOLD, C. E., 1991, Cladistic biogeography of afromontane spiders, *Australian Systematic Botany*, **4**, 73–89.
- HEER, O., 1852, Ueber die Hause-Ameise Madeiras, *Zürcherische Jugend. Naturforschende Gesellschaft*, **54**, 1–24.
- HÖLDOBLER, B., and WILSON, E. O., 1990, *The ants* (Cambridge, MA: Harvard University Press), xii + 732 pp.
- JERDON, T. C., 1851, A catalogue of the species of ants found in Southern India, *Madras Journal of Literature and Science*, **17**, 103–127.
- KARAVAEV, V., 1991, Ameisen aus Aegypten und dem Sudan, *Russkoe Entomologicheskoe Obozrenie*, **11**, 1–12.
- LATREILLE, P. A., 1802, *Histoire naturelle des fourmis, et recueil de mémoires et d'observations sur les abeilles, les araignées, les faucheurs, et autres insectes*, (Paris: Impr. Crapelet (chez T. Barrois)), xvi + 445 pp.
- LINNAEUS, C., 1758, *Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis*. Tomus I. Editio decima, reformata, (Stockholm [= Holmiae]: L. Salvii), 824 pp.
- MAGURRAN, A. E., 1988, *Ecological diversity and its measurement* (Princeton, NJ: Princeton University Press), 179 pp.
- MAMET, R., 1954, The ants (Hymenoptera Formicidae) of the Mascarene Islands, *Mauritius Institute Bulletin*, **3**, 249–259.
- MAYR, G., 1862, Myrmecologische Studien, *Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien*, **12**, 649–776.
- MAYR, G., 1866a, Myrmecologische Beiträge, *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften in Wien. Mathematisch-Naturwissenschaftliche Classe. Abteilung I*, **53**, 484–517.
- MAYR, G., 1866b, Diagnosen neuer und wenig gekannter Formiciden, *Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien*, **16**, 885–908.
- MAYR, G., 1870, Neue Formiciden, *Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien*, **20**, 939–996.
- MAYR, G., 1893, Formiciden von Herrn Dr. Fr. Stuhlmann in Ost-Afrika gesammelt, *Jahrbuch der Hamburgischen Wissenschaftlichen Anstalten*, **10**, 194–201.
- MAYR, G., 1895, Afrikanische Formiciden, *Annalen des Kaiserlich-Königlichen Naturhistorischen Museums in Wien*, **10**, 124–154.
- NYLANDER, W., 1846a, Adnotationes in monographiam formicarum borealium Europae, *Acta Societatis Scientiarum Fennicae*, **2**, 875–944.
- NYLANDER, W., 1846b, Additamentum adnotationum in monographiam formicarum borealium Europae, *Acta Societatis Scientiarum Fennicae*, **2**, 1041–1062.
- OLSON, D. M. and WARD, P. S., (in press), The ant fauna (Hymenoptera:Formicidae) of Kirindy Forest (tropical dry forest) in western Madagascar, *Primate Report*, (Spec. Vol.).
- PITMAN, W. C. III., CANDE, S., LABRECQUE, L. and PINDELL, J., 1993, Fragmentation of Gondwana: the separation of Africa from South America, in P. Goldblatt (ed), *Biological relationships between Africa and South America* (New Haven: Yale University Press), pp 15–34.
- RABINOWITZ, P. D., COFFIN, M. F. and FALVEY, D., 1983, The separation of Madagascar and Africa, *Science*, **220**, 67–69.
- ROGER, J., 1857, Einiges über Ameisen, *Berliner Entomologische Zeitschrift*, **1**, 10–20.
- ROGER, J., 1861, Die Ponera-artigen Ameisen (Schluss), *Berliner Entomologische Zeitschrift*, **5**, 1–54.
- ROGER, J., 1862, Einige neue exotische Ameisen-Gattungen und Arten, *Berliner Entomologische Zeitschrift*, **6**, 233–254.
- ROGER, J., 1863, Die neu aufgeführten Gattungen und Arten meines Formiciden-Verzeichnisses nebst Ergänzung einiger früher gegebenen Beschreibungen, *Berliner Entomologische Zeitschrift*, **7**, 131–214.
- SANTSCHI, F., 1910 (1909), Formicides nouveaux ou peu connus du Congo français, *Annales de la Société Entomologique de France*, **78**, 349–400.
- SANTSCHI, F., 1911a, Nouvelles fourmis de Madagascar, *Revue Suisse de Zoologie*, **19**, 117–134.
- SANTSCHI, F., 1911b, Formicides de diverses provenances, *Annales de la Société Entomologique de Belgique*, **55**, 278–287.
- SANTSCHI, F., 1912, Fourmis d'Afrique et de Madagascar, *Annales de la Société Entomologique de Belgique*, **56**, 150–167.

- SANTSCHI, F., 1913, Glanures de fourmis africaines, *Annales de la Société Entomologique de Belgique*, **57**, 302–314.
- SANTSCHI, F., 1914a, Voyage de Ch. Alluaud et R. Jeannel en Afrique Orientale, 1911–1912. Résultats scientifiques. *Insectes Hyménoptères. II. Formicidae*, pp. 41–148. (Paris: Libr. A. Schulz).
- SANTSCHI, F., 1914b, Formicides de l'Afrique occidentale et australe du voyage de Mr. le Professeur F. Silvestri. *Bollettino del Laboratorio di Zoologia Generale e Agraria della Reale Scuola Superiore d'Agricoltura. Portici*, **8**, 309–385.
- SANTSCHI, F., 1915, Nouvelles fourmis d'Afrique, *Annales de la Société Entomologique de France*, **84**, 244–282.
- SANTSCHI, F., 1919, Fourmis nouvelles éthiopiennes, *Revue Zoologique Africaine* (Brussels), **6**, 229–240.
- SANTSCHI, F., 1923, Descriptions de nouveaux formicides éthiopiens et notes diverses. I, *Revue Zoologique Africaine* (Brussels), **11**, 259–295.
- SANTSCHI, F., 1924, Descriptions de nouveaux formicides africains et notes diverses. II, *Revue Zoologique Africaine* (Brussels), **12**, 195–224.
- SANTSCHI, F., 1926a, Trois notes myrmécologiques, *Annales de la Société Entomologique de France*, **95**, 13–28.
- SANTSCHI, F., 1926b, Description de nouveaux formicides éthiopiens (III^{ème} partie), *Revue Zoologique Africaine* (Brussels), **13**, 207–267.
- SANTSCHI, F., 1928a, Fourmis de îles Fidji, *Revue Suisse de Zoologie*, **35**, 67–74.
- SANTSCHI, F., 1928b, Descriptions de nouvelles fourmis éthiopiennes (quatrième note), *Revue de Zoologie et de Botanique Africaines*, **16**, 54–69.
- SANTSCHI, F., 1928c, Descriptions de nouvelles fourmis éthiopiennes (suite), *Revue de Zoologie et de Botanique Africaines*, **16**, 191–213.
- SANTSCHI, F., 1930, Description de Formicides éthiopiens nouveaux ou peu connus. V, *Bulletin et Annales de la Société Entomologique de Belgique*, **70**, 49–77.
- SANTSCHI, F., 1938, Notes sur quelques Ponera Latr., *Bulletin de la Société Entomologique de France*, **43**, 78–80.
- SAS INSTITUTE, 1994, *JMP statistics and graphic guide, version 3* (Cary, NC: SAS Institute), 580 pp.
- SHATTUCK, S. O., 1994, Taxonomic catalog of the ant subfamilies Aneuretinae and Dolichoderinae (Hymenoptera:Formicidae), *University of California Publications in Entomology*, **112**, i–xix, 1–241.
- SMITH, F., 1851, *List of the specimens of British animals in the collection of the British Museum. Part VI. Hymenoptera, Aculeata* (London: British Museum (Natural History)), 134 pp.
- SMITH, F., 1857, Catalogue of the hymenopterous insects collected at Sarawak, Borneo; Mount Ophir, Malacca; and at Singapore, by A. R. Wallace. [part], *Journal and Proceedings of the Linnean Society of London. Zoology*, **2**, 42–88.
- SMITH, F., 1858, *Catalogue of hymenopterous insects in the collection of the British Museum. Part VI. Formicidae* (London: British Museum (Natural History)), 216 pp.
- SMITH, F., 1859, Catalogue of hymenopterous insects collected by Mr. A. R. Wallace at the islands of Aru and Key. [part], *Journal and Proceedings of the Linnean Society of London. Zoology*, **3**, 132–158.
- SMITH, F., 1861, Catalogue of hymenopterous insects collected by Mr. A. R. Wallace in the islands of Ceram, Celebes, Ternate, and Gilolo. [part], *Journal and Proceedings of the Linnean Society of London. Zoology*, **6**, 36–48.
- SMITH, F., 1876a, Preliminary notice of new species of Hymenoptera, Diptera, and Forficulidae collected in the island of Rodriguez by the naturalists accompanying the Transit-of-Venus expedition, *Annals and Magazine of Natural History*, (4)**17**, 447–451.
- SMITH, F., 1876b, Descriptions of new species of Cryptoceridae, belonging to the genera *Cryptoceus*, *Meranoplus* and *Cataulacus*, *Transactions of the Entomological Society of London*, **1876**, 603–612.
- SMITH, F., 1879, Descriptions of new species of aculeate Hymenoptera collected by the Rev. Thos. Blackburn in the Sandwich Islands, *Journal and Proceedings of the Linnean Society of London. Zoology*, **14**, 674–685.
- STOREY, M., MAHONEY, J. J., SAUNDERS, A. D., DUNCAN, R. A., KELLEY, S. P. and COFFIN, M. F., 1995, Timing of hot spot-related volcanism and the breakup of Madagascar and India, *Science*, **267**, 852–855.
- TRAGER, J. C., 1984, A revision of the genus *Paratrechina* (Hymenoptera:Formicidae) of the continental United States, *Sociobiology*, **9**, 49–162.

Genus	Species	Author reference	Additional reference	Nomenclature and distribution notes	Mad.	Mau.	Réu.	Sey.	Com.	Ald.	Far.	Cha.	Rod.
	<i>christi foersteri</i>	Forel 1886b:185	Wheeler 1922c										
	<i>christi maculiventris</i>	Emery 1895a:344	Wheeler 1922c										
	<i>dromedarius</i>	Forel 1891:65	Wheeler 1922c										
	<i>dromedarius pulcher</i>	Forel 1892c:234	Wheeler 1922c										
	<i>gibber</i>	Forel 1891:59	Wheeler 1922c										
	<i>kellari</i>	Forel 1886b:186	Wheeler 1922c										
	<i>kellari invalidus</i>	Forel 1897:200	Wheeler 1922c										
	<i>lubbocki</i>	Forel 1886b:186	Wheeler 1922c										
	<i>lubbocki christoides</i>	Forel 1891:61	Wheeler 1922c										
	<i>lubbocki rectus</i>	Forel 1891:217	Wheeler 1922c										
	<i>pictipes</i>	Forel 1891:217	Wheeler 1922c										
	<i>quadrinaculatus</i>	Forel 1886a:cii	Wheeler 1922c										
	<i>quadrinaculatus</i>	Forel 1892c:233	Wheeler 1922c										
	<i>immaculatus</i>	Emery 1925:123	Bolton 1995b										
	<i>opacatus</i>	Emery 1895a:344	Wheeler 1922c										
	<i>quadrinaculatus</i>	Forel 1897:187	Wheeler 1922c										
	<i>sellaris</i>	Forel 1912:166	Wheeler 1922c										
	<i>repens</i>	Forel 1891:51	Wheeler 1922c										
<i>Camponotus (Myrmamblys)</i>	<i>ihomasseti</i>	André 1887:281	Wheeler 1922c	Replacement name for <i>Myrma</i>									
<i>Camponotus (Myrmepinotus)</i>	<i>echinoploides</i>	Forel 1891:50	Wheeler 1922c	madagascariensis Forel 1886a:civ.									
	<i>edmondi</i>	Forel 1897:200	Wheeler 1922c										
	<i>edmondi ernesti</i>	Roger 1863:135	Wheeler 1922c										
	<i>ethicus</i>	Forel 1891:53	Wheeler 1922c										
	<i>robustus</i>	Forel 1895b:243	Wheeler 1922c										
	<i>subrexi</i>	Emery 1895a:344	Wheeler 1922c										
	<i>heteroclitus</i>	Emery 1899:289	Wheeler 1922c										
	<i>levellii</i>	Forel 1892c:234	Wheeler 1922c										
	<i>macquersyi</i>	Forel 1892c:236	Wheeler 1922c										
	<i>puccatus</i>	Emery 1920a:242	Wheeler 1922c										
	<i>recaurui</i>	Forel 1910c:457	Bolton 1995b										
	<i>sikorai</i>	Forel 1886b:179	Wheeler 1922c										
<i>Camponotus (Myrmopelta)</i>	<i>arminius</i>	Forel 1891:44	Wheeler 1922c										
<i>Camponotus (Myrmopitromis)</i>	<i>darvini</i>	Forel 1905:165	Wheeler 1922c	Unavailable name; described									
	<i>darvini rubropilosus</i>	Forel 1910c:456	Wheeler 1922c	from Madagascar									
	<i>var. rubesitor</i>	Santschi 1926b:263	Bolton 1995b										
	<i>descarpentrisi</i>	Forel 1891:37	Wheeler 1922c										
	<i>elliotti</i>	Santschi 1911a:133	Wheeler 1922c										
	<i>elliotti relicens</i>	Mayr 1862:665	Wheeler 1922c										
	<i>niveosetosus</i>	Forel 1886a:civ	Wheeler 1922c										
	<i>madagascariensis</i>	André 1887:281	Wheeler 1922c										
	<i>nooitsheniensis</i>												

Appendix I continued.....

Genus	Species	Author reference	Additional reference	Nomenclature and distribution notes	Mad.	Mau.	Réu.	Sey.	Com.	Ald.	Far.	Cha.	Rod.
	<i>radovae</i>	Forel 1886a:civ	Wheeler 1922c										
	<i>radovae</i>	Forel 1891:46	Wheeler 1922c										
	<i>radovae darvini</i>												
	<i>ursus</i>	Forel 1886a:civ	Wheeler 1922c										
	<i>voeltzkowii</i>	Forel 1894c:226	Wheeler 1922c										
	<i>imitator</i>	Forel 1891:209	Wheeler 1922c										
<i>Camponotus (Myrmomyrma)</i>	<i>imitator resinicola</i>	Santschi 1911a:133	Wheeler 1922c										
	<i>batesii</i>	Forel 1895b:243	Wheeler 1922c										
<i>Camponotus (Myrmosambus)</i>	<i>baatesii</i>	Roger 1863:134	Wheeler 1922c										
<i>Camponotus (Myrmosericus)</i>	<i>aurous</i>	Forel 1897:203	Wheeler 1922c										
<i>Camponotus (Myrmotrema)</i>	<i>foraminosus</i>												
	<i>aldbrandis</i>												
	<i>foraminosus raspoli</i>	Forel 1892a:349	Wheeler 1922c										
	<i>foraminosus raspoli</i> var. <i>for.</i>	Forel 1907b:90	Wheeler 1922c										
	<i>rotundata</i>												
	<i>grandidieri</i>	Forel 1886a:civ	Wheeler 1922c										
	<i>grandidieri atrabilis</i>	Santschi 1915:273	Wheeler 1922c										
	<i>grandidieri comorensis</i>	Santschi 1915:274	Wheeler 1922c										
	<i>olivieri freyeri</i>	Santschi 1915:270	Wheeler 1922c										
	<i>robecchii assumptionis</i>	Crawley 1921:95	Bolton 1995b										
	<i>sericeus</i>	(Fabricius) 1798:279	Wheeler 1922c										
<i>Camponotus (Orthotomomyrmex)</i>	<i>buteleti</i>	Forel 1905:164	Wheeler 1922c										
	<i>cervicalis</i>	Roger 1863:134	Wheeler 1922c										
	<i>cervicidis gauliei</i>	Santschi 1991a:128	Wheeler 1922c										
	<i>concolor</i>	Forel 1891:214	Bolton 1995b										
	<i>dufouri</i>	Forel 1891:16	Wheeler 1922c										
	<i>dufouri imerinensis</i>	Forel 1891:18	Wheeler 1922c										
	<i>galienii = concolor</i>	Forel 1916:457	Wheeler 1922c										
	<i>gertherti</i>	Donisthorpe 1949:271	Bolton 1995b										
	<i>goudi</i>	Forel 1886a:civ	Wheeler 1922c										
	<i>hagensii</i>	Forel 1886b:158	Wheeler 1922c										
	<i>hildebrandii</i>	Forel 1886b:161	Wheeler 1922c										
	<i>hildebrandii</i>	Emery 1920b:25	Bolton 1995b										
	<i>dichromothrix</i>												
	<i>hova</i>	Forel 1891:35	Wheeler 1922c										
	<i>hova becki</i>	Santschi 1923:292	Wheeler 1922c										
	<i>hova boivini</i>	Forel 1891:34	Wheeler 1922c										
	<i>hova fairmairi</i>	Santschi 1991a:130	Wheeler 1922c										
	<i>hova fulvus</i>	Emery 1894:72	Wheeler 1922c										
	<i>hova hovatorovoides</i>	Forel 1892c:252	Wheeler 1922c										
	<i>hova hovoides</i>	Dalla Torre 1893:249	Wheeler 1922c										
	<i>hova lateolus</i>	Emery 1925:85	Wheeler 1922c										
	<i>hova maculatooides = maculatus</i>	Emery 1920b:6	Wheeler 1922c										
	<i>maculatus strangularis</i>												
	<i>hova mixellus</i>	Dalla Torre 1893:249	Wheeler 1922c										
	<i>hova obscuratus</i>	Emery 1925:85	Wheeler 1922c										

Genus	Species	Author reference	Additional reference	Nomenclature and distribution notes	Mad.	Mau.	Réu.	Sey.	Com.	Ald.	Far.	Cha.	Rod.	
<i>Lasini</i> <i>Paratrechina</i>	<i>hova radamae</i>	Forel 1891:31	Wheeler 1922c		1									
	<i>legionarium</i>	Santschi 1911b:283 (Fabricius) 1782:491	Wheeler 1922c	West Africa	1									
	<i>maculatus</i>	Forel 1905:30	Wheeler 1922c	Unavailable name; described from Madagascar	1	1	1							
	<i>maculatus lividior</i> = <i>maculatus</i>	Santschi 1911a:128	Wheeler 1922c											
	<i>maculatus radamooides</i> = <i>maculatus</i>	Forel 1891:213	Wheeler 1922c	Syn. Baroni Urbani 1972:125	1									
	<i>maculatus strangulatus</i>	Santschi 1911a:129	Wheeler 1922c	Unavailable name; described from Seychelles										
	<i>masculinus fulvus</i> var. <i>fulvoocoronata</i>	Forel 1907a:93	Bolton 1995b	Unavailable name; described from Seychelles										
	<i>masculinus fulvus</i> var. <i>octonotatus</i>	Forel 1897:202	Wheeler 1922c											
	<i>nasicus</i>	Forel 1891:67	Wheeler 1922c		1									
	<i>perroiti</i>	Forel 1897:202	Wheeler 1922c		1									
	<i>perroiti aeshylius</i>	Forel 1913a:224	Wheeler 1922c		1									
	<i>radamae becki</i> var. <i>altior</i>	Santschi 1923:292	Bolton 1995b	Unavailable name; described from Madagascar	1									
	<i>roeseli</i>	Forel 1910a:20	Wheeler 1922c		1									
	<i>amblylops</i>	(Forel) 1892c:240	Wheeler 1922c		1									
	<i>amblylops rubescens</i>	(Forel) 1892b:516	Wheeler 1922c		1									
	<i>bourbonica</i>	(Forel) 1886b:210	Wheeler 1922c; Wilson & Taylor 1967	Pantropical tramp species	1	1	1	1					1	
	<i>bourbonica bengalensis</i>	(Forel) 1894b:408	Wheeler 1922c; Wilson & Taylor 1967; Trager 1984	Syn. Wilson & Taylor 1967:88, but see Trager 1984:151; India, Burma										
	<i>bourbonica farquharensis</i>	(Forel) 1907b:92	Wheeler 1922c											
	<i>bourbonica ngasiyana</i>	(Forel) 1907b:87	Wheeler 1922c											
	<i>comorensis</i>	(Forel) 1907b:87	Wheeler 1922c											
	<i>glabra</i>	(Forel) 1891:92	Wheeler 1922c											
	<i>gracilis</i>	(Forel) 1892b:517	Wheeler 1922c		1									
	<i>humboldtii</i>	(Forel) 1891:90	Wheeler 1922c		1									
<i>longicornis</i>	(Latreille) 1802:113	Wheeler 1922c		1										
<i>madagascarensis</i>	(Forel) 1886b:212	Wheeler 1922c	Pantropical tramp species	1	1	1	1						1	
<i>madagascarensis dilsi</i>	(Forel) 1891:87	Wheeler 1922c		1										
<i>madagascarensis</i>	Wheeler 1922c:1038	Wheeler 1922c		1										
<i>mixta</i>	(Forel) 1897:203	Wheeler 1922c												
<i>mixta sikorae</i>	(Forel) 1892c:238	Wheeler 1922c												
<i>steinheili</i>	(Forel) 1893:342	Wheeler 1922c	Antilles, West Indies											
<i>viduata</i>	Nylander 1846a:900	Donisthorpe 1946c; Mamm 1954	Cosmopolitan	1										
<i>dodo</i>	Donisthorpe 1946a:782	Donisthorpe 1946c; Mamm 1954												

Appendix I continued.....

Genus	Species	Author reference	Additional reference	Nomenclature and distribution notes	Mad.	Mau.	Réu.	Sey.	Com.	Ald.	Far.	Cha.	Rod.
<i>Anoplolepis</i>	<i>gracilipes</i>	(F. Smith) 1857:55	Wheeler 1922c; Bolton 1995b	Replacement name for <i>Formica longipes</i> Jerdon 1851:122.									1
	<i>steingroveri gertrudae capensis albivittata</i>	(Forel) 1900a:77 Mayr 1862:699 Emery 1894:71	Wheeler 1922c Wheeler 1922c Taylor 1967	Pantropical tramp species	1								
<i>Lepisiota</i> <i>Plagiotelepis</i>	<i>exigua</i>	Forel 1894b:417	Wheeler 1922c; Wilson and Taylor 1967	Widespread in Africa	1								
	<i>mauriciana</i>	Forel 1892c:241	Wheeler 1922c; Bolton 1995b	India, West Africa, western China, Hawaii	1	1	1	1					
	<i>ebraudi</i>	Forel 1886a:c	Bolton 1974										
	<i>intrudens</i>	(F. Smith) 1876b:609	Bolton 1974										
	<i>johannae = intrudens</i>	Forel 1895b:250	Bolton 1974:43	Widespread in Africa									
	<i>oberthueri</i>	Emery, in Forel 1891:146	Bolton 1974										
	<i>porcatus</i>	Emery 1899:286	Bolton 1974										
	<i>regularis</i>	Forel 1892c:252	Bolton 1974										
	<i>tenius</i>	Emery 1899:288	Bolton 1974										
	<i>voeltzkowi</i>	Forel 1907b:84	Bolton 1974										
<i>wasmanni</i>	Forel 1897:193	Bolton 1974											
<i>Crematogaster</i>	<i>adrepens</i>	Forel 1897:197	Wheeler 1922c										
	<i>egyptiaca senegalensis</i>	Roger 1863:206	Wheeler 1922c										
	<i>agnatis</i>	Forel 1892b:531	Wheeler 1922c										
	<i>castanea</i>	F. Smith 1858:136	Wheeler 1922c										
	<i>degeeri</i>	Forel 1886a:c	Wheeler 1922c	Africa									
	<i>emmae</i>	Forel 1891:227	Wheeler 1922c										
	<i>emmae laticeps</i>	Forel 1892b:529	Wheeler 1922c										
	<i>ensifera</i>	Forel 1910a:18	Wheeler 1922c										
	<i>gibba</i>	Emery 1894:70	Wheeler 1922c										
	<i>grevei</i>	Forel 1891:183	Wheeler 1922c										
	<i>hova</i>	Forel 1887:387	Wheeler 1922c										
	<i>hova latipoda</i>	Forel 1892b:535	Wheeler 1922c										
	<i>hova mossibensis</i>	Forel 1891:181	Wheeler 1922c										
<i>inops</i>	Forel 1892c:254	Wheeler 1922c											
<i>kelleri</i>	Forel 1891:197	Wheeler 1922c											
<i>madagascariensis</i>	André 1887:297	Wheeler 1922c											
<i>marthae</i>	Forel 1892b:529	Wheeler 1922c											
<i>lobata pacifica</i>	Santschi 1919:236	Santschi 1930:63											
<i>ramavalonae</i>	Forel 1887:388	Wheeler 1922c											
<i>ramavalonae paulinae</i>	Forel 1892b:530	Wheeler 1922c											
<i>ramavalonae</i>	Wasmann 1897:249	Wheeler 1922c; Bolton 1995b											
<i>paulinaeravalonae</i>													<i>nomen nudum</i>

Genus	Species	Author reference	Additional reference	Nomenclature and distribution notes	Mad.	Mau.	Réu.	Sey.	Com.	Ald.	Far.	Cha.	Rod.	
<i>travaliolae</i> <i>pepo</i> <i>rasoherinae</i> <i>rasoherinae</i> <i>brameola</i>		Forel 1922:96	Bolton 1995b											
		Forel 1891:194	Wheeler 1922c											
		Emery 1922:131	Wheeler 1922c; Bolton 1995b	Replacement name for <i>C. rasoherinae</i> Forel 1907b:79										
	<i>schencki</i>	Forel 1891:182	Wheeler 1922c											
	<i>sewelli</i>	Forel 1891:196	Wheeler 1922c											
	<i>sewelli dentata</i>	Dalla Torre 1893:86	Wheeler 1922c											
	<i>sewelli improba</i>	Forel 1907b:80	Wheeler 1922c											
	<i>lobata</i>	Emery 1895a:342	Wheeler 1922c; Bolton 1995b											
	<i>sewelli lobata</i> var.	Forel 1907b:80	Wheeler 1922c											
	<i>gigantica</i>			Unavailable name; described from Madagascar										
	<i>sewelli marnoi</i>	Mayr 1895:138	Wheeler 1922b:844											
	<i>sewelli mauritiana</i>	Forel 1907b:79	Wheeler 1922c											
	<i>medicassa</i>	Emery 1895a:342	Wheeler 1922c; Bolton 1995b											
	<i>tricolor decolor</i> = <i>castanea</i>	Forel 1891:188	Wheeler 1922c											
<i>voeltzkowi</i>	Forel 1907b:78	Wheeler 1922c												
Dactonini														
<i>Quadristrana</i>														
<i>Serrastrana</i>														
Strumigenys														
	<i>emmae</i>	(Emery) 1890:70	Bolton pers. comm.	Pantropical tramp species										
	<i>ludovici</i>	(Forel) 1904:369	Wheeler 1922c											
	<i>raymondii</i> = <i>simoni</i>	(Donisthorpe) 1946a:778	Bolton 1983:350	Syn. Bolton 1983:350										
	<i>simoni</i>	(Emery) 1895b:42	Bolton 1983:350	Eastern and southern Africa; probable introduction in Mauritius										
	<i>godefroyi</i>	Mayr 1866a:516	Wheeler 1922c	Asutralia, Polynesian, Melanesian Islands										
	<i>grandidieri</i>	Forel 1892b:517	Wheeler 1922c											
	<i>scotti</i>	Forel 1912:159	Wheeler 1922c											
	<i>cristata</i>	Santschi 1913:306	Wheeler 1922c											
	<i>emeryi</i>	Forel 1881:5	Bolton 1982											
	<i>emeryi mahdii</i> = <i>emeryi</i>	Karavaiev 1911:8	Bolton 1982											
	<i>emeryi rasalamae</i> = <i>emeryi</i>	Forel 1891:161	Bolton 1982											
	<i>mauritia</i> = <i>emeryi</i>	Donisthorpe 1946a:776	Bolton 1982											
	<i>monilicornis</i> = <i>emeryi</i>	Emery 1917:96	Bolton 1982											
	<i>nuda nereis</i> = <i>emeryi</i>	Wheeler 1927:140	Bolton 1982											
	<i>nuda shuckardoides</i>	Forel 1895b:250	Wheeler 1922c											
	<i>shuckardi</i>	Forel 1891:161	Bolton 1982											
	<i>shuckardi sculpinoides</i>	Santschi 1913:305	Wheeler 1922c	Common and widespread in Africa										
	<i>gibber</i>	Donisthorpe 1946b:242	Mamet 1954											
	<i>angulatus</i>	Mayr 1862:739	Bolton 1982:324											
	<i>latinoides</i> = <i>angulatus</i>	Mayr 1895:130	Bolton 1982:324	Sub-Saharan Africa										
	<i>retusapinosus</i>	Forel 1892c:267	Wheeler 1922c	Syn. provisional Bolton 1982:324										
	<i>albimidi</i>	(Emery) 1895a:341	Wheeler 1922c											
<i>Tetrasera</i>														

Appendix I continued.....

Genus	Species	Author reference	Additional reference	Nomenclature and distribution notes	Mad.	Mau.	Réu.	Sey.	Com.	Ald.	Far.	Cha.	Rod.
Melissoptarini <i>Melissoptarus</i>	<i>foreli</i>	(Emery) 1899:274	Wheeler 1922c										
	<i>rupipes</i>	Emery 1912:104	Wheeler 1922c										
	<i>scotti</i>	(Forel) 1912:160	Wheeler 1922c										
	<i>steinhelli</i>	(Forel) 1895c:485	Wheeler 1922c										
	<i>xalrus</i>	Bolton 1981:296											
	<i>insularis</i>	Santschi 1911a:122	Wheeler 1922c										
Meranoplini <i>Meranoplius</i>													
	<i>mayri</i>	Forel 1910a:19	Wheeler 1922c										
	<i>radamae</i>	Forel 1891:148	Wheeler 1922c										
Metaponini <i>Metapone</i>													
	<i>emersoni</i>	Gregg 1958:115											
	<i>madagascarica</i>	Gregg 1958:111											
	<i>laevithorax alluaudi</i>	Emery 1894:68	Wheeler 1922c										
	<i>pirokae</i>	Forel 1912:162	Wheeler 1922c										
Myrmecini <i>Pristomyrmex</i>													
	<i>bispinosus</i>	(Donisthorpe) 1949:272	Mamet 1954										
	<i>trispinosus</i>	(Donisthorpe) 1946d:145	Mamet 1954	<i>Dodous</i> syn. by Brown 1971:3 <i>Dodous</i> syn. by Brown 1971:3									
Myrmicini <i>Eucetramorium</i>													
	<i>moquerysi</i>	Emery 1899:281	Alpert and Rabeson in press										
	<i>monticelli</i>	Emery 1899:283	Alpert and Rabeson in press										
Phalacromyrmecini <i>Pilotrochus</i>													
	<i>besmerus</i>	Brown 1978a:221											
Phedolini <i>Aphaenogaster</i>													
	<i>belti</i>	Forel 1895b:248	Wheeler 1922c; Bolton 1982										
	<i>friederichsi</i>	Forel 1918:151	Wheeler 1922c										
	<i>goncantha</i>	(Emery) 1899:277	Wheeler 1922c										
	<i>swammerdami</i>	Forel 1886a:cvi	Wheeler 1922c										
	<i>swammerdami clara</i>	(Santschi) 1915:250	Wheeler 1922c										
	<i>swammerdami curia</i>	Forel 1891:169	Wheeler 1922c										
	<i>swammerdami spinipes</i>	Santschi 1911a:123	Wheeler 1922c										
	<i>anastassi cellarum</i>	Forel 1908:55	Donisthorpe 1946c:30	Tramp species, Neotropical, Europe									
	<i>annemariae</i>	(Forel) 1918:152	Wheeler 1922c										
	<i>bessonii</i>	Forel 1891:176, 227	Wheeler 1922c										
	<i>braueri</i>	Forel 1897:204	Wheeler 1922c										
	<i>ensifera</i>	Forel 1897:197	Wheeler 1922c										
	<i>flavens farquharensis</i>	Forel 1907b:91	Wheeler 1922c										
	<i>grallatrix</i>	Emery 1899:278	Wheeler 1922c										
	<i>jonas</i>	Forel 1907b:82	Wheeler 1922c										
	<i>longispinosa</i>	Forel 1891:170	Wheeler 1922c										

Genus	Species	Author reference	Additional reference	Nomenclature and distribution notes	Mad.	Mau.	Reu.	Sey.	Com.	Ald.	Far.	Cha.	Rod.	
Pheidolegontini <i>Oligomyrmex</i>	<i>longispinosa scabrata lucida</i>	Forel 1895b:249	Wheeler 1922c											
	<i>medicassa</i>	Forel 1895b:248	Wheeler 1922c											
	<i>megacephala</i>	Forel 1892b:525	Wheeler 1922c											
	<i>megacephala scabrator</i>	(Fabricius) 1793:361	Wheeler 1922c	Tramp species										
	<i>megacephala spinosa</i>	Forel 1891:178	Wheeler 1922c											
	<i>memorialis</i>	Forel 1891:178	Wheeler 1922c											
	<i>memorialis petax</i>	Forel 1892b:526	Wheeler 1922c											
	<i>oculata</i>	(Emery) 1899:276	Wheeler 1922c											
	<i>oswaldi</i>	Forel 1891:173, 227	Wheeler 1922c											
	<i>oswaldi decollata</i>	Forel 1892b:527	Wheeler 1922c											
	<i>picata</i>	Forel 1891:178	Wheeler 1922c											
	<i>picata bernhardae</i>	Emery 1915:245	Wheeler 1922c											
	<i>picata gutteli</i>	Forel 1905:164	Wheeler 1922c											
	<i>punctulata</i>	(Mayr) 1866b:899	Wheeler 1922c											
	<i>pusilla = megacephala</i>	Heer 1852:15	Wheeler 1992b; Mamet 1954		Africa Syn. Wheeler 1922b:812									
	<i>sikorae</i>	Forel 1891:223	Wheeler 1922c											
	<i>sikorae litigiosa</i>	Forel 1892b:526	Wheeler 1922c											
	<i>tarda</i>	Donisthorpe 1947:285	Mamet 1954											
	<i>vetaratrix</i>	Forel 1891:225	Wheeler 1922c											
	<i>vetaratrix angustinoda</i>	Forel 1892b:526	Wheeler 1922c											
<i>voeltzkowii</i>	Forel 1894c:227	Wheeler 1922c												
Pheidolegontini <i>Oligomyrmex</i>	<i>grandileri</i>	Forel 1891:201	Wheeler 1922c											
	<i>nosindambo</i>	(Forel) 1891:199												
	<i>voeltzkowii</i>	Forel 1907b:77	Wheeler 1922c											
Solenopsis <i>Monomorium</i>	<i>destructor</i>	(Jerdon) 1851:105	Mamet 1954; Bolton 1987:324											
	<i>elongatum</i>	F. Smith 1876a:448	Mamet 1954											
	<i>floricola</i>	(Jerdon) 1851:107	Wheeler 1922b:863 for list of syn. see Bolton 1987:390	Syn. Bolton 1987:390; cosmopolitan tramp species, possible Indian origin										
	<i>foissulatum seckellense</i>	Emery 1894:69	Wheeler 1922c											
	<i>hildebrandti</i>	Forel 1892c:256	Wheeler 1922c; Bolton 1987:288											
	<i>imcrinense</i>	Forel 1892c:257	Wheeler 1922c											
	<i>impressum = floricola</i>	F. Smith 1876a:447	Mamet 1954; Bolton 1987:390											
	<i>madecassum</i>	Forel 1892c:255	Wheeler 1922c; Bolton 1987:288											
	<i>malayensis</i>	Forel 1892c:255	Wheeler 1922c; Bolton 1987:288											
	<i>malayensis</i>	Forel 1892c:255	Wheeler 1922c; Bolton 1987:288											
	<i>malayensis</i>	Forel 1892c:255	Wheeler 1922c; Bolton 1987:288											

Appendix 1 continued.....

Genus	Species	Author reference	Additional reference	Nomenclature and distribution notes	Mad.	Mau.	Reu.	Sey.	Com.	Ald.	Far.	Cha.	Rod.	
Solenopsis <i>Tetramorium</i>	<i>pharaonis</i>	L. 1758:580	Bolton 1987:356											
	<i>pharaonis</i>	Forel 1892d:352	Wheeler 1922c; Bolton 1995b											
	<i>subulatum</i>	Santschi 1928c:196												
	<i>subulatum</i>	L. 1758:580	Wheeler 1922b:870											
	<i>shuckardi</i>	Forel 1895b:281	Wheeler 1922c											
	<i>ternitubum</i>	Forel 1892b:522	Wheeler 1922c											
	<i>geminata</i>	Fabricius 1804:423	Donisthorpe 1946c;											
	<i>mumeti</i>	Donisthorpe 1946a:777	Mamet 1954											
	<i>seychellensis</i>	Forel 1909:55	Wheeler 1922c											
	<i>andrei</i>	Forel 1892c:263	Bolton 1979											
	<i>anisodontum</i>	Bolton 1979:156	Bolton 1979											
	<i>antipodum = caldarium</i>	Wheeler 1927:143	Bolton 1979											
<i>bessonii</i>	Forel 1891:156	Bolton 1979												
<i>bessonii orientale = bessonii</i>	Forel 1895b:247	Bolton 1979												
<i>bicarinatum</i>	(Nylander) 1846b:1061	Bolton 1977, 1980												
Tetramorium	<i>blochmannii montanum = quadrispinosum</i>	Forel 1891:152	Bolton 1979											
	<i>blochmannii = sericeiventris caldarium</i>	Forel 1887:384	Bolton 1979											
	<i>cognatum</i>	(Roger) 1857:12	Bolton 1979, 1980											
	<i>cognatum</i>	Bolton 1979:135	Bolton 1979											
	<i>collum</i>	Bolton 1979:147	Bolton 1979											
	<i>degener</i>	Santschi 1911a:124	Bolton 1979											
	<i>delagoense</i>	Forel 1894a:80	Bolton 1979											
	<i>dysalium</i>	Bolton 1979:141	Bolton 1979, 1980											
	<i>electrum</i>	Bolton 1979:144	see Bolton 1977:94											
	<i>guineense (misidentifications) = bicarinatum</i>													
	<i>humbloii</i>	Forel 1891:154	Bolton 1979, 1980											
	<i>ibycerum</i>	Bolton 1979:148	Bolton 1979, 1980											
<i>isectum</i>	Bolton 1979:145	Bolton 1979												
<i>kelleri</i>	Forel 1887:385	Bolton 1979												
<i>lanuginosum</i>	Mayr 1870:972, 976	Bolton 1979												
<i>laticox</i>	Santschi 1926b:243	Bolton 1979												
<i>laticox</i>	Forel 1895b:247	Bolton 1979												
<i>marginatum</i>	Forel 1895c:485	Bolton 1979												

Genus	Species	Author reference	Additional reference	Nomenclature and distribution notes	Mad.	Mau.	Réu.	Sey.	Com.	Ald.	Far.	Cha.	Rod.
	<i>maricci</i> = <i>lanignosa</i>	(Domishorpe) 1946a:778	Bolton 1976										
	<i>minutum</i> = <i>caldarium</i>	Domishorpe 1942:30	Bolton 1979	Syn. Bolton 1976:351									
	<i>naganum</i>	Bolton 1979:136		Syn. Bolton 1979:169									
	<i>nasonovii</i> = <i>schauffussi</i>	Forel 1892b:521	Bolton 1979	Syn. Bolton 1979:137									
	<i>parallela</i> = <i>simillimum</i>	(F. Smith) 1859:147	Bolton 1980	Syn. Forel 1916:421									
	<i>paupei transformans</i> = <i>caldarium</i>	Santschi 1914a:104	Bolton 1980	Syn. Bolton 1980:310									
	<i>pleganon</i>	Bolton 1979:146											
	<i>plestarum</i>	Bolton 1979:150											
	<i>proximum</i>	Bolton 1979:137											
	<i>pusillum bantouana</i> = <i>simillimum</i>	Santschi 1910:382	Bolton 1980	Syn. Bolton 1980:320									
	<i>pusillum exoleta</i> = <i>simillimum</i>	Santschi 1914b:366	Bolton 1980	Syn. Bolton 1980:320									
	<i>pusillum hmitisi</i> = <i>caldarium</i>	Wheeler 1922a:193	Bolton 1979	Syn. Bolton 1979:169									
	<i>quadrispinosum</i>	Emery 1886:362	Bolton 1979	Widespread in southern Africa, fairly common on Aldabra and Cosmoledo Atoll									
	<i>quastum</i>	Bolton 1979:150											
	<i>ranarum</i>	Forel 1895c:486	Bolton 1979										
	<i>robustior</i>	Forel 1892b:521	Bolton 1979										
	<i>schauffussi</i>	Forel 1891:158	Bolton 1979										
	<i>scytalum</i>	Bolton 1979:157											
	<i>sericeiventre</i>	Emery 1877:370	Bolton 1979	Common and widespread in Africa									
	<i>severini</i>	(Emery) 1895a:343	Bolton 1979										
	<i>sitorae</i>	Forel 1892b:522	Bolton 1979										
	<i>simillimum</i>	(F. Smith) 1851:118	Bolton 1979, 1980	Pantropical tramp species									
	<i>simillimum denticulatum</i> = <i>simillimum</i>	Forel 1902:235	Bolton 1980	Syn. Bolton 1977:131									
	<i>simillimum fasciata</i> = <i>simillimum</i>	Santschi 1928a:69	Bolton 1980	Syn. Bolton 1977:131									
	<i>simillimum madecassum</i>	Forel 1895b:248	Bolton 1979	Syn. Bolton 1979:156									
	<i>simillimum opacior</i> = <i>simillimum</i>	Forel 1913b:81	Bolton 1980	Syn. Bolton 1980:320									
	<i>steinbetti</i>	Forel 1892b:520	Bolton 1979										
	<i>tantillum</i>	Emery 1899:284	Bolton 1979										
	<i>tosii</i>	Santschi 1911a:124	Bolton 1979										
	<i>xanthogaster</i>	Bolton 1979:151											
	<i>zenitum</i>												
	<i>zevarensis</i>	Ward 1994:159											
	<i>zevarensis</i>	Bolton 1979b:49											

POSERINAE
Amblyoponini
Amblyoponinae

Appendix I continued.....

Genus	Species	Author reference	Additional reference	Nomenclature and distribution notes	Mad.	Mau.	Réu.	Sey.	Com.	Ald.	Far.	Cha.	Rod.
<i>Myrtilum</i>	<i>fallax</i>	Forel 1897:192	Brown 1960										
	<i>myrsinicum</i>	Roger 1862:247	Brown 1960										
	<i>oberrubrum</i>	Forel 1897:192	Brown 1960										
	<i>rogeri</i>	Forel 1899:304	Brown 1960										
	<i>stadehmanni</i>	Forel 1895b:251	Brown 1960										
	<i>noelickoni</i>	Forel 1897:189	Brown 1960										
	<i>descarpentriest</i>	Santschi 1924:195											
<i>Prionopelta</i>	<i>arthurii</i>	Forel 1910a:15	Brown 1975										
	<i>bispicus</i>	Emery 1899:268	Brown 1975										
	<i>debilior</i> = <i>mosquersyi</i>	Forel 1907b:76	Brown 1975										
	<i>mosquersyi</i>	Emery 1899:270	Brown 1975	Syn. Brown 1975:8									
	<i>parallela</i>	(F. Smith) 1859:143	Brown 1975	See Brown 1975:8 for complete list of syn.; Indo-Austral tropical region									
	<i>sechellensis</i> = <i>parallela</i>	Forel 1912:159	Brown 1975	Syn. Brown 1975:8									
	<i>wrightsonii</i> = <i>parallela</i>	Forel 1900b:315	Brown 1975	Syn. Brown 1975:8									
<i>Ponerini</i>	<i>friederichsi</i> = <i>madagascarensis</i>	Forel 1918:155	Brown 1978b	Syn. Brown 1978b:557									
<i>Anochetus</i>	<i>grandilieri</i>	Forel 1891:108	Brown 1978b										
	<i>madagascarensis</i>	Forel 1887:382	Brown 1978b										
	<i>madecassus</i> = <i>grandilieri</i>	Santschi 1928b:54	Brown 1978b										
	<i>indagens</i>	(Forel) 1895b:246	Wheeler 1922c										
	<i>indagens bellicosa</i>	(Forel) 1895c:487	Wheeler 1922c										
	<i>johannae</i>	(Forel) 1891:220	Wheeler 1922c										
	<i>ludovicue</i>	(Forel) 1892c:249	Wheeler 1922c										
	<i>madecassa</i>	(Santschi) 1938:79											
	<i>punctatissima glabrata</i>	(Forel) 1895b:245	Wheeler 1922c										
	<i>punctatissima indifferens</i>	(Forel) 1895b:245	Wheeler 1922c										
	<i>punctatissima jugata</i>	(Forel) 1892c:251	Wheeler 1922c										
	<i>ragusai</i>	(Emery) 1894c:28	Wheeler 1922c										
	<i>sakalava</i>	(Forel) 1891:124	Wheeler 1922c										
	<i>sakalava excelisior</i>	(Forel) 1892c:248	Wheeler 1922c										
	<i>acutirostris</i>	Santschi 1912:150	Bolton 1975										
	<i>alluaudi</i>	Emery 1895a:338	Bolton 1975										
	<i>angusta</i>	(Forel) 1892b:519	Bolton 1975										
	<i>antongilensis</i>	Emery 1899:272	Bolton 1975										
	<i>arcirostris</i>	Santschi 1926a:25	Bolton 1975										
	<i>coeruleocans</i>	Emery 1895a:339	Bolton 1975										
	<i>conrensis</i> = <i>stuhmanni</i>	Forel 1907b:76	Bolton 1975	Syn. Bolton 1975:262									
	<i>tribrata</i> = <i>maxillosa</i>	Emery 1895b:20	Bolton 1975	Provisional syn. Bolton 1975:254									
	<i>falcata</i> = <i>maxillosa</i>	Roger 1861:42	Bolton 1975	Syn. Forel 1910b:46									
	<i>falcigera</i>	Roger 1861:42	Bolton 1975	Pantropical tramp species									
	<i>gracilis</i>	Emery 1899:271	Bolton 1975										

Leptogenys

Genus	Species	Author reference	Additional reference	Nomenclature and distribution notes	Mad.	Mau.	Réu.	Sey.	Com.	Ald.	Far.	Cha.	Rod.	
<i>Odontomachus</i>	<i>grandidleri</i>	Forel 1910a:17	Bolton 1975		1									
	<i>incisa</i>	Forel 1891:113	Bolton 1975		1									
	<i>incisa imerincensis</i> =	Forel 1892c:242	Bolton 1975	Syn. Bolton 1975:298										
	<i>incisa</i>			Syn. Bolton 1975:298										
	<i>incisa suarensis</i> =	Emery 1895a:338	Bolton 1975											
	<i>insularis</i> = <i>jaligera</i>	F. Smith 1879:675	Bolton 1975											
	<i>maxillosa</i>	(F. Smith) 1858:93	Bolton 1975	Syn. Wilson and Taylor 1967:30										
	<i>oswaldi</i>	Forel 1891:119	Bolton 1975	Pantropical tramp species	1	1	1	1	1	1			1	
	<i>ridens</i>	Forel 1910a:16	Bolton 1975		1									
	<i>saussurei</i>	(Forel) 1891:121	Bolton 1975		1									
	<i>stahlmanni</i>	Mayr 1893:198	Bolton 1975		1									
	<i>truncatirostris</i>	Forel 1897:195	Bolton 1975		1									
	<i>vinsonella</i> = <i>maxillosa</i>	Dufour 1864:210	Bolton 1975		1									
	<i>voeltzkovi</i>	Forel 1897:194	Bolton 1975		1									
	<i>coquereli</i>	Roger 1861:30	Brown 1976		1									
	<i>minor</i> = <i>coquereli</i>	Emery 1899:273	Brown 1976		1									
	<i>stanleyi</i> = <i>trogodytes</i>	Wheeler 1922a:102	Brown 1976		1									
	<i>trogodytes</i>	Santschi 1914a:58	Brown 1976	Reported in lit. as <i>O. haematodes</i> ; widespread and often common in Africa	1									
	<i>Pachycondyla</i>	<i>ambigua</i>	André 1890:316	Bolton 1995b		1								
		<i>camboneti</i>	(Forel) 1891:133	Wheeler 1922c	Tropical Africa, se Arabia	1								
<i>comorensis</i>		(André) 1887:292	Wheeler 1922c		1									
<i>darwini madecassa</i>		(Emery) 1899:268	Wheeler 1922c; Bolton 1995b		1									
<i>eliseae</i>		(Forel) 1891:221	Wheeler 1922c; Bolton 1995b		1									
<i>melanaria macra</i>		(Emery) 1894a:68	Wheeler 1922c; Bolton 1995b		1									
<i>obscurans</i>		(Walker) 1859:372	Mamet 1954; Bolton 1995b	Oriental-Melanesian regions, ne Australia, spread by commerce to se US	1									
<i>perroti</i>		(Forel) 1891:131	Wheeler 1922c; Bolton 1995b		1									
<i>perroti admista</i>		(Forel) 1892c:251	Wheeler 1922c; Bolton 1995b		1									
<i>sikoreae</i>		(Forel) 1891:127	Wheeler 1922c; Bolton 1995b		1									
<i>Procerastium</i>	<i>testacea</i>	(Bernard) 1953:191	Wilson and Taylor 1967; Bolton 1995b	From Africa to northern Australia in commerce. Probable introduction in Madagascar	1									
	<i>wasmanni</i>	(Forel) 1887:383	Wheeler 1992c; Bolton 1995b		1									
	<i>ardium</i>	Brown 1974:72												
	<i>Tetraponera</i>	<i>aplopyx</i>	Brown 1980:337			1								
		<i>arrogans</i>	(Santschi) 1911a:117	Wheeler 1922c		1								
		<i>demens</i>	(Santschi) 1911b:282	Wheeler 1922c		1								
		<i>dana</i>	(Santschi) 1911a:119	Wheeler 1922c		1								
		<i>exasciata</i>	(Forel) 1892c:261	Wheeler 1922c		1								
		<i>ficatrix</i>	(Forel) 1897:198	Wheeler 1922c		1								
		<i>flexuosa</i>	(Santschi) 1911a:120	Wheeler 1922c		1								
<i>grandidleri</i>		(Forel) 1891:203	Wheeler 1922c		1									
<i>grandidleri hillebrandii</i>		(Forel) 1891:204	Wheeler 1922c		1									
<i>grandidleri variegata</i>		(Forel) 1895c:487	Wheeler 1922c		1									
<i>hysterica</i>	(Forel) 1892c:258	Wheeler 1922c		1										
<i>hysterica dimidiata</i>	(Forel) 1895c:487	Wheeler 1922c		1										
<i>hysterica inflata</i>	(Emery) 1899:273	Wheeler 1922c		1										
<i>mandibularis</i>	(Emery) 1895a:340	Wheeler 1922c		1										
<i>morondaviensis</i>	(Forel) 1891:206	Wheeler 1922c		1										
<i>perlonga</i>	Santschi 1928b:59			1										
<i>plicatidens</i>	(Santschi) 1926a:26			1										
<i>rakotonis</i>	(Forel) 1891:206	Wheeler 1922c		1										
<i>rufoignra</i>	(Jerdon) 1851:111	Wheeler 1922c	India; probable introduction in Seychelles	1										
<i>sahlbergii</i>	(Forel) 1887:386	Wheeler 1922c		1										
<i>sahlbergi spuria</i>	(Forel) 1897:199	Wheeler 1922c		1										
<i>sahlbergi deplanata</i>	(Forel) 1904:375	Wheeler 1922c		1										
<i>sahlbergi longula</i>	(Emery) 1895a:340	Wheeler 1922c		1										

8

7

4

16

24

38

15

42

320

381

Total valid species group names

Grand total valid species-group names

Appendix I continued.....

Genus	Species	Author reference	Additional reference	Nomenclature and distribution notes	Mad.	Mau.	Réu.	Sey.	Com.	Ald.	Far.	Cha.	Rod.	
<i>PSEUDOMYRMECINAE</i>	<i>Pseudomyrmecina</i>	<i>Tetraponera</i>	<i>aplopyx</i>	Brown 1980:337		1								
			<i>arrogans</i>	(Santschi) 1911a:117	Wheeler 1922c		1							
			<i>demens</i>	(Santschi) 1911b:282	Wheeler 1922c		1							
			<i>dana</i>	(Santschi) 1911a:119	Wheeler 1922c		1							
			<i>exasciata</i>	(Forel) 1892c:261	Wheeler 1922c		1							
			<i>ficatrix</i>	(Forel) 1897:198	Wheeler 1922c		1							
			<i>flexuosa</i>	(Santschi) 1911a:120	Wheeler 1922c		1							
			<i>grandidleri</i>	(Forel) 1891:203	Wheeler 1922c		1							
			<i>grandidleri hillebrandii</i>	(Forel) 1891:204	Wheeler 1922c		1							
			<i>grandidleri variegata</i>	(Forel) 1895c:487	Wheeler 1922c		1							
<i>hysterica</i>	(Forel) 1892c:258	Wheeler 1922c		1										
<i>hysterica dimidiata</i>	(Forel) 1895c:487	Wheeler 1922c		1										
<i>hysterica inflata</i>	(Emery) 1899:273	Wheeler 1922c		1										
<i>mandibularis</i>	(Emery) 1895a:340	Wheeler 1922c		1										
<i>morondaviensis</i>	(Forel) 1891:206	Wheeler 1922c		1										
<i>perlonga</i>	Santschi 1928b:59			1										
<i>plicatidens</i>	(Santschi) 1926a:26			1										
<i>rakotonis</i>	(Forel) 1891:206	Wheeler 1922c		1										
<i>rufoignra</i>	(Jerdon) 1851:111	Wheeler 1922c	India; probable introduction in Seychelles	1										
<i>sahlbergii</i>	(Forel) 1887:386	Wheeler 1922c		1										
<i>sahlbergi spuria</i>	(Forel) 1897:199	Wheeler 1922c		1										
<i>sahlbergi deplanata</i>	(Forel) 1904:375	Wheeler 1922c		1										
<i>sahlbergi longula</i>	(Emery) 1895a:340	Wheeler 1922c		1										

8

7

4

16

24

38

15

42

320

381

Total valid species group names

Grand total valid species-group names

/Appendix 2 continued.....

	NEO	NEA	PAL	AFR	(S.A.)	(W.A)	(E.A)	MAL	(India)	ORI	INA	AUS
<i>Rhoptromyrmex</i>			1	1	1	1			1	1	1	1
<i>Rhytidoponera</i>											1	1
<i>Rogeria</i>	1	1									1	1
<i>Romblonella</i>											1	1
<i>Rossomyrmex</i>			1									
<i>Rostromyrmex</i>										1		
<i>Rotastruma</i>										1	1	
<i>Santschiella</i>				1		1						
<i>Scyphodon</i>											1	
<i>Secostruma</i>											1	
<i>Sericomyrmex</i>	1											
<i>Serrastruma</i>				1	1	1	1	1				
<i>Simopelta</i>	1											
<i>Simopone</i>				1	1	1		1		1	1	
<i>Smithistruma</i>	1	1	1	1	1	1	1	1	1	1	1	
<i>Solenopsis</i>	1	1	1	1	1	1	1	1	1	1	1	1
<i>Sphinctomyrmex</i>	1			1		1			1	1	1	1
<i>Stegomyrmex</i>	1											
<i>Stenamma</i>	1	1	1						(1)	1	1	
<i>Stereomyrmex</i>									1	1		
<i>Stigmacros</i>												1
<i>Streblognathus</i>				1	1							
<i>Strongylognathus</i>			1									
<i>Strumigenys</i>	1	1	1	1	1	1	1	1	1	1	1	1
<i>Talaridris</i>	1											
<i>Tapinoma</i>	1	1	1	1	1	1	1	1	1	1	1	1
<i>Tatuidris</i>	1											
<i>Technomyrmex</i>	1		1	1	1	1	1	1	1	1	1	1
<i>Teleutomyrmex</i>			1									
<i>Terataner</i>				1	1	1	1	1				
<i>Teratomyrmex</i>												1
<i>Tetheamyрма</i>												1
<i>Tetramorium</i>	1	1	1	1	1	1	1	1	1	1	1	1
<i>Tetraponera</i>			1	1	1	1	1	1	1	1	1	1
<i>Thaumatomyrmex</i>	1											
<i>Tingimyrmex</i>	1											
<i>Trachymyrmex</i>	1	1										
<i>Tranopelta</i>	1											
<i>Trichoscapa</i>		[1]	1	[1]		[1]						
<i>Tricytarus</i>												
<i>Turneria</i>												1
<i>Typhlomyrmex</i>	1											
<i>Vollenhovia</i>			1						1	1	1	1
<i>Vombisidris</i>									1	1	1	1
<i>Wasmannia</i>	1	[1]										
<i>Willowsiella</i>												1
<i>Xenomyrmex</i>	1	1										
<i>Yavnella</i>			1						1	1		
<i>Zacryptocerus</i>	1	1										
undescribed myrmicine genus								1				
undescribed ponerine genus				1		1						
Total	128	63	73	89	60	83	55	48	74	97	123	88

Anillomyrma and *Decamorium* have recently been recorded from E.A.