

BAJCARIDRIS

kraussii. *Formica kraussii* Forel, 1895d: 231 (w.) ALGERIA.

Emery, 1899b: 18 (m.); Forel, 1902a: 155 (m.).

Combination in *Formica (Proformica)*: Emery, 1909b: 204;

combination in *Proformica*: Kuznetsov-Ugamsky, 1928a: 8;

combination in *Bajcaridris*: Agosti, 1994a: 99.

Status as species: Emery, 1899b: 18; Forel, 1902a: 155; Emery, 1909b: 204; Wheeler, W.M. 1913f: 547 (redescription); Emery, 1925b: 244; Kuznetsov-Ugamsky, 1928a: 8; Agosti, 1994a: 99; Bolton, 1995b: 80; Borowiec, L. 2014: 21.

menozzii. *Proformica menozzii* Santschi, 1923a: 137 (w.) ALGERIA.

Combination in *Formica (Proformica)*: Emery, 1925b: 244;

combination in *Bajcaridris*: Agosti, 1994a: 99.

Status as species: Emery, 1925b: 244; Cagniant, 1970c: 38; Agosti, 1994a: 99; Bolton, 1995b: 80; Cagniant, 2006a: 194; Borowiec, L. 2014: 21.

theryi. *Formica (Proformica) theryi* Santschi, 1936c: 209, fig. 6 (w.) MOROCCO.

Espadaler & Cagniant, 1987: 133 (m.).

Combination in *Proformica*: Cagniant, 1964: 110;

combination in *Bajcaridris*: Agosti, 1994a: 99.

Status as species: Agosti, 1994a: 99; Bolton, 1995b: 80; Cagniant, 2006a: 194; Borowiec, L. 2014: 21.