

**STIGMATOMMA**

**amblyops.** *Stigmatomma amblyops* Karavaiev, 1935a: 57, fig. 1 (w.) VIETNAM.

Type-material: 2 syntype workers.

Type-locality: Vietnam ("Cochinchine"): Cauda, Nr 5675 (*K. Davydov*).

Type-depository: SIZK.

Combination in *Amblyopone*: Brown, 1960a: 167;

combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.

**Status as species:** Chapman & Capco, 1951: 24; Brown, 1960a: 167; Bolton, 1995b: 61; Xu, 2001d: 552 (in key); Xu & Chu, 2012: 1182 (in key).

Distribution: Vietnam.

**annae.** *Amblyopone (Stigmatomma) annae* Arnol'di, 1968: 1801, fig. 2 (w.) TURKMENISTAN.

Type-material: holotype worker.

Type-locality: Turkmenistan: Kara-Kala, nr Syunt, 13.iii.1966 (*A. Tikhomenova*).

Type-depository: ZMUM.

Combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.

**Status as species:** Dlussky & Zabelin, 1985: 209; Dlussky, Soyunov & Zabelin, 1990: 173; Bolton, 1995b: 61; Xu & Chu, 2012: 1186 (in key).

Distribution: Turkmenistan.

**arizonense.** *Stigmatomma pallipes* subsp. *arizonense* Wheeler, W.M. 1915b: 389 (w.) U.S.A. (Arizona).

Type-material: holotype worker.

Type-locality: U.S.A.: Arizona, Huachuca Mts, Ramsay Cañon, 5800 ft (*W.M. Mann*).

Type-depository: MCZC.

Subspecies of *pallipes*: Cole, 1937a: 98.

Junior synonym of *pallipes*: Creighton, 1940b: 3; Creighton, 1950a: 33; Smith, M.R. 1951a: 783; Brown, 1960a: 169; Smith, D.R. 1979: 1335; Bolton, 1995b: 61.

**awa.** *Amblyopone awa* Xu & Chu, 2012: 1192, figs. 51-56 (w.q.) CHINA (Yunnan, Tibet).

Type-material: holotype worker, 5 paratype workers, 2 paratype queens.

Type-locality: holotype China: Yunnan Prov., Cangyuan County, Banlao Town, Huguang Village, 1720 m., 17.iii.2011, No. A11-855, soil sample, monsoon evergreen broadleaf forest (*Y. Hao*); paratypes: 1 worker, 1 queen with same data, 1 worker Yunnan Prov., Cangyuan County, Banhong Town, Nanban Village, 1250 m., 17.iii.2011, No. A11-927 (*Y. Hao*), 1 worker Yunnan Prov., Ximeng County, Lisuo Town, Nankang Village, 740 m., 22.iii.2011, No. A11-1671 (*L. Zhang*), 2 workers, 1 queen Tibet, Linzhi County, Lulang Town, Zhaqu Village, 2380 m., 25.ix.2007, No. A07-432 (*X. Liu*).

Type-depository: SFCY.

Combination in *Stigmatomma*: Yoshimura & Fisher, 2014: 15 (in text); Bharti & Rilta, 2015b: 508.

**Status as species:** Bharti & Rilta, 2015b: 508 (redescription); Bharti, Guénard, *et al.* 2016: 18.

Distribution: China, India.

*barreto* Bruch, 1921; see under **FULAKORA**.

**bellii.** *Amblyopone (Stigmatomma) bellii* Forel, 1900c: 55 (w.) INDIA (Karnataka).

Type-material: syntype workers (number not stated).

Type-locality: India: Kanara (*Bell*).

Type-depository: MHNG.

Combination in *Stigmatomma*: Bingham, 1903: 38;

combination in *Amblyopone*: Brown, 1960a: 167;

combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.

**Status as species:** Bingham, 1903: 38; Emery, 1911d: 24; Chapman & Capco, 1951: 24; Brown, 1960a: 167, 193; Bolton, 1995b: 61; Tiwari, 1999: 32; Xu, 2001d: 552 (in key); Bharti & Wachkoo, 2011: 590 (in key); Xu & Chu, 2012: 1179 (in key); Bharti, Guénard, *et al.* 2016: 18.

Distribution: India.

***besucheti*.** *Amblyopone besucheti* Baroni Urbani, 1978a: 49, figs. 15, 16 (w.) SEYCHELLES IS (La Digue I.).

Type-material: holotype worker, 9 paratype workers.

Type-locality: holotype Seychelles: La Digue I., 28.i.1975 (*P. Schauenberg*); paratypes with same data.

Type-depositories: MHNG (holotype); BMNH, MHNG, NHMB (paratype).

Esteves & Fisher, 2016: 42 (putative q.).

Combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.

**Status as species:** Bolton, 1995b: 61; Dorow, 1996a: 75; Taylor, 1979: 833; Esteves & Fisher, 2016: 34 (redescription).

Distribution: Seychelles, Singapore.

***bierigi*** Santschi, 1930; see under ***FULAKORA***.

***binodosus*.** *Arotropus binodosus* Provancher, 1881a: 206, figs. 32, 33 (q.) CANADA (no state data, probably Québec).

Type-material: holotype male.

[Note: the holotype was misprinted with a queen symbol in the original description, but was actually a male (Provancher, 1887: 240, 241). Lectotype designation of a worker by Gahan & Rohwer, 1917: 306, is set aside as no worker was represented in the original description. Lectotype male designation by Francoeur & Béique, 1966: 141, appears unnecessary as Provancher described only a single specimen, automatically the holotype.]

Type-locality: Canada (no further data) (*L. Provancher*).

[Note: type-locality was probably Québec, Cap Rouge (Provancher, 1887: 241).]

Type-depository: QMNH.

[Misspelled as *binodus* by Creighton, 1940b: 3.]

Provancher, 1887: 240 (w.).

Combination in *Amblyopone*: Provancher, 1887: 240;

combination in *Stigmatomma*: Emery, in Dalla Torre, 1893: 14.

Status as species: Provancher, 1887: 240; Cresson, 1887: 258; Dalla Torre, 1893: 14.

Junior synonym of *pallipes*: Emery, 1895c: 261; Wheeler, W.M. 1908e: 400; Emery, 1911d: 24; Wheeler, W.M. 1911f: 159; Creighton, 1940b: 3; Creighton, 1950a: 33; Smith, M.R. 1951a: 783; Brown, 1960a: 169; Francoeur & Béique, 1966: 141; Smith, D.R. 1979: 1335; Bolton, 1995b: 61.

***bolabola*.** *Stigmatomma bolabola* Esteves & Fisher, 2016: 45, figs. 28-33 (w.) MADAGASCAR.

Type-material: holotype worker.

Type-locality: Madagascar: Montagne d'Akirindro, 7.6 km. 341° NNW Ambinanitelo, -15.28833, 49.54833, 17.iii.2003, BLF08250, CASENT0034580, sifted litter (leaf mold, rotten wood) (*Fisher, Griswold, et al.*)

Type-depository: CASC.

Distribution: Madagascar.

***boltoni*.** *Amblyopone boltoni* Bharti & Wachkoo, 2011: 586, figs. 1-3 (w.) INDIA (Himachal Pradesh).

Type-material: holotype worker, 8 paratype workers.

Type-locality: holotype India: Himachal Pradesh, Terrace, 31.9234°N, 75.9294°E, 430 m., 17.vii.2010, Winkler (*A.A. Wachkoo*); paratypes: 2 workers Himachal Pradesh, Ghati Bilwan, 31.9300°N, 75.9302°E, 400 m., 11.x.2008 (*A.A. Wachkoo*); 6 workers Himachal Pradesh, Khatiar, 32.0057°N, 75.9388°E, 450 m., 18.x.2010, soil core (*A.A. Wachkoo*).

Type-depositories: PUAC (holotype); CASC, PUAC (paratypes).

Combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.

**Status as species:** Xu & Chu, 2012: 1185 (in key); Bharti, Guénard, *et al.* 2016: 18.

Distribution: India.

**bruni.** *Stigmatomma bruni* Forel, 1912a: 45 (w.) TAIWAN.

Type-material: holotype worker.

Type-locality: Taiwan (“Formosa”): Pilam (*H. Sauter*).

Type-depository: probably DEIB.

Forel, 1913f: 183 (m.).

Combination in *Amblyopone*: Brown, 1960a: 167;

combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.

**Status as species:** Forel, 1913f: 183; Wheeler, W.M. 1929g: 58; Teranishi, 1940: 56; Chapman & Capco, 1951: 24; Brown, 1960a: 167, 196; Terayama, 1989b: 344; Bolton, 1995b: 61; Xu, 2001d: 552 (in key); Lin & Wu, 2003: 66; Xu, 2006: 152 (in key); Terayama, 2009: 93; Zhou & Ran, 2010: 101; Xu & Chu, 2012: 1180 (in key).

**Senior synonym of juergii:** Brown, 1960a: 168; Bolton, 1995b: 61; Terayama, 2009: 93.

Distribution: Taiwan.

**caliginosum.** *Amblyopone caliginosa* Onoyama, 1999: 158, figs. 1-9 (w.q.) JAPAN.

Type-material: holotype worker, 5 paratype workers, 1 paratype queen.

Type-locality: holotype Japan: Kanagawa Pref., Cape Manazuru, 13.xii.1981 (*K. Masuko*); paratypes: 4 workers, 1 queen with same data, 1 worker with same data but 19.ix.1982.

Type-depositories: OMNH (holotype); OMNH, OUOJ (paratypes).

Combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19;

combination in *Bannapone*: Eguchi, *et al.* 2015: 82;

combination in *Stigmatomma*: Ward & Fisher, 2016: 691.

**Status as species:** Imai, *et al.* 2003: 219; Xu & Chu, 2012: 1177 (in key); Eguchi, *et al.* 2015: 85 (in key); Hsu, F. *et al.* 2017: 91 (in key).

Distribution: Japan.

*celata* Mann, 1919; see under **FULAKORA**.

**crenatum.** *Amblyopone crenata* Xu, 2001d: 553, figs. 18-20 (w.) CHINA (Yunnan).

Type-material: holotype worker, 3 paratype workers.

Type-locality: holotype China: Yunnan Prov., Mengla County, Manzhuang Village, 900 m., 14.viii.1997, No. A97-1680, semi-evergreen monsoon forest of Xishuangbanna Nature Reserve (*Y. He*); paratypes Yunnan Prov., Mengla County, Menglun Town, Man’gang Village, 1000 m., 9.vii.1997, No. A97-3042, mountain rainforest fragment of Xishuangbanna Nature Reserve (*G. Hu*).

Type-depository: SFCY.

Combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.

**Status as species:** Xu, 2006: 152 (in key); Zhou & Ran, 2010: 102; Guénard & Dunn, 2012: 23; Xu & Chu, 2012: 1182 (in key).

Distribution: China.

**crypticum.** *Bannapone cryptica* Eguchi, *et al.* 2015: 79, figs. 1-11 (w.q.) VIETNAM.

Type-material: holotype worker, 2 paratype workers, 2 paratype queens.

Type-locality: holotype Vietnam: Nghe An, Tuong Duong “Sang Le Forest”, <ca 220 m., 2.iv.2006, Eg02iv06 (*K. Eguchi*); paratypes with same data.

Type-depositories: VNMN (holotype); MCZC (paratypes).

Combination in *Stigmatomma*: Ward & Fisher, 2016: 691.

**Status as species:** Hsu, F. *et al.* 2017: 91 (in key).

Distribution: Vietnam.

**denticulatum.** *Stigmatomma denticulatum* Roger, 1859: 251, pl. 7, fig. 2 (w.) GREECE (Zakynthos I.).

Type-material: holotype (?) worker.

[Note: no indication of number of specimens is given.]

Type-locality: Greece: Zante I. (= Zakynthos I.), Monte Scapo (v. *Kiesenwetter*).

Type-depository: MNHU.

Emery, 1875a: 74 (m.); Emery, 1909c: 357 (q.).

Combination in *Amblyopone*: Emery & Forel, 1879: 455; Brown, 1960a: 168;

combination in *Stigmatomma*: Dalla Torre, 1893: 14; Yoshimura & Fisher, 2012a: 19.

**Status as species:** Mayr, 1861: 53 (in key); Roger, 1861a: 53; Roger, 1863b: 20; Mayr, 1863: 454; André, 1874: 187 (in key); Emery, 1875a: 74; Emery, 1875b: 471; Emery, 1878b: 50; Emery & Forel, 1879: 455; Emery, 1881b: 528; André, 1882c: 233 (in key); Forel, 1886e: clxvii; Saunders, E. 1890: 203; Dalla Torre, 1893: 14; Emery, 1895b: 59; Emery, 1909c: 357; Emery, 1911d: 24; Forel, 1913d: 427; Emery, 1914d: 156; Santschi, 1915a: 55 (in key); Emery, 1916b: 101 (in key); Bondroit, 1918: 80; Müller, 1921: 48; Finzi, 1923: 1; Müller, 1923b: 24; Finzi, 1924a: 15; Emery, 1924c: 164; Finzi, 1928b: 129; Finzi, 1930d: 311; Grandi, 1935: 98; Zimmermann, 1935: 7; Finzi, 1939b: 87; Consani & Zangheri, 1952: 39; Ceballos, 1956: 296; Brown, 1960a: 168, 197; Collingwood & Yarrow, 1969: 55; Baroni Urbani, 1971c: 9; Aktaş, 1977: 118; Collingwood, 1978: 74 (in key); Baroni Urbani, 1978a: 42 (redescription); Schembri & Collingwood, 1981: 420; Agosti & Collingwood, 1987a: 53; Agosti & Collingwood, 1987b: 264 (in key); Kugler, J. 1988: 256; Dlussky, Soyunov & Zabelin, 1990: 174; Tinaut, 1990d: 190; Atanassov & Dlussky, 1992: 61; Mei, 1992a: 412; Bolton, 1995b: 61; Poldi, *et al.* 1995: 2; Bračko, 2006: 134; Cagniant, 2006a: 195; Petrov, 2006: 83 (in key); Vonshak, *et al.* 2009: 38; Lapeva-Gjonova, *et al.* 2010: 4; Legakis, 2011: 3; Borowiec, L. & Salata, 2012: 536; Kiran & Karaman, 2012: 3; Borowiec, L. & Salata, 2013: 365; Borowiec, L. 2014: 161; Lebas, *et al.* 2016: 106; Borowiec, L. & Salata, 2018: 7; Salata & Borowiec, 2018c: 48.

**Senior synonym of *gheorghieffii*:** Emery, 1915a: 255, Emery, 1916b: 100; Brown, 1960a: 168, 197; Baroni Urbani, 1978a: 42; Bolton, 1995b: 61.

**Senior synonym of *gracilicornis*:** Brown, 1960a: 168, 197; Baroni Urbani, 1978a: 42; Bolton, 1995b: 61.

Distribution: Albania, Bulgaria, Croatia, Gibraltar, Greece, Israel, Italy, Lebanon, Malta, Morocco, Romania, Serbia, Spain, Turkey.

**\**electrinum*.** \**Amblyopone electrina* Dlussky, 2009: 1047, figs. 2c,d (m.) BALTIC AMBER (Eocene).

Combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.

**Status as species:** Dlussky & Rasnitsyn, 2009: 1031; Perkovsky, 2016: 115.

*elongata* Santschi, 1912; see under **FULAKORA**.

***emeryi*.** *Amblyopone emeryi* Saunders, E. 1890: 203, fig. (w.) MOROCCO.

Type-material: syntype workers (number not stated, "a small colony").

Type-locality: Morocco: Tangier (*J.J. Walker*).

Type-depositories: BMNH, MHNG.

Emery, 1915a: 254 (m.).

Combination in *Amblyopone*: Brown, 1960a: 168;

combination in *Stigmatomma*: Emery, in Dalla Torre, 1893: 14; Yoshimura & Fisher, 2012a: 19.

**Status as species:** Dalla Torre, 1893: 14; Emery, 1909c: 358; Emery, 1911d: 24; Santschi, 1914a: 430; Santschi, 1915a: 55 (in key); Emery, 1915a: 254; Brown, 1960a: 168; Baroni Urbani, 1978a: 46 (redescription); Tinaut, 1990d: 190; Bolton, 1995b: 62; Cagniant, 2006a: 195; Borowiec, L. 2014: 162; Lebas, *et al.* 2016: 106.

Distribution: Morocco.

- eminia***. *Amblyopone eminia* Zhou, 2001b: 23, 225, figs. 4, 5 (w.) CHINA (Guangxi).  
Type-material: holotype worker.  
Type-locality: China: Guangxi, Huaping Natural Reserve, 17.ix.1998 (*S. Zhou*).  
Type-depository: GNUC.  
Combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.  
**Status as species**: Xu, 2006: 152 (in key); Zhou & Ran, 2010: 102; Guénard & Dunn, 2012: 23; Xu & Chu, 2012: 1179 (in key).  
Distribution: China.
- feae***. *Stigmatomma feae* Emery, 1895k: 454 (w.) MYANMAR.  
Type-material: 3 syntype workers.  
Type-locality: Myanmar (“Birmanian”): Carin, Asciiui Ghecu, 1400-1500 m. (*L. Fea*).  
Type-depository: MSNG.  
Combination in *Amblyopone (Stigmatomma)*: Forel, 1900c: 56;  
combination in *Stigmatomma*: Emery, 1911d: 24; Yoshimura & Fisher, 2012a: 19.  
**Status as species**: Forel, 1900c: 56; Bingham, 1903: 38; Emery, 1911d: 24; Chapman & Capco, 1951: 25; Brown, 1960a: 168; Bolton, 1995b: 62; Xu, 2001d: 552 (in key); Xu & Chu, 2012: 1181 (in key).  
Distribution: Myanmar.
- fulvidum***. *Amblyopone fulvida* Terayama, 1987: 31, figs. 1-5 (w.) JAPAN.  
Type-material: holotype worker.  
Type-locality: Japan: Okinawa Pref., Henzu-jima Is, Yonagusuku-son, 4.i.1985 (*H. Takamine*).  
Type-depository: NIAS.  
Combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19;  
combination in *Bannapone*: Eguchi, *et al.* 2015: 82;  
combination in *Stigmatomma*: Ward & Fisher, 2016: 691.  
**Status as species**: Morisita, *et al.* 1989: 11; Bolton, 1995b: 62; Xu, 2001d: 552 (in key); Imai, *et al.* 2003: 219; Xu & Chu, 2012: 1177 (in key); Eguchi, *et al.* 2015: 85 (in key).  
Distribution: Japan.
- gaetulicum***. *Amblyopone gaetulica* Baroni Urbani, 1978a: 47, figs. 6, 12 (w.) MOROCCO.  
Type-material: holotype worker.  
Type-locality: Morocco: Tangier, iii.1896 (*H. Vaucher*).  
Type-depository: NHMB.  
Combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.  
**Status as species**: Tinaut, 1990d: 190; Bolton, 1995b: 62; Cagniant, 2006a: 195; Borowiec, L. 2014: 162; Lebas, *et al.* 2016: 106.  
Distribution: Gibraltar, Morocco, Spain.
- gheorghieffii***. *Amblyopone gheorghieffii* Forel, 1892i: 309 (m.) BULGARIA.  
Type-material: holotype male.  
Type-locality: Bulgaria: Sliven, 10.viii.1891 (*A. Forel*).  
Type-depository: MHNG.  
Combination in *Stigmatomma*: Emery, in Dalla Torre, 1893: 14.  
**Status as species**: Dalla Torre, 1893: 14; Emery, 1909c: 358; Emery, 1911d: 24.  
Junior synonym of *denticulatum*: Emery, 1915a: 255, Emery, 1916b: 100; Brown, 1960a: 168, 197; Baroni Urbani, 1978a: 42; Bolton, 1995b: 62.
- gracilicornis***. *Stigmatomma denticulatum* var. *gracilicornis* Menozzi, 1936d: 268 (w.) GREECE (Rhodes I.).  
Type-material: 8 syntype workers.  
Type-locality: Greece: Rhodes I., M. Profeta Elia (*C. Menozzi*).  
Type-depository: IEUB.

Junior synonym of *denticulatum*: Brown, 1960a: 168, 197; Baroni Urbani, 1978a: 42; Bolton, 1995b: 62.

**\*groehni.** \**Amblyopone groehni* Dlussky, 2009: 1046, figs. 2a,b (w.) BALTIC AMBER (Eocene).

Combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.

**Status as species:** Dlussky & Rasnitsyn, 2009: 1031; Perkovsky, 2016: 115.

**impressifrons.** *Stigmatomma impressifrons* Emery, 1869b: 13, pl. 1, fig. 8 (w.q.) ITALY.

Type-material: syntype workers, syntype queens (numbers not stated).

Type-locality: Italy: vic. Naples, woods of Capodimonte (C. Emery).

Type-depositories: MHNG, MSNG.

Emery, 1915a: 254 (m.).

Combination in *Amblyopone*: Emery & Forel, 1879: 455;

combination in *Stigmatomma*: Emery, in Dalla Torre, 1893: 14; Emery, 1909c: 357; Yoshimura & Fisher, 2012a: 19.

**Status as species:** Emery, 1875a: 74; Emery, 1875b: 472; Emery, 1878b: 50; Emery & Forel, 1879: 455; André, 1882c: 234 (in key); Forel, 1886e: clxvii; Dalla Torre, 1893: 14; Emery, 1895b: 59; Emery, 1909c: 357; Emery, 1911d: 24; Santschi, 1915a: 55 (in key); Emery, 1915a: 254; Emery, 1916b: 101 (in key); Bondroit, 1918: 80; Müller, 1923b: 24 (in key); Kutter, 1927: 98; Grandi, 1935: 98; Brown, 1960a: 168; Baroni Urbani, 1971c: 10; Baroni Urbani, 1978a: 44 (redescription); Agosti & Collingwood, 1987a: 53; Agosti & Collingwood, 1987b: 264 (in key); Tinaut, 1990d: 192; Mei, 1992a: 412; Atanassov & Dlussky, 1992: 62; Bolton, 1995b: 62; Poldi, *et al.* 1995: 2; Petrov, 2006: 83 (in key); Lapeva-Gjonova, *et al.* 2010: 5; Legakis, 2011: 3; Borowiec, L. & Salata, 2012: 536; Borowiec, L. 2014: 162; Lebas, *et al.* 2016: 106; Salata & Borowiec, 2018c: 48.

Distribution: Bulgaria, Greece, Italy (+ Sicily), Spain.

**irayhady.** *Stigmatomma irayhady* Esteves & Fisher, 2016: 54, figs. 35-42 (w.q.) MADAGASCAR.

Type-material: holotype worker, 4 paratype workers, 2 paratype queens.

Type-locality: holotype Madagascar: Antsiranana, Forêt de Binara, 9.4 km. 235° SW Daraina, 1100 m., -13.26333, 49.6, 5.xii.2003, MW25 sample transect, 5 m., BLF09800, CASENT0042899, montane rainforest, sifted litter (leaf mold, rotten wood) (B.L. Fisher); paratypes with same data but CASENT0042842, -843, -845, -847, -898, and 0797614.

Type-depositories: CASC (holotype); BMNH, CASC, MHNG, NHMB (paratypes).

Distribution: Madagascar.

**janovitsika.** *Stigmatomma janovitsika* Esteves & Fisher, 2016: 66, figs. 44-58 (w.q.m.) MADAGASCAR.

Type-material: holotype worker, 2 paratype workers, 1 paratype queen.

Type-locality: holotype Seychelles: Mahé I., Morne Blanc, 660 m., -4.6574, 55.43325, 10.ii.2010, BLF24151, CASENT0161533, sifted litter (leaf mold, rotten wood) (B.L. Fisher, *et al.*); paratypes with same data but CASENT0161532, 0156022, -023.

Type-depositories: CASC (holotype); CASC, MHNG (paratypes).

Distribution: Seychelles.

**juergi.** *Stigmatomma bruni* r. *juergi* Forel, 1922: 87 (w.) TAIWAN.

Type-material: holotype worker.

Type-locality: Taiwan ("Formosa"): Pilam (H. Sauter).

Type-depository: MHNG.

Subspecies of *bruni*: Wheeler, W.M. 1929g: 58; Chapman & Capco, 1951: 25.

Junior synonym of *bruni*: Brown, 1960a: 168; Bolton, 1995b: 62; Terayama, 2009: 93.

**kangba.** *Amblyopone kangba* Xu & Chu, 2012: 1187, figs. 13-15 (w.) CHINA (Tibet, Yunnan).

Type-material: holotype worker, 4 paratype workers.

- Type-locality: holotype China: Tibet, Zayu County, Zhuwagen Town, Cibaqiao, 1750 m., 30.viii.2010, No. A10-3405, soil sample, *Pinus yunnanensis* forest (X. Liu); paratypes: 2 workers with same data, 2 workers China: Yunnan Prov., Xichou County, Xisa Town, Jiaokui Village, 1480 m., 31.iii.2010, No. A10-1763, *Cyclobalanopsis glaucooides* forest (Z. Xu).  
Type-depository: SFCY.  
Combination in *Stigmatomma*: Yoshimura & Fisher, 2014: 15 (in text).  
Distribution: China.
- liebe.** *Stigmatomma liebe* Esteves & Fisher, 2016: 83, figs. 59-72 (w.q.m.) MADAGASCAR.  
Type-material: holotype worker, 5 paratype workers, 1 paratype queen.  
Type-locality: holotype Madagascar: Fianarantsoa, 8.0 km. NE Ivohibe, 1200 m., -22.42167, 46.89833, 3.xi.1997, MW50 sample transect, 5 m., BLF01753, CASENT0318428, montane rainforest, sifted litter (leaf mold, rotten wood) (B.L. Fisher); paratypes with same data but CASENT0746699 to -702.  
Type-depositories: CASC (holotype); BMNH, CASC, MHNG (paratypes).  
Distribution: Madagascar.
- luyiae.** *Stigmatomma luyiae* Hsu, F. *et al.* 2017: 83, figs. 1-4, 8B (w.) TAIWAN.  
Type-material: holotype worker, 1 paratype worker.  
Type-locality: holotype Taiwan: Nantou County, LFDP (Lienhuachih Forest Dynamics Plot), 10.i.2015, Winkler sample (sifted soil) (F.C. Hsu); paratype with same data.  
Type-depositories: NMNS (holotype); TARI (paratype).  
Distribution: Taiwan.
- luzonicum.** *Stigmatomma (Fulakora) luzonicum* Wheeler, W.M. & Chapman, 1925: 56, pl. 1, fig. 18 (q.) PHILIPPINES (Luzon I.).  
Type-material: holotype queen.  
Type-locality: Philippines: Luzon, Laguna Prov., Los Baños (C.F. Baker).  
Type-depository: MCZC.  
Combination in *Amblyopone*: Brown, 1960a: 168;  
combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.  
**Status as species:** Chapman & Capco, 1951: 24; Brown, 1960a: 168, 196; Baltazar, 1966: 234; Bolton, 1995b: 62; Xu & Chu, 2012: 1186 (in key).  
**Senior synonym of williamsi:** Brown, 1960a: 168, 196; Bolton, 1995b: 62.  
Distribution: Philippines (Luzon, Negros).
- meilianum.** *Amblyopone meiliana* Xu & Chu, 2012: 1190, figs. 33-35 (w.) CHINA (Yunnan).  
Type-material: holotype worker.  
Type-locality: China: Yunnan Prov., Deqin County, Yunling Town, Mingyong Village, E slope of Snow Mt Meili, 3250 m., 10.x.2004, No. A04-536, ground sample, mixed forest (S. Shi).  
Type-depository: SFCY.  
Combination in *Stigmatomma*: Yoshimura & Fisher, 2014: 15 (in text).  
Distribution: China.
- minor** Santschi, 1922; see under **FULAKORA**.
- minutum.** *Stigmatomma minuta* Forel, 1913k: 4 (w.) INDONESIA (Sumatra).  
Type-material: syntype workers (number not stated).  
Type-locality: Indonesia: Sumatra, Soengei Bamban, 9.iv.-14.vi.1912 (von Buttel-Reepen).  
Type-depository: MHNG.  
Combination in *Stigmatomma (Fulakora)*: Mann, 1919: 279;  
combination in *Amblyopone*: Brown, 1960a: 168;  
combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.  
**Status as species:** Chapman & Capco, 1951: 24; Brown, 1960a: 168, 196; Bolton, 1995b: 62; Xu & Chu, 2012: 1183 (in key); Bharti, Guénard, *et al.* 2016: 18.  
Distribution: India, Indonesia (Sumatra).
- montigena.** *Stigmatomma pallipes* subsp. *montigena* Creighton, 1940b: 7, figs. 6, 8 (w.q.) U.S.A. (North Carolina).  
Type-material: holotype ergatoid queen, paratype workers (number not stated).

- Type-locality: holotype U.S.A.: North Carolina, Little Switzerland, nr Spruce Pines (*W.S. Creighton*); paratypes with same data.  
Type-depositories: AMNH (holotype); AMNH, LACM, MCZC (paratypes).  
Subspecies of *pallipes*: Creighton, 1950a: 33; Smith, M.R. 1951a: 782.  
Junior synonym of *pallipes*: Brown, 1949c: 84; Brown, 1960a: 169; Smith, M.R. 1967: 346; Smith, D.R. 1979: 1335; Bolton, 1995b: 62.
- mulanae***. *Bannapone mulanae* Xu, 2000b: 301, figs. 10-12 (q.) CHINA (Yunnan).  
Type-material: holotype queen.  
Type-locality: China: Yunnan Prov., Mengla County, Manzhuang Village, 900 m., 10.iii.1998, No. A98-418, soil sample, semi-evergreen monsoon forest (*T. Liu*).  
Type-depository: SFCY.  
Combination in *Stigmatomma*: Ward & Fisher, 2016: 691.  
**Status as species**: Zhou & Ran, 2010: 102; Guénard & Dunn, 2012: 23 Guénard, *et al.* 2013: 373; Eguchi, *et al.* 2015: 85 (in key); Hsu, F. *et al.* 2017: 91 (in key).  
Distribution: China.
- muticum* Santschi, 1914; see under *XYMMER*.
- noonadan***. *Amblyopone noonadan* Taylor, 1965a: 6, figs. 3, 4 (w.) NEW GUINEA (Papua New Guinea: New Britain I.).  
Type-material: holotype worker, 1 paratype worker.  
Type-locality: holotype Papua New Guinea: New Britain, Yalom, 1000 m., 19.v.1962, Danish Noona Dan Expd. (no collector's name); paratype with same data.  
Type-depositories: ZMUC (holotype); MCZC (paratype).  
Taylor, 1979: 826 (m.).  
Combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.  
**Status as species**: Taylor, 1979: 824 (redescription); Bolton, 1995b: 62; Xu & Chu, 2012: 1182 (in key).  
Distribution: Papua New Guinea.
- normandi***. *Stigmatomma normandi* Santschi, 1915a: 54, fig. 1 (w.) TUNISIA.  
Type-material: syntype workers (number not stated).  
Type-localities: Tunisia: Le Kef (*Normand*), Tunisia: Sousse (*Normand*).  
Type-depository: NHMB.  
Combination in *Amblyopone*: Brown, 1960a: 169;  
combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.  
**Status as species**: Brown, 1960a: 168; Baroni Urbani, 1978a: 45 (redescription); Kugler, J. 1988: 256; Bolton, 1995b: 62; Vonshak, *et al.* 2009: 38; Borowiec, L. 2014: 162.  
Distribution: Isrel, Tunisia.
- octodontatum***. *Amblyopone octodontata* Xu, 2006: 152, figs. 1-7 (w.q.) CHINA (Yunnan).  
Type-material: holotype worker, 4 paratype workers, 1 paratype queen.  
Type-locality: holotype China: Yunnan Prov., Kunming City, Xishan Mountain Forest Park, Nie'ermu, 2150 m., 3.v.2001, No. A00476, *Pinus armandii* forest (*Y. Zhao*); paratypes with same data.  
Type-depository: SFCY.  
Combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.  
**Status as species**: Zhou & Ran, 2010: 102; Guénard & Dunn, 2012: 23; Xu & Chu, 2012: 1186 (in key), 1194.  
Distribution: China.
- ophthalmicum***. *Amblyopone ophthalmica* Baroni Urbani, 1978a: 44, figs. 3, 9 (w.) ISRAEL.  
Type-material: holotype worker, 1 paratype worker.  
Type-locality: holotype Israel: Galilee, Ginosar, 24.v.1973 (*I. Löbl*); paratype with same data.  
Type-depositories: MHNG (holotype); NHMB (paratype).  
Combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.

**Status as species:** Kugler, J. 1988: 256; Bolton, 1995b: 62; Vonshak, *et al.* 2009: 38; Xu & Chu, 2012: 1184 (in key); Borowiec, L. 2014: 162.

Distribution: Israel.

**oregonense.** *Stigmatomma pallipes* subsp. *oregonense* Wheeler, W.M. 1915b: 389 (w.q.) U.S.A. (Oregon, Washington).

Type-material: 13 syntype workers, 4 syntype queens.

Type-localities: U.S.A.: Oregon, Marion (*P.J. Schmitt*), U.S.A.: Washington, Olympia (*T. Kincaid*).

Type-depositories: AMNH, MCZC.

[*Stigmatomma pallipes* subsp. *oregonense* Wheeler, W.M. 1910g: 561. *Nomen nudum*.]

Combination in *Amblyopone*: Brown, 1960a: 169;

combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.

**Subspecies of *pallipes*:** Essig, 1926: 869; Creighton, 1940b: 7; Creighton, 1950a: 34; Smith, M.R. 1951a: 782.

**Status as species:** Brown, 1960a: 169, 183, 192 (in key); Smith, M.R. 1967: 346; Smith, D.R. 1979: 1335; Wheeler, G.C. & Wheeler, J. 1986g: 17 (in key); Ward, 1988: 105; Lattke, 1991c: 6 (in key); Bolton, 1995b: 62; Lacau & Delabie, 2002: 40 (in key); Ward, 2005: 25.

Distribution: Canada, U.S.A.

***pallidipes***, misspelling, see under *pallipes*.

***pallipes*.** *Typhlopone pallipes* Haldeman, 1844: 54 (w.) U.S.A. (no state data).

Type-material: syntype (?) workers.

[Note: no indication of number of specimens is given.]

Type-locality: U.S.A.: (no further data; the only information supplied is “found in old stumps in June”).

Type-depository: unknown (no type-material known to exist).

[Note: Creighton, 1940b: 4, comments that he examined two ‘autotypes’ that had been sent by Haldeman to Harris, which were deposited in the Boston Society of Natural History (now MSBM). While not syntypes, these were probably specimens collected and identified by Haldeman himself.]

[Unjustified emendation of spelling to *pallidipes*: Dalla Torre, 1893: 14; Santschi, 1914a: 429.]

Emery, 1895c: 261 (q.m.); Wheeler, W.M. 1900c: 61 (l.); Wheeler, G.C. & Wheeler, J. 1952a: 114 (l.).

Combination in *Amblyopone*: Haldeman, 1849a: 201 (footnote); Cresson, 1887: 259; Brown, 1960a: 169;

combination in *Stigmatomma (Amblyopone)*: Wheeler, W.M. 1900c: 56;

combination in *Stigmatomma*: Roger, 1863b: 20; Emery, in Dalla Torre, 1893: 14; Yoshimura & Fisher, 2012a: 19.

**Status as species:** Roger, 1863b: 20; Mayr, 1863: 457; Mayr, 1886d: 439; Cresson, 1887: 259; Dalla Torre, 1893: 14; Emery, 1895c: 261; Wheeler, W.M. 1900c: 56; Wheeler, W.M. 1904e: 299; Wheeler, W.M. 1905f: 374; Wheeler, W.M. 1906b: 1; Wheeler, W.M. 1908e: 399; Wheeler, W.M. 1910g: 561; Emery, 1911d: 24; Wheeler, W.M. 1913c: 112; Wheeler, W.M. 1917i: 460; Smith, M.R. 1928b: 243; Smith, M.R. 1930a: 2; Wheeler, W.M. 1932a: 2; Dennis, 1938: 275; Wesson, L.G. & Wesson, R.G. 1940: 90; Creighton, 1940b: 3 (redescription); Brown, 1949c: 84; Creighton, 1950a: 33; Smith, M.R. 1951a: 783; Cole, 1953c: 84; Smith, M.R. 1958c: 110; Whelden, 1958: 1; Brown, 1960a: 169, 183, 192 (in key); Carter, 1962a: 6 (in list); Francoeur & Béique, 1966: 141; Smith, M.R. 1967: 346; Hunt & Snelling, 1975: 20; Francoeur, 1977b: 206; Francoeur, 1979: 34 (redescription); Smith, D.R. 1979: 1335; Traniello, 1982: 65; Wheeler, G.C. & Wheeler, J. 1986g: 17; DuBois & LaBerge, 1988: 135; Ward, 1988: 105; Deyrup, *et al.* 1989: 92; Lattke, 1991c: 6 (in key); Wheeler, G.C., *et al.* 1994: 302; Bolton, 1995b: 62; Lacau &

Delabie, 2002: 40 (in key); Mackay & Mackay, 2002: 31; Deyrup, 2003: 44; Coover, 2005: 23; MacGown & Forster, 2005: 67; Ward, 2005: 25; Ellison, *et al.* 2012: 89; Deyrup, 2017: 13.

**Senior synonym of *arizonense*:** Creighton, 1940b: 3; Creighton, 1950a: 33; Smith, M.R. 1951a: 783; Brown, 1960a: 169; Smith, D.R. 1979: 1335; Bolton, 1995b: 62.

**Senior synonym of *binodosus*:** Emery, 1895c: 261; Wheeler, W.M. 1908e: 400; Emery, 1911d: 24; Wheeler, W.M. 1911f: 159; Creighton, 1940b: 3; Creighton, 1950a: 33; Smith, M.R. 1951a: 783; Brown, 1960a: 169; Francoeur & Béique, 1966: 141; Smith, D.R. 1979: 1335; Bolton, 1995b: 62.

**Senior synonym of *montigena*:** Brown, 1949c: 84; Brown, 1960a: 169; Smith, M.R. 1967: 346; Smith, D.R. 1979: 1335; Bolton, 1995b: 62.

**Senior synonym of *serratum*:** Mayr, 1886d: 439; Emery, 1895c: 261; Emery, 1911d: 24; Creighton, 1950a: 33; Smith, M.R. 1951a: 783; Brown, 1960a: 169; Smith, D.R. 1979: 1335; Bolton, 1995b: 62.

**Senior synonym of *subterranea*:** Brown, 1960a: 169; Smith, M.R. 1967: 346; Smith, D.R. 1979: 1335; Bolton, 1995b: 62.

**Senior synonym of *wheeleri*:** Creighton, 1940b: 3; Creighton, 1950a: 33; Smith, M.R. 1951a: 783; Brown, 1960a: 169; Smith, D.R. 1979: 1335; Bolton, 1995b: 62.

Distribution: Canada, U.S.A.

*paranense* Santschi, 1925; see under ***FULAKORA***.

***pertinax***. *Amblyopone pertinax* Baroni Urbani, 1978a: 48, figs. 13, 14 (w.) INDIA (West Bengal).

Type-material: holotype worker.

Type-locality: India: Darjeeling Dist., Chim Khona, nr Ghum, 2200 m., 28.v.1975 (*W. Wittmer*).

Type-depository: NHMB.

Combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19;

combination in *Bannapone*: Eguchi, *et al.* 2015: 84;

combination in *Stigmatomma*: Ward & Fisher, 2016: 691.

**Status as species:** Bolton, 1995b: 62; Xu, 2001d: 552 (in key); Bharti & Wachkoo, 2011: 590 (in key); Xu & Chu, 2012: 1177 (in key); Eguchi, *et al.* 2015: 85 (in key); Bharti, Guénard, *et al.* 2016: 18; Hsu, F. *et al.* 2017: 91 (in key).

Distribution: India.

***pluto***. *Amblyopone pluto* Gotwald & Lévioux, 1972: 384, figs. 1-26 (w.q.m.) IVORY COAST.

Type-material: holotype worker, paratype workers, paratype queens, paratype males (numbers not stated).

Type-locality: holotype Ivory Coast: Lamto Field Station of the University of Abidjan (ca 50 km. S

Toumodi, 6°13'N, 5°41'W), 16.viii.1965, nest AA 104 (*J. Lévioux*); paratypes: some with same data but others with variant dates and nest numbers. The authors do not provide details but mention "type-nest and 4 additional samples", comprising at least 15 workers, 5 queens and 2 males.]

Type-depositories: MNHN (holotype); BMNH, MCZC, MNHN (paratypes).

Combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.

**Status as species:** Bolton, 1995b: 62.

Distribution: Ivory Coast.

***quadratum***. *Stigmatomma quadratum* Karavaiev, 1935a: 58 (w.) VIETNAM.

Type-material: holotype worker.

Type-locality: Vietnam: Poulo Dama archipelago, Gulf of Siam (= Gulf of Thailand), S Phuquoé I. (= Phu Quoc I.), Nr 5673 (*K. Davydov*).

Type-depository: SIZK.

Combination in *Amblyopone*: Brown, 1960a: 169;

combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.

**Status as species:** Chapman & Capco, 1951: 25; Brown, 1960a: 169; Bolton, 1995b: 62; Xu, 2001d: 552 (in key); Xu & Chu, 2012: 1181 (in key).

Distribution: Vietnam.

***reclinatum***. *Amblyopone reclinata* Mayr, 1879: 667 (w.) INDONESIA (Java).

Type-material: holotype (?) worker.

Type-locality: Indonesia: Java (*S. van Vollenhoven?*).

Type-depository: NHMW.

Stitz, 1925: 110 (m.); Imai, *et al.* 1984: 67 (k.).

Combination in *Stigmatomma*: Emery, in Dalla Torre, 1893: 14;

combination in *Amblyopone*: Brown, 1960a: 169, 193;

combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.

**Status as species:** Dalla Torre, 1893: 14; Forel, 1905c: 3; Emery, 1911d: 24; Forel, 1913k: 4; Stitz, 1925: 110; Karavaiev, 1925a: 75; Chapman & Capco, 1951: 25; Brown, 1960a: 169; Baltazar, 1966: 234; Bolton, 1995b: 62; Jaitrong & Nabhitabhata, 2005: 12; Xu & Chu, 2012: 1179 (in key).

Distribution: Indonesia (Java), Philippines (Binaluan, Luzon, Palawan), Thailand.

***roahady***. *Stigmatomma roahady* Esteves & Fisher, 2016: 103, figs. 73-87 (w.q.m.) MADAGASCAR.

Type-material: holotype worker, 8 paratype workers.

Type-locality: holotype Madagascar: Toamasina, Forêt Ambatovy, 14.3 km. 57° Moramanga, 1075 m., -18.85083, 48.32, 12.iv.2005, BLF11961, CASENT0318421, montane rainforest, ex rotten log (*B.L. Fisher*); paratypes with same data but CASENT0318422 to -424, 0318916 to -918, and 0227519.

Type-depositories: CASC (holotype); CASC, MCZC, MHNG, NHMB, NHMW, MZSP, USNM (paratypes).

Distribution: Madagascar.

***rothneyi***. *Amblyopone (Stigmatomma) rothneyi* Forel, 1900c: 56 (w.) INDIA (West Bengal).

Type-material: lectotype worker (by designation of Brown, 1960a: 193 (in text)).

[Note: Forel gives no indication of the original number of syntypes.]

Type-locality: India: Barrackpore (*Rothney*).

Type-depository: MHNG.

Combination in *Stigmatomma*: Bingham, 1903: 37;

combination in *Amblyopone*: Brown, 1960a: 169;

combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.

**Status as species:** Rothney, 1903: 97; Bingham, 1903: 37; Emery, 1911d: 25; Wheeler, W.M. & Chapman, 1925: 55; Wheeler, W.M. 1928c: 4; Wheeler, W.M. 1930h: 58; Chapman & Capco, 1951: 25; Brown, 1960a: 169, 193; Baltazar, 1966: 234; Bolton, 1995b: 62; Xu, 2001d: 552 (in key); Xu, 2006: 152 (in key); Zhou & Ran, 2010: 102; Bharti & Wachkoo, 2011: 590 (in key); Guénard & Dunn, 2012: 23; Xu & Chu, 2012: 1182 (in key); Yoshimura & Fisher, 2012a: 19; Bharti, Guénard, *et al.* 2016: 18.

Distribution: China, India.

***rubiginoum***. *Amblyopone rubiginous* Wu, J. & Wang, 1992: 1303, fig. 4567 (w.) CHINA (Hunan).

Type-material: holotype worker.

Type-locality: China: Hunan Prov., Liuyang County, vi.1985 (*Jiang Yun*).

Type-depository: CAFB.

Combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.

**Status as species:** Bolton, 1995b: 62; Wu, J. & Wang, 1995: 33; Xu, 2001d: 552 (in key); Xu, 2006: 152 (in key); Zhou & Ran, 2010: 102; Guénard & Dunn, 2012: 23; Xu & Chu, 2012: 1179 (in key).

Distribution: China.

**sakaii.** *Amblyopone sakaii* Terayama, 1989b: 344, figs. 1, 2 (w.) TAIWAN.

Type-material: holotype worker.

Type-locality: Taiwan: Nantou Hsien, Nanshanchi, 21.viii.1987 (*H. Sakai*).

Type-depository: NIAS.

Combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.

**Status as species:** Bolton, 1995b: 62; Onoyama, 1999: 161; Xu, 2001d: 552 (in key); Imai, *et al.* 2003: 219; Lin & Wu, 2003: 67; Xu, 2006: 152 (in key); Terayama, 2009: 93; Zhou & Ran, 2010: 102; Xu & Chu, 2012: 1177 (in key); Bharti & Rilta, 2015b: 510 (in key); Hsu, F. *et al.* 2017: 90 (in key).

Distribution: Japan, Taiwan.

**sakalava.** *Stigmatomma sakalava* Esteves & Fisher, 2016: 159, figs. 88-100 (w.) MADAGASCAR.

Type-material: holotype worker, 5 paratype workers.

Type-locality: holotype Madagascar: Antsiranana, Ampasindava, Andranomatavy Forest, 275 m., -13.6648, 47.98702±500 m., 6.x.2013, BLF31671, CASENT0366766, disturbed dry forest, sifted litter (*B.L. Fisher, et al.*); paratypes with same data but CASENT000366764, -765, and 0318441 to -443.

Type-depositories: CASC (holotype); BMNH, CASC, MHNG, NHMB (paratypes).

Distribution: Madagascar.

**santschii.** *Stigmatomma santschii* Menozzi, 1922a: 347 (w.) SENEGAL.

Type-material: 2 syntype workers.

Type-locality: Senegal: Dakar (no collector's name).

Type-depository: IEUB.

Combination in *Amblyopone*: Brown, 1960a: 169;

combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.

**Status as species:** Brown, 1960a: 169; Bolton, 1995b: 63.

Distribution: Ghana, Nigeria, Senegal.

**scrobiceps.** *Bannapone scrobiceps* Guénard, *et al.* 2013: 373, figs. 2-4 (w.) CHINA (Yunnan).

Type-material: holotype worker, 1 paratype worker.

Type-locality: holotype China: Yunnan Prov., Mengla County, Xishuangbanna Dai Autonomous Prefecture, Xishuangbanna Tropical Botanical Garden, 21°55'07.1''N, 101°16'20.6''E, 550 m., 5.vi.2013, tropical rainforest, leaf litter (*B. Guénard*); paratype with same data (preserved in alcohol).

Type-depositories: USNM (holotype); OIST (paratype).

Combination in *Stigmatomma*: Ward & Fisher, 2016: 691.

**Status as species:** Eguchi, *et al.* 2015: 85 (in key); Liu, C., Guénard, *et al.* 2015: 30; Hsu, F. *et al.* 2017: 91 (in key).

Distribution: China.

**serratum.** *Stigmatomma serratum* Roger, 1859: 251 (w.) U.S.A. (Maryland).

Type-material: holotype worker.

Type-locality: U.S.A.: Maryland, Baltimore (no collector's name).

Type-depository: MNHU.

Status as species: Roger, 1861a: 53; Roger, 1863b: 20; Mayr, 1863: 454.

Junior synonym of *pallipes*: Mayr, 1886d: 439; Emery, 1895c: 261; Emery, 1911d: 24; Creighton, 1950a: 33; Smith, M.R. 1951a: 783; Brown, 1960a: 169; Smith, D.R. 1979: 1335; Bolton, 1995b: 63.

**silvestrii.** *Stigmatomma silvestrii* Wheeler, W.M. 1928d: 97 (w.) JAPAN.

Type-material: 2 syntype workers.

Type-locality: Japan: Nara (Hondo) (*F. Silvestri*).

Type-depository: MCZC (perhaps also in DEUN).

Ogata, 1987: 103 (m.); Terayama, 1989b: 345 (q.); Masuko, 1990b: 221 (l.).

Combination in *Amblyopone*: Brown, 1960a: 169, 197;

combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.

**Status as species:** Azuma, 1951: 86; Chapman & Capco, 1951: 25; Azuma, 1953: 1; Brown, 1960a: 169; Azuma, 1977: 112; Onoyama, 1980: 196; Morisita, *et al.* 1989: 11; Terayama, 1989b: 345; Masuko, 1990b: 221; Masuko, 1993: 35; Tang, J., Li, *et al.* 1995: 24; Bolton, 1995b: 63; Kim, Kim & Kim, 1998: 146; Xu, 2001d: 552 (in key); Imai, *et al.* 2003: 219; Lin & Wu, 2003: 67; Xu, 2006: 152 (in key); Terayama, 2009: 93; Zhou & Ran, 2010: 102; Guénard & Dunn, 2012: 24; Xu & Chu, 2012: 1186 (in key).

Distribution: China, Japan, Korea, Taiwan.

*subterranea*. *Stigmatomma pallipes* subsp. *subterranea* Creighton, 1940b: 8, fig. 4 (w.) U.S.A. (Kansas).

Type-material: holotype worker, 14 paratype workers.

Type-locality: holotype U.S.A.: Kansas, Elmo, 6.ix.1927 (*W.S. Creighton*); paratypes with same data.

Type-depositories: AMNH (holotype); AMNH, LACM, MCZC (paratypes).

**Status as species:** Brown, 1949c: 85.

Subspecies of *pallipes*: Buren, 1944a: 279; Creighton, 1950a: 34; Smith, M.R. 1951a: 783; Smith, M.R. 1958c: 110.

Junior synonym of *pallipes*: Brown, 1960a: 169; Smith, M.R. 1967: 346; Smith, D.R. 1979: 1335; Bolton, 1995b: 63.

**testaceum**. *Amblyopone? testacea* Motschoulsky, 1863: 15 (q.) SRI LANKA. *Nomen dubium*.

Type-material: holotype queen.

Type-locality: Sri Lanka (“Ceylon”): nr Colombo (*V. Motschoulsky*).

Type-depository: unknown (apparently not in ZMUM).

[Note: *Amblyopone? testacea* Motschoulsky, 1863: 15 is not conspecific nor congeneric with *Cryptopone testacea* Emery, 1893a: cclxxv; discussed by Wilson, 1958d: 360; noted under genus *Cryptopone*, and *Cryptopone testacea*.]

Forel, 1900d: 328 (w.).

Combination in *Cryptopone*: Chapman & Capco, 1951: 53 (miscombination);

combination in *Pachycondyla*: Kugler, J. 1988: 256 (miscombination);

combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.

**Status as species:** Smith, F. 1871a: 324; Dalla Torre, 1893: 13; Forel, 1900d: 328; Forel, 1907a: 2; Wheeler, W.M. 1933g: 6; Wilson, 1958d: 360 (*nomen dubium*); Bolton, 1995b: 63.

Distribution: Sri Lanka.

**trigonignathum**. *Amblyopone (Stigmatomma) trigonignatha* Brown, 1949c: 81, fig. 1 (w.) U.S.A. (North Carolina).

Type-material: holotype worker.

Type-locality: U.S.A.: North Carolina, Concord (*D.L. Wray*).

Type-depository: MCZC.

Combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.

**Status as species:** Smith, M.R. 1958c: 110; Brown, 1960a: 169, 185, 192 (in key); Smith, D.R. 1979: 1335; Lattke, 1991c: 6 (in key); Bolton, 1995b: 63; Lacau & Delabie, 2002: 40 (in key).

Distribution: U.S.A.

**trilobum**. *Amblyopone triloba* Xu, 2001d: 552, figs. 11-13 (w.) CHINA (Yunnan).

Type-material: holotype worker.

Type-locality: China: Yunnan Prov., Lushui County, Pianma, 2500 m., 25.iv.1999, No. A99-27, in subalpine moist evergreen broadleaf forest of Gaoligong Mountain Nature Reserve (*Z. Xu*).

Type-depository: SFCY.

Combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.

**Status as species:** Xu, 2006: 152 (in key); Zhou & Ran, 2010: 102; Guénard & Dunn, 2012: 24; Xu & Chu, 2012: 1182 (in key).

- Distribution: China.
- tsyhady.** *Stigmatomma tsyhady* Esteves & Fisher, 2016: 174, figs. 101-114 (w.q.m.)  
MADAGASCAR.  
Type-material: holotype worker, 8 paratype workers.  
Type-locality: holotype Madagascar: Toamasina, Ambatovy, 12.4 km. NE Moramanga, 1080 m., -18.83937, 48.30842, 8.iii.2007, BLF16936, CASENT0121332, montane rainforest, ground forager (*B.L. Fisher, et al.*); paratypes with same data but CASENT0318425 to -427, 0318440, and 0121349.  
Type-depositories: CASC (holotype); BMNH, CASC, MHNG, NHMB (paratypes).  
Distribution: Madagascar.
- wheeleri.* *Stigmatomma pallidipes* [sic] var. *wheeleri* Santschi, 1914a: 429 (w.q.m.) U.S.A. (Connecticut).  
Type-material: syntype workers, syntype queens, syntype males (numbers not stated).  
Type-locality: U.S.A.: Connecticut, Colebrook (*W.M. Wheeler*).  
Type-depository: MCZC, NHMB.  
Subspecies of *pallipes*: Wheeler, W.M. 1916m: 581.  
Junior synonym of *pallipes*: Creighton, 1940b: 3; Creighton, 1950a: 33; Smith, M.R. 1951a: 783; Brown, 1960a: 169; Smith, D.R. 1979: 1335; Bolton, 1995b: 63.
- williamsi.* *Stigmatomma (Fulakora) williamsi* Wheeler, W.M. & Chapman, 1925: 56, pl. 1, fig. 19 (w.q.) PHILIPPINES (Luzon I.).  
Type-material: 13 syntype workers, 2 syntype queens.  
Type-locality: Philippines: Luzon, Mountain Prov., Baguio, 1370 m. (*F.X. Williams*).  
Type-depository: MCZC.  
Status as species: Chapman & Capco, 1951: 24; Baltazar, 1966: 234.  
Junior synonym of *luzonicum*: Brown, 1960a: 168, 196; Bolton, 1995b: 63; Bolton, 1995b: 63.
- xui.** *Stigmatomma xui* Bharti & Rilta, 2015b: 507, figs. 1-3 (w.) INDIA (Sikkim).  
Type-material: holotype worker.  
Type-locality: India: East Sikkim, Rorathang, 27°11'49.91''N, 88°36'12.44''E, 587 m., 12.vi.2012, Winkler (*J.S. Rilta*).  
Type-depository: PUAC.  
**Status as species:** Bharti, Guénard, *et al.* 2016: 18; Hsu, F. *et al.* 2017: 90 (in key).  
Distribution: India.
- zaojun.** *Amblyopone zaojun* Terayama, 2009: 94, figs. 10, 11 (w.) TAIWAN.  
Type-material: holotype worker.  
Type-locality: Taiwan: Taichung Hsien, Deji, 28.vii.-2.viii.1988 (*M. Terayama*).  
Type-depository: NIAS.  
Combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19.  
**Status as species:** Zhou & Ran, 2010: 102; Xu & Chu, 2012: 1186 (in key).  
Distribution: Taiwan.
- zoma.** *Amblyopone zoma* Xu & Chu, 2012: 1189, figs. 28-30 (w.) CHINA (Tibet).  
Type-material: holotype worker.  
Type-locality: China: Tibet, Medog County, Beibing Town, Gangouhe, 740 m., 19.vii.2011, No. A11-3676, soil sample in valley, tropical rainforest (*Z. Xu*).  
Type-depository: SFCY.  
Combination in *Stigmatomma*: Yoshimura & Fisher, 2014: 15 (in text).  
Distribution: China.
- zwaluwenburgi.** *Stigmatomma (Fulakora) zwaluwenburgi* Williams, F.X. 1946: 639 (w.) HAWAII.  
Type-material: holotype worker (lacks head), 1 paratype worker.  
[Note: originally 5 workers but Williams states that "three of the five specimens collected are not now available.]

Type-locality: holotype Hawaii: Oahu. Honolulu, Experiment Station H.S.P.A., ii.1941(*R.H. Van Zwaluwenburg*); paratype with same data but 26.x.1945.

Type-depository: BPBM.

Combination in *Amblyopone*: Brown, 1960a: 169;

combination in *Stigmatomma*: Yoshimura & Fisher, 2012a: 19;

combination in *Bannapone*: Eguchi, *et al.* 2015: 84;

combination in *Stigmatomma*: Ward & Fisher, 2016: 691.

**Status as species:** Brown, 1960a: 169, 213; Wilson & Taylor, 1967: 19; Bolton, 1995b: 63; Onoyama, 1999: 158 (redescription); Framenau & Thomas, 2008: 57; Sarnat & Economo, 2012: 22; Eguchi, *et al.* 2015: 85 (in key).

Distribution: Christmas I., Fiji Is, Hawaii.