





Unidentifiable taxon, *incertae sedis* in Formicidae: Dlussky, Wappler & Wedmann, 2009: 14.

\***pulcher**. \**Gesomyrmex pulcher* Dlussky, Wappler & Wedmann, 2009: 11, figs. 4A, 6G (q.) GERMANY (Messel, Eocene).

**spatulatus**. *Gesomyrmex spatulatus* Cole, 1949: 72, fig. 1 (s.w.) INDIA (Assam).

**Status as species**: Chapman & Capco, 1951: 208; Bolton, 1995b: 207; Bharti, Guénard, et al. 2016: 27.

\**theryi*. \**Dimorphomyrmex theryi* Emery, 1905b: 188, fig. 1 (w.) BALTIC AMBER (Eocene).

[Misspelled as *thyperi* by Keilbach, 1982: 280.]

Status as species: Handlirsch, 1907: 868; Wheeler, W.M. 1910c: 132; Wheeler, W.M. 1915h: 104; Burnham, 1979: 113 (error); Keilbach, 1982: 280 (error).

Junior synonym of \**hoernesi*: Wheeler, W.M. 1929a: 11; Bolton, 1995b: 207; Dlussky, Wappler & Wedmann, 2009: 6.

**tobiasi**. *Gesomyrmex tobiasi* Dubovikoff, 2004b: 220, figs. 1, 2 (q.) VIETNAM.

**Status as species**: Dlussky, Wappler & Wedmann, 2009: 13 (in key).