

Taxonomic review of the *Cataglyphis livida* complex (Hymenoptera, Formicidae), with a description of a new species from Iran

Sebastian Salata^{1,2}, Haniyeh Kiyani³, Kambiz Minaei³, Lech Borowiec¹

1 Department of Biodiversity and Evolutionary Taxonomy, University of Wrocław, Przybyszewskiego 65, 51–148, Wrocław, Poland **2** California Academy of Sciences, 55 Music Concourse Dr, San Francisco, 94118, CA, USA **3** Department of Plant Protection, Faculty of Agriculture, Shiraz University, Shiraz, Iran

Corresponding author: Sebastian Salata (sdsalata@gmail.com)

Academic editor: B.L. Fisher | Received 4 September 2020 | Accepted 7 December 2020 | Published 13 January 2021

<http://zoobank.org/FA1B6869-B8C7-4BC1-B7DC-E3016C5F6CB9>

Citation: Salata S, Kiyani H, Minaei K, Borowiec L (2021) Taxonomic review of the *Cataglyphis livida* complex (Hymenoptera, Formicidae), with a description of a new species from Iran. ZooKeys 1010: 117–131. <https://doi.org/10.3897/zookeys.1010.58348>

Abstract

Cataglyphis fici **sp. nov.**, a member of the *Cataglyphis livida* complex, is described from the Estahban county of south-western Iran. The species is characterized by small body size and yellow to yellowish red body coloration with distinctly infuscated legs. Additionally, the taxonomic status of all known members of the *Cataglyphis livida* complex is updated. *Cataglyphis lutea* Pisarski, 1967, **stat. rev.** is raised to the species level and *Cataglyphis viaticoides* (André, 1881) is proposed as a senior synonym of *Cataglyphis livida bulgarica* Atanassov, 1982, **syn. nov.** and *Cataglyphis albicans mixtus* (Forel, 1895), **syn. nov.** Finally, a provisional key to members of the *livida* complex is provided.

Keywords

Cataglyphis, eastern Mediterranean, key to species, south-western Iran, taxonomy

Introduction

The ant genus *Cataglyphis* Foerster, 1850 currently includes 112 valid species and sub-species (Bolton 2020) distributed mostly in the semideserts and deserts of the Palearctic Region and the sub-Saharan area, India, and Pakistan. *Cataglyphis* species are among

the most common ants occurring in arid and open, Mediterranean-type habitats of North Africa, the Arabian Peninsula, and Central Asia. Less frequently, *Cataglyphis* was recorded from the high altitude, mountain steppes, or forest steppes (Agosti 1990; Radchenko 1997; Brown 2000). Members of the genus are thermophilous, nest in ground and feed mainly on dead insects and other invertebrates. They are characterized by a strictly diurnal activity and are recognized for their superior navigating abilities (Lenoir et al. 2009; Wehner 2020).

The very first worldwide revision of the genus (Santschi 1929) is outdated. The only modern and comprehensive revision of *Cataglyphis* was presented by Agosti (1990) who provided, among others, its redefined species group division. Unfortunately, most of the groups recognized by Agosti remain unrevised and consist of taxa of uncertain status. On the regional level *Cataglyphis* was revised in the following countries: Armenia (Arakelian 1994), Bulgaria (Atanassov and Dlussky 1992), China (Chang and He 2002), Iraq (Pisarski 1965), Kingdom of Saudi Arabia (Collingwood and Agosti 1996), Morocco (Cagniant 2009), Portugal (Collingwood and Prince 1998), and Turkmenistan (Dlussky et al. 1992). There are also available revisions or checklists covering the former European U.S.S.R. (Arnol'di and Dlussky 1978), Iberian Peninsula (Collingwood 1978), Asia (Radchenko 1997, 1998), Central and North Europe (Seifert 2018), and Arabian Peninsula (Collingwood et al. 2011; Sharaf et al. 2015).

Recent publications, presenting descriptions of new species and changes in taxonomic statuses of these already described, proved that the diversity of *Cataglyphis* is underestimated and requires further studies (Radchenko and Paknia 2010; Amor and Ortega 2014; Sharaf et al. 2015; Ionescu and Eyer 2016; Salata and Borowiec 2018). Iran, due to its location and predominance of open and arid habitats, hosts one of the highest numbers of *Cataglyphis* species. So far, there are 35 *Cataglyphis* taxa recorded for this country, but some records need confirmation or correction (Paknia et al. 2008, 2009; Moradloo et al. 2015; Janicki et al. 2016; Rad et al. 2018; Khalili-Moghadam et al. 2021).

Cataglyphis fici, a new species collected in Fars Province of Iran, is a member of the *Cataglyphis albicans* species group sensu Agosti (1990). Its members are characterized by small body size (WL < 3.5 mm), monomorphic workers and colonies lacking distinct major or soldier castes, nodiform petiole with angled dorsal outline and short peduncle, subtly microsculptured and shiny body, and uniformly yellow to black mesosoma. Within the *albicans* species group, *C. fici* is most similar to species listed within the *livida* complex. Agosti (1990) characterized this complex based on uniform, yellow body coloration and included there three species: *C. argentata* (Radoszkowsky), *C. livida* (André), and *C. viaticoides* (André), four subspecies: *C. albicans aurata* Menozzi, *C. albicans fezzanensis* Bernard, *C. albicans mixtus* (Forel), and *C. livida lutea* Pisarski, and four quadrinomial unavailable names: *C. livida lutea agnata* Santschi, *C. livida lutea ambigua* Santschi, *C. livida lutea arabica* (Emery), and *C. livida lutea arenaria* Finzi. Members of the *livida* complex are distributed from Morocco to Indus river including the Arabian Peninsula, and inhabit semi-deserts, deserts, and rocky open areas such as dry hills or coastal cliffs.

The work presented here is a contribution to studies on members of the *C. livida* complex. We list an updated synopsis of members of this complex and provide a provisional key to their identification. Additionally, we describe *Cataglyphis fici* sp. nov., a new member of the *C. livida* complex, based on material recently collected from Iran.

Materials and methods

Investigated specimens were collected in fig orchards located in Estahban city, Fars Province, Iran and are part of the material gathered for a scientific project conducted by the second author. The city is placed 1730 m a.s.l and is characterized by a dry climate, with a yearly precipitation amount of 224 millimeters and summer temperatures frequently exceeding 25.0 °C.

The dominant method was direct sampling (hand collecting). Individual specimens were collected on the ground and preserved in 75% EtOH. Photographs were taken using a Nikon SMZ 1500 stereomicroscope, Nikon D5200 photo camera, and Helicon Focus software. All given label data are in the original spelling, presented in square brackets; a vertical bar (|) separates data on different rows and double vertical bars (||) separate labels. Type specimens' photographs are available online on AntWeb (www.AntWeb.org) and are accessible using the unique CASENT identifying specimen code.

Examined specimens are housed in the following collections:

MNHW Museum of Natural History, University of Wrocław, Poland, in temporary deposit by Department of Biodiversity and Evolutionary Taxonomy, University of Wrocław, Poland;

MHNG Muséum d'Historie Naturelle, Genève, Switzerland.

Measurements:

- HL** head length; measured in a straight line from mid-point of anterior clypeal margin to mid-point of posterior margin in full-face view;
- HW** head width; measured in full-face view at the center of the eyes;
- SL** scape length; maximum straight-line length of scape excluding the basal condylar bulb;
- PW** pronotum width; maximum width of pronotum in dorsal view;
- PRL** propodeum length; measured in lateral view, from metanotal groove to posterior-most point of propodeum;
- PRW** propodeum width; maximum width of propodeum in dorsal view;
- PTH** petiole height; the chord of ventral petiolar profile at node level is the reference line perpendicular to which the maximum height of petiole is measured, measured in lateral view;
- PTW** petiole width; maximum width of the petiolar node in lateral view;

- WL** Weber's length; measured as diagonal length from the anterior end of the neck shield to the posterior margin of the propodeal lobe;
HFL hind femur length; measured on dorsal side from trochanter to apex of femur.

All measurements are given in mm.

Ratios

- CI** cephalic index, HL/HW;
SI scape index, SL/HL;
PI petiole index, PTH/PTW;
FI femur index, HFL/WL.

Synopsis of species of the *Cataglyphis livida* complex

- Cataglyphis arenaria* Finzi, 1940
Cataglyphis argentata (Radoszkowsky, 1876)*
Cataglyphis aurata Menozzi, 1932
Cataglyphis fici sp. nov.
Cataglyphis livida (André, 1881)
Cataglyphis lutea Pisarski, 1967, stat. rev.
Cataglyphis viaticoides (André, 1881)
= *Cataglyphis livida bulgarica* Atanassov, 1982, syn. nov.
= *Cataglyphis albicans mixtus* (Forel, 1895), syn. nov.

Taxonomy

Diagnosis of workers of the *Cataglyphis livida* complex

Small body size (WL < 3.0 mm); colonies with monomorphic workers, lacking distinct major or soldier castes; petiole nodiform with angled dorsal outline and short peduncle; body subtly microsculptured and shiny; body uniformly yellow to red (never brown to black) or bicolored with entirely to partially black gaster.

Distribution: from Morocco to Asia Minor and the Middle East, in semideserts, deserts and rocky open areas such as dry hills or coastal cliffs.

Note 1. Agosti (1990), as the first one, noticed that the data from labels of type specimens of *C. viaticoides* did not correspond with the original description of the species. This problem was later investigated and clarified by Bračko et al. (2016). Based on evidence gathered and discussed by the authors, the definition of the *livida* complex proposed by Agosti (1990) was modified to accommodate *C. viaticoides*.

Note 2. Based on its description, *C. albicans fezzanensis* Bernard is characterized by the presence of polymorphic worker caste, and additional study on the type specimen

* Due to ambiguous description of this species it is tentatively assigned to the *livida* complex.

indicated that its body sculpture is stronger and less shiny than in other members of the *livida* complex. Based on this data, we decided not to list this species as a member of this complex.

Provisional key to the *Cataglyphis livida* complex

- 1 At least mid and hind legs infuscated. Iran..... *C. fici*
- Legs in the same coloration as mesosoma..... 2
- 2 Head and mesosoma uniformly yellowish red to reddish yellow, gaster entirely or mostly black, Balkans to Asia Minor..... *C. viaticoides*
- Gaster in the same coloration as the rest of body (yellow to red) or its apex slightly infuscated 3
- 3 Mesosoma and head without layer of silvery hair, northeastern Mediterranean to Middle East *C. lutea*
- At least mesosoma with a layer of silvery hair, Morocco to Asia Minor..... 4
- 4 Mesosoma and posterior head with thick layer of silvery hair, North Africa....
..... *C. arenaria*, *C. argentata*, *C. aurata**
- A layer of silvery hair limited to mesosoma, Asia Minor..... *C. livida*

Review of species

Cataglyphis fici sp. nov.

<http://zoobank.org/81413366-4DA0-44D9-BF48-3ACD395E86F4>


Figs 1–6

Type material. *Holotype*: IRAN •worker, Fars, Estahban, 29.1331/54.389, 1730 m a.s.l., 16 Aug. 2019, H. Kiyani leg., LBC-IR00179, CASENT6006519 (MNHW); *paratypes*: 5 workers, the same data as holotype, CASENT6006520–CASENT6006524 (MNHW, MHNG); *paratype*: worker: the same data as holotype except LBC-IR00180, CASENT6006525 (MNHW); *paratype*: worker, the same data as holotype except 6 Sep. 2018 and LBC-IR00182, CASENT6006526 (MNHW).

Holotype worker labels: IRAN, Fars, 1730 m | Estahban | 29.1331 / 54.389 | 16 VIII 2019, H. Kiyani || Collection L. Borowiec | Formicidae | LBC-IR00179 || CASENT6006519.

Diagnosis. *Cataglyphis fici* is a member of the *Cataglyphis albicans* group and can be separated from all species clustered in the *cinnamomea* and *fortis* complexes and most of members of the *albicans* complex by yellow to yellowish red body coloration; while other species have body completely brown to black. From bicolored species of the *albicans* complex it differs in smaller body size (WL < 3 mm) and presence of infuscated to


* Note. The taxonomic status of *C. arenaria*, *C. argentata*, and *C. aurata* is unclear and requires further studies. Due to the lack of distinct morphological differences between types of these taxa and their ambiguous descriptions, we decided to list all three species in one couplet. For more details, see comments in the species accounts.



Figures 1, 2. Holotype worker of *Cataglyphis fici* sp. nov. **1** dorsal **2** lateral. Scale bars: 1 mm.

yellowish brown mid and hind legs. However, *C. fici* is most similar to species included in the *C. livida* complex and can be easily separated based on at least mid and hind legs partly to mostly infuscated to yellowish brown. In contrast, all remaining members of the *C. livida* complex have legs uniformly colored and always in the same shade as mesosoma and head. Additionally, *C. fici* differs from *C. lutea*, *C. arenaria*, *C. argentata*, *C. aurata*, and *C. livida* in gaster darker than head and mesosoma sometimes infuscated.

Description. Worker (n = 8): **Measurements.** HL: 1.405 (1.29–1.54); HW: 1.325 (1.23–1.43); SL: 1.483 (1.36–1.59); PW: 0.897 (0.84–0.97); PRL: 0.737



3




4

Figures 3, 4. Holotype worker of *Cataglyphis fici* sp. nov. **3** head **4** petiole Scale bars: 0.5 mm (**4**), 1 mm (**3**).

(0.68–0.79); PRW: 0.595 (0.56–0.65); PTH: 0.383 (0.32–0.654); PTW: 0.360 (0.34–0.38); WL: 1.933 (1.81–2.09); HFL: 2.038 (1.83–2.24); CI: 1.060 (1.038–1.077); SI: 1.056 (1.032–1.072); PI: 1.065 (0.889–1.158); FI: 1.054 (1.036–1.111).

Color. Head, mesosoma and petiolar scale from yellow to yellowish red, in the darkest specimens sides of mesonotum and propodeum indistinctly infuscated, gaster in the palest specimens mostly yellow with infuscated two apical sternites (Fig. 6), in dark specimens yellowish at base then gradually infuscated apically, with dark brown apical tergites and sternites (Fig. 5). Forelegs in the palest specimens completely yellow, mid and hind legs with brown femora and tibiae and yellowish tarsi, in dark specimens all legs at least partly infuscated. Usually fore coxa mostly brown with reddish spots of



Figures 5, 6. Paratype workers of *Cataglyphis fici* sp. nov. **5** the darkest form **6** the palest form Scale bar: 1 mm.

diffused borders laterally, fore femora mostly brown with yellowish apices, fore tibiae yellowish brown and fore tarsi yellowish; mid and hind femora dark brown, tibiae yellowish brown to brown, tarsi yellowish to yellowish brown (Figs 2, 5). Antennal scape yellow, funicles in the palest specimens slightly darker than scape, yellowish basally and yellowish brown apically, in dark specimens only first segment of funiculus yellowish, remaining segments gradually yellowish brown to dark brown. **Head.** Subrectangular, approximately $1.05 \times$ as long as wide, sides below eyes almost parallel, above eyes gently convex, occipital margin convex (Fig. 3). Anterior margin of the clypeus convex, with small median emargination, clypeal anterior margin with a row of short, white setae and additional six long, white setae, the longest as long as $0.7 \times$ length

of clypeus. Whole surface of clypeus densely microreticulate with shiny background, covered with very sparse and short, adpressed hairs. Eyes large, oval, approximately $1.4 \times$ as long as wide. Frontal carinae short, not extending beyond frontal lobes, interocular area without shiny line or carina and with a pair of long white setae. Antennal fossa shallow, microreticulate with shiny background. Whole head surface finely microreticulate with shiny background, occipital part of the head and are behind eyes with reduced sculpture and shinier, covered with extremely sparse, indistinct, adpressed hairs. Ocellar region with a pair of moderately long white setae, occipital part of head with 2–6 long, white erect setae, underside only with a pair of long, white setae close to lateral margin of head. Antennal scape moderately long, in frontal view almost straight, approximately $1.1 \times$ as long as length of the head; thin, in apex only slightly and gradually widened; its base without tooth. Funiculus long, first funicle segment elongated, approximately $0.8 \times$ as long as segments II and III combined, and $1.7 \times$ as long as segment II (Fig. 3). Surface of scape densely microsculptured; shiny to indistinctly opalescent; covered with strong, moderately dense, decumbent setae. Mandibles rounded, only in basal part smooth and shiny, apical $\frac{3}{4}$ with deep grooves, surface shiny with several long yellow setae, masticatory margin with four large teeth. **Mesosoma.** Long, $2.1 \times$ as long as wide; metanotal groove shallow (Fig. 2). Pronotum convex on sides (Fig. 1). In lateral view promesonotum slightly arched in profile; propodeum positioned lower than promesonotum, moderately convex in lateral view; propodeal spiracle strongly elongated and slit-shaped, approximately $4.2 \times$ as long as wide (Fig. 2). Whole mesosoma opalescent, with dense, fine microreticulation and shiny background; covered with extremely sparse and short adpressed microsetae, on sides of pro- and mesonotum appears almost hairless, only anterior part of pronotum, posterior angles of mesonotum and propodeum with sparse setosity. Pro- and mesonotum without erect setae, propodeum without erect setae or in its posterior part with one or two short, white, erect setae. **Scale.** In form of a short cuneiform node, in lateral view almost trapezoidal with very short peduncle. Anterior face close to base distinctly convex, posterior face slightly concave, top of scale in lateral view obtusely rounded, without erect setae (Fig. 4). In anterior and posterior view top margin of scale without emargination. Surface of petiole distinctly microreticulate and shiny. **Gaster.** With fine transverse microreticulation and striation and very shiny background. Whole surface of gaster with hardly visible, extremely short, sparse, adpressed microsetae, tergites I and II without erect setae, tergites III and IV with a pair of long white setae centrally, each gastral sternite with 2–4 long, white, erect setae. **Legs.** Dorsal and lateral surface of femora and tibiae covered with sparse, white adpressed setae. Ventral surface of femora and tibiae with rows of elongate, white, erect spiniform setae.

Biology. Little known, workers were collected on the ground in fig orchard at altitude 1730 m.

Etymology. The species name *fici* is a noun in the genitive case named after the generic name of the fig tree, *Ficus* sp., the dominant plant in the type locality of this ant species.

Cataglyphis arenaria Finzi, 1940

Cataglyphis (*Cataglyphis*) *albicans* var. *arenaria* Finzi, 1940: 164 [first available use of *Myrmecocystus albicans lividus arenaria* Forel, 1909: 384].

Status as species: Collingwood and Agosti (1996): 378.

Syntype worker, Biskra, Algeria (MHNG) [Syntype worker images examined, AntWeb, CASENT0911101, photographs by Alexandra Westrich, available on AntWeb.org].

Diagnosis. Whole body yellow, only gaster sometimes with indistinctly infuscated apex; mesosoma, posterior part of the head and coxae covered with a layer of silvery hair.

Distribution. North Africa region, from Mauritania to Jordan. Based on photographs available on AntWeb specimens from Arabian Peninsula probably refer to *C. lutea*.

Note. *Cataglyphis arenaria* was separated from *C. livida* and *C. viaticoides* based on the presence of a thick layer of silvery hair on mesosoma and posterior part of the head. Two years after the original description of *C. arenaria*, Karavaiev (1911) described *Myrmecocystus albicans* ssp. *lividus* var. *aurata*, which was later validated by Menozzi (1932) as trinominal. The latter species also was separated from *C. livida* based on presence of silvery hair on its body. Probably, Karavaiev was unaware of the existence of *C. arenaria* during his work on *C. aurata*. Study of type specimens and descriptions of both *C. arenaria* and *C. aurata* did not provide any characters useful in separating these two species. Thus, we conclude that both taxa could be conspecific (see also note in *Cataglyphis argentata* (Radoszkowsky, 1876) and *C. aurata* Menozzi, 1932). However, this hypothesis requires verification based on larger material collected from the whole area of their distribution, supported with studies on male genitalia, and genetic analyses.

Cataglyphis argentata (Radoszkowsky, 1876)

Camponotus argentata Radoszkowsky, 1876: 140.

Cataglyphis argentata: Dalla Torre (1893): 217.

Type specimens. Unavailable.

Diagnosis. Whole body yellow, only gaster sometimes with indistinctly infuscated apex; mesosoma, body covered with a layer of silvery hair.

Distribution. Egypt.

Note. Type specimens of this species are considered lost and, as suggested by Agosti (1990), due to ambiguous description of this species, its assignation to the *livida* complex is tentative. The silvery hair mentioned in the description can suggest an affiliation of *C. argentata* with the *bombycina* or *laevior* complexes. If *C. argentata* is a member of the *livida* complex, then its description could indicate that it is probably conspecific with two other North African taxa: *C. arenaria* and *C. aurata*. If this assumption is correct, then the name *C. argentata* has priority over *C. arenaria* and *C. aurata*.

***Cataglyphis aurata* Menozzi, 1932**

Cataglyphis (*Cataglyphis*) *albicans aurata* Menozzi, 1932: 95 [first available use of *Myrmecocystus albicans* ssp. *lividus* var. *auratus* Karavaiev, 1911: 10].

Syntype worker, Assuan, Egypt (MHNG) [syntype worker images examined, AntWeb, CASENT0911100, photos by Zach Lieberman, available on AntWeb.org].

Diagnosis. Whole body yellow, only gaster sometimes with indistinctly infuscated apex; mesosoma, posterior head and coxa covered with a layer of silvery hair.

Distribution. North Africa. Probably records from Asia Minor refer to *Cataglyphis lutea*.

Note. *Cataglyphis aurata* was separated from *C. livida* based on the presence of a thick layer of silvery hair on its body. Probably, Karavaiev, during his work on *C. aurata*, was unaware of the existence of *C. arenaria*, another species described from the North African region characterized by the same feature. Study on type specimens and descriptions of both *C. arenaria* and *C. aurata* did not provide any characters useful separating these two species. Thus, we conclude that they could be conspecific. However, this hypothesis requires verification based on larger material collected from the whole area of their distribution, supported with studies on male genitalia, and genetic analyses. See also note in *C. argentata*.

***Cataglyphis livida* (André, 1881)**

Myrmecocystus albicans var. *lividus* André, 1881: 58.

Status as species: Arnol'di (1964): 1810.

Syntype workers, Jaffa, Israel (MHNG) [syntype workers images examined, AntWeb, CASENT0911099 and CASENT0912207, photographs by Zach Lieberman and Will Ericson, available on AntWeb.org].

Diagnosis. Whole body yellow, only gaster sometimes with indistinctly infuscated apex; mesosoma and coxa covered with a layer of silvery hair.

Distribution. Unknown. Due to mislabeling of type specimens of *C. livida* and *C. viaticoides*, both species were wrongly interpreted, and most of their historic records require verification. Based on available material, we can confirm its presence in Egypt, coastal parts of Israel, Syria, and Antalya Province in Turkey.

Note. A study on type specimens of *C. livida* revealed that this species could be easily separated from most members of the *livida* complex based on the presence of a layer of silvery hair on propodeum and katapisternum, and lack of these on posterior head. Lack of comment on this feature in the original description combined with mislabeling of type specimens (see Bračko et al. 2016) caused confusion, leading to the long-lasting misinterpretation of *C. livida*.

***Cataglyphis lutea* Pisarski, 1967, stat. rev.**

Cataglyphis livida subsp. *lutea* Pisarski, 1967: 418 [first available use of *Myrmecocystus albicans viaticoides lutea* Emery, 1906: 53].

Junior synonym of *Cataglyphis livida*: Radchenko, 1997: 428.

Syntype worker, Shiraz, Iran (MSNG) [Syntype worker images examined, AntWeb, CASENT0905718, photographs by Will Ericson, available on AntWeb.org].

Diagnosis. Whole body yellow, only gaster sometimes with indistinctly infuscated apex; body never with a layer of silvery hair.

Distribution. Species known from Arabian Peninsula east to Afghanistan.

Note. *Cataglyphis lutea* was described from Shiraz, Fars Province in Iran as an unavailable quadrinomial name (Emery 1906), later validated by Pisarski (1967) as a subspecies of *C. livida*, and finally considered as its junior synonym (Radchenko 1997). A study on type specimen revealed that *C. lutea* distinctly differs from *C. livida* in lack of a layer of silvery hair on mesosoma, and its distribution does not overlap with confirmed records of *C. livida*. Thus, we decided to raise it to the species status. AntCat resources indicated that, except type locality, *C. lutea* is also known from Aran va Bidgol, Maranjab, Iran (CDA000106) and Saudi Arabia (CASENT0906455).

***Cataglyphis viaticoides* (André, 1881)**

Myrmecocystus albicans var. *viaticoides* André, 1881: 57.

Syntype worker, Beirut, Lebanon (MNHN) [syntype worker images examined, AntWeb, CASENT0912236, photographs by Zach Lieberman, available on AntWeb.org].

= *Cataglyphis livida bulgarica* subsp. *bulgarica* Atanassov, 1982: 213, syn. nov.

Type specimens unavailable.

= *Myrmecocystus albicans* var. *mixtus* Forel, 1895: 229, syn. nov.

Syntype worker, Edirne, Turkey (MHNG) [syntype worker images examined, AntWeb, CASENT0911104, photographs by Zach Lieberman, available on AntWeb.org].

Diagnosis. Head and mesosoma uniformly yellowish red to reddish yellow, gaster entirely or mostly dark; thin layer of silvery hair limited to propodeum.

Distribution. Balkans and Asia Minor.

Note. Radchenko (1997), based on confusion related to the type labels of *C. livida* and *C. viaticoides* (see Bračko et al. 2016), considered *C. livida bulgarica* and *C. albicans mixtus* as junior synonyms of *C. livida*. Results presented by Bračko et al. (2016) clarified that the only member of the *livida* complex with entirely or mostly black gaster is *C. viaticoides* and thus *Cataglyphis livida bulgarica* Atanassov, 1982 and *Cataglyphis albicans mixtus* (Forel, 1895) should be considered as its junior synonyms.

Acknowledgements

Haniyeh Kiyani and Kambiz Minaei thank Shiraz University, Iran for supporting their scientific project focused on studies on ant diversity in fig orchards near Estabhan. The authors would like to thank Brian L. Fisher, Donat Agosti, and one anonymous reviewer for reviewing and improving a previous version of this manuscript.

References

- Agosti D (1990) Review and reclassification of *Cataglyphis* (Hymenoptera, Formicidae). *Journal of Natural History* 24: 1457–1506. <https://doi.org/10.1080/00222939000770851>
- Amor F, Ortega P (2014) *Cataglyphis tartessica* sp. n., a new ant species (Hymenoptera: Formicidae) in south-western Spain. *Myrmecological News* 19: 125–132. <https://doi.org/10.5281/zenodo.16346>
- André E (1881) Catalogue raisonné des Formicides provenant du voyage en Orient de M. Abeille de Perrin et description des espèces nouvelles. *Annales de la Société Entomologique de France* (6)1: 53–78. <https://doi.org/10.5281/zenodo.27096>
- Arakelian GR (1994) Fauna of The Republic of Armenia. Hymenopterous insects. Ants (Formicidae). Erevan: Gitutium, 153 pp. [In Russian]
- Arnol'di KV (1964) Higher and specialized representatives of the ant genus *Cataglyphis* (Hymenoptera, Formicidae) in the fauna of the USSR. *Zoologicheskii Zhurnal* 43: 1800–1815. [In Russian.] <https://antcat.org/references/122339>
- Arnol'di KV, Dlussky GM (1978) Superfam. Formicoidea. 1. Fam. Formicidae – ants. In: Medvedev GS (Ed.) Keys to the insects of the European part of the USSR (Vol. 3). Hymenoptera. Part 1. *Opredeliteli Faune SSSR* 119: 519–556. [In Russian]
- Atanassov N, Dlussky GM (1992) Fauna of Bulgaria. Hymenoptera, Formicidae. *Fauna of Bulgaria* 22: 1–310.
- Bolton B (2020) An online catalog of the ants of the world. <https://antcat.org>. [accessed 25 August 2020]
- Bračko G, Kiran K, Karaman C, Salata S, Borowiec L (2016) Survey of the ants (Hymenoptera: Formicidae) of the Greek Thrace. *Biodiversity Data Journal* 4: e7945. <https://doi.org/10.3897/BDJ.4.e7945>
- Brown Jr WL (2000) Diversity of ants. In: Agosti D, Majer J, Alonso E, Schultz TR (Eds) *Ants: Standard Methods for Measuring and Monitoring Biodiversity*. Biological diversity handbook series. Smithsonian Institution Press, Washington, 280 pp.
- Cagniant H (2009) Le genre *Cataglyphis* Foerster, 1850 au Maroc (Hyménoptères Formicidae). *Orsis* 24: 41–71. <https://ddd.uab.cat/record/54485>
- Chang YD, He DH (2002) Three new species of the genus *Cataglyphis* Foerster from Northwest China (Hymenoptera: Formicidae: Formicinae). *Zoological Research* 23: 61–64. [In Chinese] <http://www.zoores.ac.cn/en/article/id/826>

- Collingwood CA (1978) A provisional list of Iberian Formicidae with a key to the worker caste (Hym. Aculeata). EOS. Revista Española de Entomología 52: 65–95. <https://doi.org/10.5281/zenodo.26690>
- Collingwood CA, Agosti D (1996) Formicidae (Insecta: Hymenoptera) of Saudi Arabia (Part 2). Fauna of Saudi Arabia 15: 300–385. <https://antcat.org/references/130931>
- Collingwood C, Prince A (1998) A guide to ants of continental Portugal (Hymenoptera: Formicidae). Boletim da Sociedade Portuguesa de Entomologia. Suplemento 5: 1–49. <https://doi.org/10.5281/zenodo.1244612>
- Collingwood C, Agosti D, Sharaf MR, van Harten A (2011) Order Hymenoptera, family Formicidae. Arthropod fauna of the UAE 4: 405–474. <https://doi.org/10.5281/zenodo.1168586>
- Dalla Torre KW (1893) Catalogus Hymenopterorum Hucusque Descriptorum Systematicus et Synonymicus (Vol. 7). Formicidae (Heterogyna). W. Engelmann, Leipzig, 289 pp. <https://www.biodiversitylibrary.org/page/10937376>
- Dlussky GM, Soyunov OS, Zabelin SI (1990) Ants of Turkmenistan. Ylym Press, Ashkabad, 273 pp. [In Russian]
- Emery C (1906) Rassegna critica delle specie paleartiche del genere *Myrmecocystus*. Memorie della Reale Accademia delle Scienze dell'Istituto di Bologna (6)3: 47–61. <https://doi.org/10.5281/zenodo.25506>
- Finzi B (1940) Formiche della Libia. Memorie della Società Entomologica Italiana 18 (1939): 155–166. <https://doi.org/10.5281/zenodo.26322>
- Forel A (1895) Südpalaearktische Ameisen. Mitteilungen der Schweizerischen Entomologischen Gesellschaft 9: 227–234. <https://doi.org/10.5281/zenodo.14245>
- Forel A (1909) Études myrmécologiques en 1909. Fourmis de Barbarie et de Ceylan. Nidification des Polyrhachis. Bulletin de la Société Vaudoise des Sciences Naturelles 45: 369–407. <https://doi.org/10.5281/zenodo.14425>
- Ionescu A, Eyer P-A (2015) Notes on *Cataglyphis* Foerster, 1850 of the *bicolor* species-group in Israel, with description of a new species (Hymenoptera: Formicidae). Israel Journal of Entomology 46: 109–131. <https://doi.org/10.5281/zenodo.221456>
- Janicki J, Narula N, Ziegler M, Guénard B, Economo EP (2016) Visualizing and interacting with large-volume biodiversity data using client-server web-mapping applications: The design and implementation of antmaps.org. Ecological Informatics 32: 185–193. <https://doi.org/10.1016/j.ecoinf.2016.02.006>
- Karavaiev V (1911) Ameisen aus Aegypten und dem Sudan. Russkoe Entomologicheskoe Obozrenie 11: 1–12. <https://doi.org/10.5281/zenodo.25954>
- Khalili-Moghadam A, Salata S, Borowiec L (2021) Three new species of *Cataglyphis* (Hymenoptera: Formicidae) from Chaharmahal va Bakhtiari Province of Iran. ZooKeys 1009: 1–28. <https://doi.org/10.3897/zookeys.1009.59205>
- Lenoir A, Aron S, Cerda X, Hefetz A (2009) *Cataglyphis* desert ants: a good model for evolutionary biology in Darwin's anniversary year—a review. Israel Journal of Entomology 39: 1–32. <http://digital.csic.es/bitstream/10261/65135/1/israel.pdf>
- Menozi C (1932) Missione scientifica del Prof. E. Zavattari nel Fezzan (1931). Hymenoptera-Formicidae. Bollettino della Società Entomologica Italiana 64: 93–95. <https://doi.org/10.5281/zenodo.25776>

- Moradloo S, Fard RN, Rad SP, Taylor B (2015) Records of ants (Hymenoptera: Formicidae) from Northern Iran. *Zoology in the Middle East* 2015: 1–6. <https://doi.org/10.1080/09397140.2015.1020611>
- Paknia O, Radchenko A, Alipanah H, Pfeiffer M (2008) A preliminary check list of the ants (Hymenoptera: Formicidae) of Iran. *Myrmecological News* 151–159. https://myrmecologicalnews.org/cms/index.php?option=com_content&view=category&id=254&Itemid=355
- Paknia O, Radchenko A, Pfeiffer M (2009) New records of ants (Hymenoptera, Formicidae) from Iran. *Asian Myrmecology* 3: 29–38. <http://doi.org/10.20362/am.003005>
- Pisarski B (1965) Les fourmis du genre *Cataglyphis* Foerst. en Irak (Hymenoptera, Formicidae). *Bulletin de l'Académie Polonaise des Sciences. Série des Sciences Biologiques* 13: 417–422. <https://antcat.org/references/127818>
- Pisarski B (1967) Fourmis (Hymenoptera: Formicidae) d'Afghanistan récoltées par M. Dr. K. Lindberg. *Annales Zoologici (Warsaw)* 24: 375–425. <https://doi.org/10.5281/zenodo.25692>
- Rad SP, Taylor B, Torabi R, Aram E, Abolfathi, Afshari, Borjali F, Ghatei M, Hediary F, Jazini F, Kiah VH, Mahmoudi Z, Sararyan F, Seiri M (2018) Further records of ants (Hymenoptera: Formicidae) from Iran. *Zoology in the Middle East* 2015: 1–15. <https://doi.org/10.1080/09397140.2018.1442301>
- Radchenko AG (1997) Review of ants of the genus *Cataglyphis* Foerster (Hymenoptera, Formicidae) of Asia. *Entomologicheskoe Obozrenie* 76: 424–442. [In Russian.]
- Radchenko AG (1998) A key to the ants of the genus *Cataglyphis* Foerster (Hymenoptera, Formicidae) of Asia. *Entomologicheskoe Obozrenie* 77: 502–508. [In Russian]
- Radchenko AG, Paknia O (2010) Two new species of the genus *Cataglyphis* Foerster, 1850 (Hymenoptera: Formicidae) from Iran. *Annales Zoologici* 60: 69–76. <https://doi.org/10.3161/000345410X499533>
- Radoszkowsky O (1876) Comte-rendu des hyménoptères recueillis en Égypte et Abyssinie en 1873. *Trudy Russkago Entomologicheskago Obshchestva* 12: 111–150. <https://rcin.org.pl/ibd/dlibra/publication/142800/edition/114697>
- Salata S, Borowiec L (2018) Taxonomic and faunistic notes on Greek ants (Hymenoptera: Formicidae). *Annals of the Upper Silesian Museum in Bytom Entomology* 27(online 008): 1–51. <http://doi.org/10.5281/zenodo.2199191>
- Santschi F (1929) Étude sur les *Cataglyphis*. *Revue Suisse de Zoologie* 36: 25–70. <https://www.biodiversitylibrary.org/page/43721762>
- Seifert B (2018) *The Ants of Central and North Europe*. Iutra Verlags- und Vertriebsgesellschaft, Tauer, 407 pp.
- Sharaf MR, Collingwood CA, Aldawood AS (2015) Notes on the ant genus *Cataglyphis* Foerster, 1850 (Hymenoptera, Formicidae) in the Arabian Peninsula with description of a new species and a key to species of the *C. pallida*-group. *ZooKeys* 545: 101–117. <https://doi.org/10.3897/zookeys.545.6308>
- Wehner R (2020) *Desert Navigator. The Journey of an Ant*. The Belknap Press of Harvard University Press, Cambridge, Massachusetts and London, 392 pp. <https://doi.org/10.4159/9780674247918>