

TYPHLOMYRMEX

bruchi. *Typhlomyrmex bruchi* Santschi, 1922d: 242. *Nomen nudum*.

[Note: the specimens were later described by Santschi, 1923c: 245, as *Prionopelta bruchi*: Lacau *et al.* 2005: 39.]

clavicornis. *Typhlomyrmex clavicornis* Emery, 1906c: 112 (footnote) (q.) BOLIVIA.

Brown, 1965c: 71 (w.).

Status as species: Emery, 1911d: 34; Wheeler, W.M. 1925a: 3 (in key); Brown, 1965c: 69; Kempf, 1972a: 256; Bolton, 1995b: 422; Wild, 2007b: 27; Feitosa, 2015c: 98.

Senior synonym of *divergens*: Brown, 1965c: 69; Kempf, 1972a: 256; Bolton, 1995b: 422.

Senior synonym of *richardsi*: Brown, 1965c: 71; Kempf, 1972a: 256; Bolton, 1995b: 422.

divergens. *Typhlomyrmex clavicornis* var. *divergens* Forel, 1906d: 248 (q.m.) PARAGUAY.

Subspecies of *clavicornis*: Emery, 1911d: 34; Wheeler, W.M. 1925a: 4 (in key).

Junior synonym of *clavicornis*: Brown, 1965c: 69; Kempf, 1972a: 256; Bolton, 1995b: 422.

foreli. *Typhlomyrmex foreli* Santschi, 1925b: 6 (q.) BRAZIL (Paraná).

Status as species: Brown, 1965c: 76; Kempf, 1972a: 256; Bolton, 1995b: 422; Feitosa, 2015c: 98.

major. *Typhlomyrmex pusillus* st. *major* Santschi, 1923c: 246 (w.) BRAZIL (Santa Catarina).

Brown, 1965c: 72 (q.).

Subspecies of *pusillus*: Wheeler, W.M. 1925a: 4 (in key).

Status as species: Brown, 1965c: 71; Kempf, 1972a: 256; Bolton, 1995b: 422; Arias-Penna, 2008b: 102; Lacau, *et al.* 2008: 252 (redescription); Feitosa, 2015c: 98; Guénard & Economo, 2015: 226; Feitosa & Prada-Achiardi, 2019: 678.

manco. *Typhlomyrmex robustus* subsp. *manco* Wheeler, W.M. 1925a: 2 (w.m.) BOLIVIA, PERU.

Junior synonym of *rogenhoferi*: Brown, 1957e: 229; Brown, 1965c: 74; Kempf, 1972a: 256; Bolton, 1995b: 422.

meire. *Typhlomyrmex meire* Lacau, Villemant & Delabie, 2004: 5, figs. 1 – 5 (w.q.m.) BRAZIL (Bahia).

Status as species: Feitosa, 2015c: 98.

prolatus. *Typhlomyrmex prolatus* Brown, 1965c: 72, fig. 6 (q.) COSTA RICA.

Status as species: Kempf, 1972a: 256; Bolton, 1995b: 422.

pusillus. *Typhlomyrmex pusillus* Emery, 1894c: 141, pl. 1, fig. 2 (w.) BOLIVIA.

Emery, 1906c: 111 (q.); Wheeler, G.C. & Wheeler, J. 1964b: 447 (l.).

Status as species: Emery, 1906c: 111; Emery, 1911d: 34; Santschi, 1912e: 521; Bruch, 1914: 212; Gallardo, 1918b: 15; Santschi, 1923c: 246; Wheeler, W.M. 1925a: 4 (in key); Santschi, 1929d: 276; Menozzi, 1931b: 261; Kusnezov, 1953b: 336; Kempf, 1961b: 493; Brown, 1965c: 73; Kempf, 1972a: 256; Brandão, 1991: 383; Bolton, 1995b: 422; Wild, 2007b: 27; Arias-Penna, 2008b: 102; Lacau, *et al.* 2008: 253 (redescription); Branstetter & Sáenz, 2012: 255; Bezděčková, *et al.* 2015: 111; Feitosa, 2015c: 98; Feitosa & Prada-Achiardi, 2019: 678.

Senior synonym of *schmidti*: Brown, 1965c: 73; Kempf, 1972a: 256; Bolton, 1995b: 422.

richardsi. *Typhlomyrmex richardsi* Donisthorpe, 1939b: 161 (m.) GUYANA.

Junior synonym of *clavicornis*: Brown, 1965c: 71; Kempf, 1972a: 256; Bolton, 1995b: 422.

robustus. *Typhlomyrmex rogenhoferi* r. *robustus* Emery, 1890b: 40 (w.) COSTA RICA.

[Also described as new by Emery, 1894k: 46.]

Forel, 1899c: 3 (q.); Wheeler, G.C. & Wheeler, J. 1952a: 121 (l.).

Subspecies of *rogenhoferi*: Emery, in Dalla Torre, 1893: 16 (footnote); Forel, 1899c: 3.

Status as species: Dalla Torre, 1893: 16; Emery, 1911d: 34; Mann, 1922: 2; Wheeler, W.M. 1925a: 2.

Junior synonym of *rogenhoferi*: Brown, 1957e: 229; Brown, 1965c: 74; Kempf, 1972a: 256; Bolton, 1995b: 422.

rogenhoferi. *Typhlomyrmex rogenhoferi* Mayr, 1862: 737 (w.) BRAZIL (no state data).

[Misspelled as *rongenhoferi* by Menozzi, 1927c: 267.]

Mayr, 1887: 538 (q.m.); Wheeler, G.C. & Wheeler, J. 1976a: 42 (l.).

Status as species: Roger, 1863b: 21; Mayr, 1863: 457; Mayr, 1887: 538; Dalla Torre, 1893: 16; Forel, 1895b: 111; Emery, 1896g: 33; Forel, 1899c: 2; Emery, 1906c: 111; Emery, 1911d: 34; Forel, 1912b: 81; Bruch, 1914: 212; Mann, 1916: 403; Gallardo, 1918b: 14; Borgmeier, 1923: 54; Wheeler, W.M. 1925a: 2; Menozzi, 1927c: 267; Menozzi, 1931b: 261; Brown, 1957e: 229; Kempf, 1961b: 493; Brown, 1965c: 74; Kempf, 1970b: 325; Kempf, 1972a: 256; Brandão, 1991: 383; Bolton, 1995b: 422; Wild, 2007b: 27; Arias-Penna, 2008b: 102; Lacau, *et al.* 2008: 256 (redescription); Branstetter & Sáenz, 2012: 255; Bezděčková, *et al.* 2015: 111; Feitosa, 2015c: 98; Feitosa & Prada-Achiardi, 2019: 678.

Senior synonym of *manco*: Brown, 1957e: 229; Brown, 1965c: 74; Kempf, 1972a: 256; Bolton, 1995b: 422.

Senior synonym of *robustus*: Brown, 1957e: 229; Brown, 1965c: 74; Kempf, 1972a: 256; Bolton, 1995b: 422.

Senior synonym of *sculpturatus*: Brown, 1965c: 74; Kempf, 1972a: 256; Bolton, 1995b: 422.

schmidti. *Typhlomyrmex schmidti* Menozzi, 1927c: 268 (q.m.) COSTA RICA.

Junior synonym of *pusillus*: Brown, 1965c: 73; Kempf, 1972a: 256; Bolton, 1995b: 422.

sculpturatus. *Centromyrmex sculpturatus* Santschi, 1931c: 266 (q.) PANAMA.

Combination in *Typhlomyrmex*: Brown, 1965c: 74.

Junior synonym of *rogenhoferi*: Brown, 1965c: 74; Kempf, 1972a: 256; Bolton, 1995b: 422.