

Chemical Ecology in Insect-microbe Interactions in the Neotropics

Authors

Carla Menegatti* , Taise T. H. Fukuda* , Mônica T. Pupo 

Affiliation

Departamento de Ciências Farmacêuticas, Faculdade de Ciências Farmacêuticas de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, Brazil

Key words

antimicrobials, attine ants, interspecies interactions, microbial symbiosis, social insects, stingless bees

received

May 29, 2020

accepted after revision

July 29, 2020

published online

Bibliography

Planta Med 2020

DOI 10.1055/a-1229-9435

ISSN 0032-0943

© 2020. Thieme. All rights reserved.

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

Correspondence

Professor Dr. Mônica T. Pupo

Departamento de Ciências Farmacêuticas, Faculdade de Ciências Farmacêuticas de Ribeirão Preto, Universidade de São Paulo

Avenida do Café, 14040-903 Ribeirão Preto, SP, Brazil

Phone: + 55 16 33 15 47 10, Fax: + 55 16 33 15 41 78

mtpupo@fcrfp.usp.br

ABSTRACT

Small molecules frequently mediate symbiotic interactions between microorganisms and their hosts. Brazil harbors the highest diversity of insects in the world; however, just recently, efforts have been directed to deciphering the chemical signals involved in the symbioses of microorganisms and social insects. The current scenario of natural products research guided by chemical ecology is discussed in this review. Two groups of social insects have been prioritized in the studies, fungus-farming ants and stingless bees, leading to the identification of natural products involved in defensive and nutritional symbioses. Some of the compounds also present potential pharmaceutical applications as antimicrobials, and this is likely related to their ecological roles. Microbial symbioses in termites and wasps are suggested promising sources of biologically active small molecules. Aspects related to public policies for insect biodiversity preservation are also highlighted.

Introduction

Brazil is one of the world's 17 megadiverse countries that host between 15–20% of the entire world's biological diversity, including the greatest number of endemic species [1], offering innumerable research opportunities and sustainable technological development. Natural products research in Brazil has been historically focused on plant-derived compounds and evaluation of their biological activities [2], with only recent efforts directed to microbial-derived compounds and marine organisms [3–5].

Recent estimates based on DNA sequence information have shown that microorganisms consist in the highest number of living species, most of them yet to be described [6], but there are no records on the number of microbial species in Brazil. This untapped ecological niche represents a huge potential for natural products discovery, as suggested by sequencing of environmental bacterial DNA collected in different biomes [7].

Microorganisms are the most ancient form of life on earth and have established symbiotic interactions with several other organisms, from mammals to arthropods and plants [8]. Symbiosis involves any intimate species interaction, either positive or negative, including mutualism, commensalism, and parasitism, as defined by Bary [9]. Indeed, symbioses with microorganisms have contributed with innovations in eukaryotic evolution [10]. Much of the communication between microorganisms and between microorganisms and their environment is based on chemical interactions [11], the subject of research in the interdisciplinary field of chemical ecology [12]. Deciphering the chemical language between species is key to understanding how they interact and provides an opportunity for ecologically-guided natural products discovery for biotechnological purposes.

* These authors contributed equally to this work.

Insects dominate the known diversity of living organisms. Class Insecta comprises about 1 million described species and makes up 83.5% of all species in the Phylum Arthropoda [13]. Brazil harbors the highest diversity of insects in the world [14], but natural products research on insects is mainly focused on pheromones in the country [15]. As with other organisms, insects have developed a plethora of interspecies interactions with microorganisms, and social insects are fascinating examples of multipartite symbiosis.

Eusociality is an evolutionarily advanced level of social organization nearly confined to insects, especially ants, bees, wasps, and termites. Eusocial adult insects in a colony belong to different overlapping generations, care cooperatively for the offspring, and are divided into reproductive and nonreproductive castes [16]. Some social insects have also evolved symbiotic mutualistic interactions with fungi in which nutrient exchange between species is the key behind this association. Insects provide food resources—usually plant-derived material difficult to digest—to the fungi and protect them against opportunistic pathogens. In return, fungi can supply nutrients and molecules important for insect's physiological functions [17, 18]. Social insects are particularly attractive for microbial disease agents, since they live in high populational densities in relatively homeostatic nest environments and store food resources [19]. To avoid pathogens, social insects have evolved several strategies, such as grooming, nest hygiene, and chemical defenses [19, 20]. Another defensive strategy of insects is the symbiotic association with bacteria that produce and secrete biologically active small molecules that are selective against pathogens [20].

The biosynthetic potential of insect-associated microbes worldwide is covered elsewhere [17, 20–22], and here we highlight some of the work done on the Brazilian biodiversity. The biological activity of some microbial strains isolated from Brazilian social insects has been investigated but not exhaustively studied. These microbial isolates comprise yeasts, proteobacteria, and actinobacteria, which produce a variety of biologically active small molecules.

Natural Products Mediating Microbial Symbiosis Fungus-Farming Ants

E. O. Wilson, a recognized American biologist, naturalist, and writer, stated in 1975: “There are more species of ants in a square kilometer of Brazilian forest than all the species of primates in the world, more workers in a single colony of driver ants than all the lions and elephants in Africa” [23].


Fungus-farming “attine” ants (Formicidae: Myrmicinae: Attini: Attina) originated in a single ancestral attine in Amazon around 45 million y ago [24]. Approximately 250 species of fungus-farming ants are found in the “New World”, ranging from North America to South America [25]. Attine ants collect plant and other material they forage from the environment to nurture basidiomycete fungi they cultivate for food. During the evolution of these interactions, the foraged material has been diversified, giving rise to the so-called “basal” and “highly evolved” agricultural systems, based on the substrate that the fungal crop is fed [26]. Other microbes found in this multipartite symbiosis are a specialized path-

ogenic fungus from the genus *Escovopsis*, which can suppress the crop fungal cultivar and destroy the ant colony [27], and a symbiotic actinobacterium usually belonging to the genus *Pseudonocardia*, which produces small molecules that selectively inhibit the pathogenic fungus over the crop fungus [28, 29]. Dentigerumycin, produced by *Pseudonocardia* associated with *Apterostigma* ants collected in Central America, was the first selective antifungal characterized as mediator of the defensive symbiosis in attine ants (► Fig. 1) [30]. Even though there is a reasonable number of publications examining the chemistry involved in this symbiosis from attine colonies sampled in Central America [30–34], the potential of the microbial symbionts from Brazilian attine ants still remains to be explored.

Some attine ants have their exoskeletons covered by the actinobacteria, whereas others do not show the same obvious association. Ants of *Atta* genus, for instance, do not show specialized crypts—morphological structures that harbor actinobacteria—in their bodies. The bacterial symbionts are supposed to be internalized inside ants' bodies [35] or even in other places of the colony. The hypothesis that opportunistic pathogens were inhibited by microorganisms living in the colony was therefore tested. Different parts of *Atta sexdens* colonies (fungal garden, waste deposit, and surface of leaves collected by ants) were sampled and resulted in the isolation of 99 yeast strains. These strains had their inhibitory activities tested against 6 reporter yeast strains and also against each other. The results showed that 77 strains (78%) inhibited the growth of the competing strain. This high number of active strains pointed toward a role in maintaining the nest microbial community [36]. Investigations were also conducted into the role of bacterial strains associated with the fungal gardens of *Atta sexdens* ants collected in eucalyptus plantations in Rio de Janeiro state, Brazil. A high number of colonies was found to be associated with *Burkholderia* sp. strains, which inhibited different entomopathogenic fungi, such as *Metarhizium anisopliae*, *Beauveria bassiana*, the saprophytic fungus *Verticillium lecanii*, and the specific ant pathogen *Escovopsis weberi*. Although these strains had antifungal activity against different fungi, they did not show any activity against the fungal cultivar. *Burkholderia* sp. strains were isolated from 32 out of 57 ant nests (56%), also suggesting an important ecological role [37].

As well as producing small molecules with defensive roles, Proteobacteria can also fulfill other ecological functions. *Serratia marcescens* isolated from *Atta sexdens rubropilosa* colonies produce volatile pyrazines, including 2,5-dimethylpyrazine (1) and 3-ethyl-2,5-dimethylpyrazine (2) (► Fig. 2), that are components of ants' trail pheromone [38]. However, the dependence of ants on the microbial biosynthesis of trail pheromones remains to be elucidated. The microbial involvement in the production of pheromones has been recognized for some insects, but more efforts are needed to experimentally validate connections between the presence of specific symbionts, changes in the host's chemistry, and behavioral effects [39].

Leaf-cutter ants start new colonies with queen ants fecundated during the nuptial flight; however, mortality is extremely high during the nuptial flight and immediately afterward [24]. *A. sexdens rubropilosa* queen ants were found to be infected by the entomopathogenic fungus *Aspergillus nomius* after nuptial



► **Fig. 1** Ecological relationships between microorganisms and attine ants. Mutualistic interactions are represented by green arrows and red T-bars represent harmful interactions. Investigations into this system led to the isolation of the antifungal dentigerumycin [30]. Source: Monica Tallarico Pupo

flight. The fungus produced aflatoxin B1 (3) and aflatoxin G1 (4) (► **Fig. 2**) both *in situ* and *in vitro*. These compounds may play a pivotal role in the fungal pathogenicity observed for the *Atta* queens [40].

Different from *Atta* ants, leaf-cutter ants of genus *Acromyrmex* usually carry the symbiotic actinobacteria in crypts in their exoskeletons. *Acromyrmex subterraneus brunneus* worker ants at USP-campus, Piracicaba-SP, Brazil were sampled for the presence of actinobacteria, leading to the isolation of 20 actinobacteria strains. Among these bacteria, 17 strains belong to the genus *Streptomyces*, and the remaining are *Pseudonocardia*, *Kitasatospora*, and *Propionicimonas*. The majority of *Streptomyces* isolates inhibited the growth of the nest pathogen *Escovopsis weberi* [41]. These studies suggested that the ant-associated microbes isolated from samples collected in Brazil—other than *Pseudonocardia*—produce secondary metabolites with biological activity; however, chemical compounds responsible for antimicrobial activities are still elusive.

Actinobacteria can produce compounds showing great chemical diversity and a large variety of biological activities, and have evolved protective symbiotic interactions with different organisms [42]. Indeed, *Streptomyces* has been proven to be a good source of antimicrobial defenses in insects [22]. The first compounds from *Streptomyces* strains associated with fungus-farming ants in Brazil were the antimycins urauchimycin A (5) and urauchimycin B (6) (► **Fig. 2**) [43], with broad and potent antifungal activity against medically important *Candida* strains.


Recently, interdisciplinary research groups in Brazil have been mainly focused on the characterization of biologically active natural products from microbial symbionts of social insects [44]. Using an ecological-driven approach, the actinobacterial symbionts of

attine ants have been systematically screened against the specialized pathogenic fungus *Escovopsis* and then selected for further screening against other bacterial, fungal, and protozoan human pathogens.

The search for symbionts of *Acromyrmex subterraneus brunneus* ants, collected at USP-campus, Ribeirão Preto-SP, led to the isolation of *Streptomyces chartreusis* ICBG377, recovered from the fungal garden. The actinobacterium produces the antibiotic streptazolin (7), its *E*-isomer (8), strepchazolin A (9), strepchazolin B (10), and the inorganic compound cyclooctasulfur (11), the active compound against *Escovopsis* (► **Fig. 2**) [45]. Compound 11 was also produced by *S. chartreusis* ICBG323, isolated from the exoskeleton of winged males of *Mycocetopus goeldii* [45].

The actinobacterium *S. puniceus* ICBG378, isolated from *Acromyrmex rugosus rugosus* ants, produces griseorhodin A (12) and griseorhodin C (13), natural products known by their cytotoxic activity against cancer cell lines [46], and dinactin (14), active against *Escovopsis* (► **Fig. 2**) [47]. Dinactin (14) was also active against *Leishmania donovani*, one of the etiological agents of leishmaniasis, a neglected tropical disease that causes thousands of deaths yearly in developing countries.

Similarly, *Cyphomyrmex*-associated *Streptomyces* sp. ICBG292 produced Mer-A2026B (15), piericidin-A1 (16), and nigericin (17) (► **Fig. 2**), all active against *Escovopsis* and against intracellular amastigotes of *L. donovani*. Compounds 15 and 16 were also isolated from *Atta*-associated *Streptomyces*, while 14 and 17 showed the most potent leishmanicidal activities, with good selectivity indexes [47]. The biological activity of these compounds highlights the importance of exploring different sources for prospecting compounds that can help treating human diseases [47].



► **Fig. 2** Compounds isolated from microbial symbionts of attine ants collected in Brazil.

The hypothesis that the ant microbiome is a good source for exploring new medically useful antimicrobial agents was further reiterated. The isolation of cyphomycin (**18**) (► **Fig. 2**), a new antifungal polyketide, proved that the Brazilian biodiversity should be explored in order to find new candidates for the treatment of fungal infections. Cyphomycin (**18**) was produced by a *Cyphomyrmex*-associated *Streptomyces* strain and showed potent activity against human fungal pathogens both *in vitro* and *in vivo* [22].

Natural Products Mediating Microbial Symbiosis in Stingless Bees

Although a wide diversity of microorganisms, such as bacteria, fungi, and yeasts are found to be associated with bees, little is known about their role as beneficial symbionts [48]. Indeed, honey bees across the world harbor a rich bacterial community

[49–52]. Lactic acid bacteria (LAB) are frequently isolated from the guts of honey bees and bumble bees, and it is believed that this specific microbiota coevolved with their hosts, playing roles in nutrition and defense against pathogens [53–56].

More information is available regarding microbial pathogens of honey bees, which span several kingdoms, including the most damaging threats such as viruses, bacteria, and fungi [57]. *Paenibacillus larvae* and *Melissococcus plutonius*, the infective agents of American and European foulbrood diseases, respectively, are major bacterial threats to honey bees (*Apis* species) [57]. The American foulbrood disease (AFB) is widespread in honey bee larvae [58, 59] and drastically impacts the apiculture and the pollination of crops and wild plants [60]. The long-lived spores produced by the bacterium are infectious only for larvae, especially in early larval stages [61]. Burning the infected colonies is one usual treatment for AFB [62]. Antibiotics such as oxytetracycline are used in some countries for prevention and treatment of contami-

nated colonies still, but this approach is not effective against spores [63–67]. Other problems can be caused using antibiotics as their chemical residues accumulate in honey, reducing the longevity of the bees and selecting resistant *P. larvae* strains [61, 68]. The European foulbrood disease is caused by the globally distributed Gram-positive, non-spore-forming bacterium *M. plutonius* [69]. Ingestion of larval food contaminated with *M. plutonius* causes infection in larvae. Bumble bees (*Bombus terrestris*), important pollinators of the northern hemisphere, are colonized by the virulent parasite trypanosomatid *Crithidia bombi* (Trypanosomatidae, Zoomastigophorea) [70]. These infections have a variety of consequences, such as the mortality of the colonies.

While microbial diseases are better understood for honey bees, little is known about microbial diseases affecting stingless bees. Stingless bees (Apidae: Meliponini) are a large group of bees with more than 500 species described, around 300 of them occur in Brazil [71]. Although *A. mellifera* is originally from Africa, this species is an important pollinator widespread around the world [72]. The global distribution of honey bees favors their microbial pathogens to spillover stingless bees native to tropical and subtropical regions [73]. Indeed, some honey bee pathogens already detected in stingless bees include the disease-causing bacterium *Lysinibacillus sphaericus* in Australia [74], the acute bee paralysis virus (ABPV) in Brazil [75], the bacterium *M. plutonius* in Brazil [76], and the fungus *Nosema ceranae* in laboratory colonies [73]. Therefore, an ecological approach to study bacterial symbionts of stingless bees involved in defensive responses can be based on microbial pathogens of honey bees and bumble bees.

Research has been directed toward the role of the associated microbiota in protecting bees against pathogens. LAB have a potential role in controlling the bacterial pathogens causing American and European foulbrood diseases [77]. Stingless bees from different geographical regions also carry LAB [78] that might possess similar functions. Bacteriocin-like compounds, active against *P. larvae* were also identified from LAB isolated from honey bees in Argentina [79]. Actinobacteria have also been isolated from stingless bees from other tropical and subtropical regions. Antibiotic-producing *Streptomyces* spp. were isolated from the stingless bee *Tetragonisca angustula* in Costa Rica, showing antimicrobial activities against a variety of human pathogens [80]. Actinobacteria strains active against *P. larvae* and *M. plutonius* were isolated from colonies of honey bees (*A. mellifera*, *A. cerana*, *A. floreae*) and stingless bees (*Trigona laeviceps* and *T. fuscobalteata*) in Thailand [81]. However, the small molecules mediating these defensive symbioses have not been comprehensively studied.

The first unprecedented example of nutritional symbiosis in stingless bees is the Brazilian bee *Scaptotrigona depilis*, which is surrounded by a complex microbial community. *S. depilis* cultivates a fungus of the genus *Zygosaccharomyces* in the brood cell, which provides ergosterol (19) (► Fig. 3) as a precursor for ecdysteroid biosynthesis and, consequently, for proper larval development and metamorphosis [82, 83]. Two additional fungi are also active in the cerumen of brood cells, *Candida* sp. and *Monascus ruber*, which modulate *Zygosaccharomyces* sp. growth. *Candida* sp. produces volatile alcohols such as ethanol and isoamyl alcohol that stimulate the growth of *Zygosaccharomyces* sp., while *Monascus ruber* inhibits *Zygosaccharomyces* sp. and *Candida* sp. by the


production of lovastatin (20) and monascin (21), respectively (► Fig. 3) [84]. Fungi of the genus *Monascus* were also found in association with other species of stingless bees in Brazil, but their chemical-ecological functions remain unknown [85]. The larvae of *S. depilis* also engage in associations with microbes. Genome analyses of the *Bacillus* sp. SDLI1 isolated from *S. depilis* larvae indicated the presence of biosynthetic gene clusters that encode the production of a variety of antibiotics, suggesting a bacterial defensive symbiosis [86]. *Bacillus* spp. have been commonly associated with honey bees' larvae and inhibit *P. larvae* [87, 88]. The isolation of *Bacillus* DNA in fossils showed close phylogenetic relationships with strains typically isolated from stingless bees, which could provide information about the evolution of microbe-insect symbiosis [89].

The stingless bee *Melipona scutellaris* inhabits in Northeast Brazil and engages in a relationship with various bacteria. The ecological-driven approach of bioassays against entomopathogens led to the identification of some bacterial strains as possible defensive symbionts and their chemical signals. The bacterium *Paenibacillus polymyxa* was isolated from the larval food of *M. scutellaris* and produces (L)-(-)-phenyllactic acid (22) and a family of 9 lipodepsipeptides known as fusaricidins, including the major compounds fusaricidin A (23) and fusaricidin B (24) (► Fig. 3), active against the entomopathogenic fungus *B. bassiana* and bacterium *P. larvae*. Interestingly, fusaricidins 23 and 24 were also detected directly in the larval food of different sampled colonies, suggesting a beneficial defensive role against pathogens [90]. Adult *M. scutellaris* bees also carry several actinobacteria in their bodies. *Streptomyces* sp. ICBG1323 and *Micromonospora* sp. ICBG1321 were isolated from nurse and forager bees, respectively. Two families of structurally complex bioactive polyketides were isolated from the associated strains: lobophorins (25–28) from *Streptomyces* sp. ICBG1323 and anthracyclines from *Micromonospora* sp. ICBG132 (29–39), including the rare quinocyclines 29–34 and the novel compound 39 (► Fig. 3). The compounds presented variable levels of activities against *P. larvae*. Compounds 28 and 30 showed the higher antibacterial activity, better than the control antibiotics [91]. Finally, two new compounds were isolated from *Streptomyces* sp. ICBG1318 strain in association with *M. scutellaris* nurse bees. The novel cyclodepsipeptides named meliponamycin A (40) and meliponamycin B (41) (► Fig. 3) were strongly active against *P. larvae* and human pathogens, such as *Staphylococcus aureus* and *L. infantum* [92].

The examples highlight that more research on the stingless bees-associated microbiota is essential to enhance the current knowledge of the molecular signals involved in these symbiotic interactions. This knowledge might contribute to design policies for the preservation of these important pollinators of native flora and agricultural crops.

Microbial Symbiosis in Wasps

Previous research efforts on digger wasps of the genus *Philanthus* (beewolves; Hymenoptera, Crabronidae), which consists of more than 100 species widespread in Europe, Africa, Asia, and North America, showed association with the symbiotic actinobacteria *Candidatus Streptomyces philanthi* [93, 94]. The symbionts are cul-




► Fig. 3 Compounds isolated from microbial symbionts of Brazilian stingless bees.

tivated in specialized antennal gland reservoirs and transferred to the brood cells where they produce antibiotics such as streptochlorin and piericidin derivatives responsible for protecting the wasps' larvae against pathogens [95,96]. The Brazilian digger wasps *Trachypus boharti* also present bacteria in the antennal gland reservoirs. Gene sequences revealed that among all antennal symbionts described, the Brazilian wasps cultivate the most distantly related actinobacteria [97]. However, the chemistry behind this protective symbiosis remains to be uncovered.

Another example of wasp-microbe symbiosis is established between the parasitic wasp *Asobara tabida* with the bacterium *Wolbachia*. The bacterium is vertically transmitted via the eggs by wasps and plays a fundamental role in oogenesis completion [98]. The treatment with antibiotics to eliminate *Wolbachia* found

that aposymbiotic females of *A. tabida* are reproductively sterile, being unable to produce viable offspring [99]. An intense apoptosis process is responsible for the absence of egg production, and there is evidence that *Wolbachia* inhibits the programmed cell death by the disruption of cellular physiology of the host [100]. The symbiosis with the bacterium *Wolbachia* was found for wasps of the genus *Encarsia*; meanwhile, another bacterium described as "*Encarsia bacterium*" was found to be associated with a population of *Encarsia* wasps, including *E. pergandiella* collected in Brazil. The bacterium is related to parthenogenesis [101]. The symbiont "*Candidatus Cardinium hertigii*" associated with *Encarsia* wasps from Brazil and USA is also linked to reproductive alterations in the host [102].



► **Fig. 4** Compounds isolated from microbial symbionts of termites.

Parasitoid wasps lay their eggs into other arthropods who are hosts for wasp larval development [103]. Braconid wasps engaged in an ancient relationship with polydnavirus that suppress host defense mechanisms and permit the larval development [104–106]. This symbiosis is so old (about 70 million y) that the genes involved in viral replication have been incorporated into the wasp genome [107].

Invasive wood wasps *Sirex noctilio* collected in USA are associated with *Streptomyces* strains with specific enzymatic activities responsible for degrading cellulose, which is used by the insect as source of energy [108]. Wood wasps still hold a close relationship with a fungal symbiont *Amylostereum chailletii* that feeds wasp larvae, providing them with digestive enzymes [109].

Very few efforts have been pursued on revealing the chemical signaling in interactions between microbes and wasps in Brazil. The country harbors the richest fauna of social wasps (Polistinae) in the world, with more than 300 species, 104 of them endemic from Brazil [110], offering several opportunities for chemical ecology based natural products discovery.

Microbial Symbiosis in Termites

Although fungus-farming ants and fungus-growing termites share behavioral similarities, and both seem to rely on the presence of symbiotic actinobacteria to chemically defend their nests against fungal pathogens [54, 111], they do not share a common ancestor with the same characteristics [112]. Moreover, these insects differ from each other in the geographical distribution. While attine ants originated and are found in the “New World”, fungus growing Macrotermitinae termites (Termitidae: Macrotermitinae) originated in Africa and comprise about 330 species distributed in the “Old World”, including Africa and Asia [113]. Macrotermitinae termites are subdivided into 11 genera [114].

Brazil houses around 300 species of termites belonging to the families Kalotermitidae, Rhinotermitidae, Serritermitidae, and Termitidae [115]. Termites contribute to structure and composition of soils by efficiently degrading biomass with the aid of resi-

dent gut microbiota, so most of the research in Brazil has focused on the enzymatic potential of termite-associated microbiota [116]. Termite microbiota might also have a contribution in defensive symbiosis by the production of secondary metabolites. Indeed, two *Streptomyces* strains from termite mounds collected in Bahia State showed percentages of inhibition above 98% against bovine viral diarrhea virus (BVDV), but the active compound has not been identified [117].

There are some examples of natural products produced by actinobacteria in association with African fungus-growing termites [118–122]. *Amycolatopsis* sp. produced macrotermycins A–D (42–45) (► **Fig. 4**). Besides the antifungal ecological role, these 4 macrolactams presented antibacterial activity against *S. aureus* [123]. The polyketide fridamycin A (46) (► **Fig. 4**) was isolated from the termite-associated *Actinomadura* sp.; it demonstrated glucose uptake stimulation and could be an option for type 2 diabetes therapeutics [124]. Microtermolides A (47) and B (48) (► **Fig. 4**) were isolated from a fungus-growing termite-associated *Streptomyces* sp. [125, 126]. Both compounds are depsipeptides; moreover, microtermolide B is a rare linear depsipeptide and seems to be the first one of this class produced by a *Streptomyces* strain. The lack of extensive work on termite-associated actinobacteria in Brazil, however, can be due to the absence of fungus-growing termites in this region of the world [127]. Brazilian termites, as wasps, are eusocial insects. Even though there is not much work on Brazilian termites, this system seems promising since their colonies are susceptible to parasitic pressure, and previous works have demonstrated the beneficial association between social insects and antibiotic-producing microbes.

Conclusion

Brazil harbors an impressive reservoir of genetic resources in different biomes, including the highest number of known insects in the world and an undescribed microbial diversity. The chemistry of microbial natural products and chemical ecology of microbial symbiosis are complementary research fields in their early days in

the country. Two decades have passed since the publication of the first natural product from a microbial source [128], and 7 y since the first report on natural products mediating microbial interactions [129] in Brazil.

The chemistry involved in interspecies interactions between insects and microbes remains largely to be unveiled. The examples showed here are just a glimpse of the chemodiversity involved in nutritional and defensive microbial symbiosis in some species of attine ants and stingless bees in Brazil. Several other species of these insects should be investigated, as well as other Brazilian social insects such as termites and wasps.

The understanding of how species interact in nature is instrumental to design sustainable approaches for their uses. Besides improving the knowledge about interspecies interactions, the chemical ecology approach to studying insect-microbe symbiosis might lead to the identification of biologically active compounds with privileged scaffolds for biotechnological development, mainly as agrochemicals and pharmaceuticals. The huge biodiversity remaining in Brazil potentially encodes useful products to be developed based on sustainable practices. Efforts of multidisciplinary research groups in chemistry, microbiology, molecular biology, entomology, and pharmacology are instrumental to achieve such results.

Not less important is the regular and prioritized governmental financial support for research in the field. The biodiversity of insects has declined worldwide; therefore, researchers and government in Brazil might act synergistically. According to Sánchez-Bayoa & Wyckhuys [130], the main drivers of insect decline are: i) habitat loss and conversion to intensive agriculture and urbanization; ii) pollution, mainly that by synthetic pesticides and fertilizers; iii) biological factors, including pathogens and introduced species; and iv) climate change. It is out of scope to discuss each one of those factors in this article but needless to explain that all of them occur in Brazil.

Strong public policies are urgently needed to protect Brazilian biodiversity. Specifically, it is important to consider the preservation of native insects and, consequently, the benefits they provide—in association with resident microbes—in the structure and functioning of the ecosystems.

Contributors' Statement

Conception and design of the work: M. T. Pupo; Data collection: C. Menegatti, T. T. H. Fukuda, M. T. Pupo; Analysis and interpretation of the data: C. Menegatti, T. T. H. Fukuda, M. T. Pupo; Drafting the manuscript: C. Menegatti, T. T. H. Fukuda, M. T. Pupo; Critical revision of the manuscript: M. T. Pupo.

Acknowledgements

The research on microbial symbionts of insects is financially supported by the São Paulo Research Foundation (FAPESP) grants #2013/50954-0 (MTP), #2013/07600-3 (MTP; CEPID-CIBFar), #2016/15576-3 (CM), and #2015/26349-5 (TTHF) and also by the Conselho Nacional de Pesquisa e Desenvolvimento Tecnológico (CNPq) grant #303792/2018-2 (MTP). This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior-Brasil (CAPES), Finance Code 001.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

- [1] Mittermeier RA, Da Fonseca GAB, Rylands AB, Brandon K. A brief history of biodiversity conservation in Brazil. *Conserv Biol* 2005; 19: 601–607
- [2] Pinto AC, Silva DHS, Bolzani VD, Lopes NP, Epifanio RD. Current status, challenges and trends on natural products in Brazil. *Quim Nova* 2002; 25: 45–61
- [3] Berlinck RGS, Borges WD, Scotti MT, Vieira PC. The chemistry of natural products in Brazil in the XXI century. *Quim Nova* 2017; 40: 706–710
- [4] de Oliveira LG, Pupo MT, Vieira PC. Exploring microbial natural products in the frontiers of chemistry and biology. *Quim Nova* 2013; 36: 1577–1586
- [5] Ioca LP, Allard PM, Berlinck RGS. Thinking big about small beings—the (yet) underdeveloped microbial natural products chemistry in Brazil. *Nat Prod Rep* 2014; 31: 646–675
- [6] Castelle CJ, Banfield JF. Major new microbial groups expand diversity and alter our understanding of the tree of life. *Cell* 2018; 172: 1181–1197
- [7] Charlop-Powers Z, Owen JG, Reddy BVB, Ternei M, Guimaraes DO, de Frias UA, Pupo MT, Seepe P, Feng ZY, Brady SF. Global biogeographic sampling of bacterial secondary metabolism. *Elife* 2015; 4: e05048
- [8] Moran NA. Symbiosis. *Curr Biol* 2006; 16: R866–R871
- [9] Wilkinson DM. At cross purposes—how do we cope with scientific terms that have two different definitions? *Nature* 2001; 412: 485
- [10] Russell JA, Sanders JG, Moreau CS. Hotspots for symbiosis: function, evolution, and specificity of ant-microbe associations from trunk to tips of the ant phylogeny (Hymenoptera: Formicidae). *Myrmecol News* 2017; 24: 43–69
- [11] Mithofer A, Boland W. Do you speak chemistry? Small chemical compounds represent the evolutionary oldest form of communication between organisms. *EMBO Rep* 2016; 17: 626–629
- [12] Bergstrom G. Chemical ecology = chemistry plus ecology! *Pure Appl Chem* 2007; 79: 2305–2323
- [13] Stork NE. How many species of insects and other terrestrial arthropods are there on earth? *Annu Rev Entomol* 2018; 63: 31–45
- [14] Rafael JA, Aguiar AP, Amorim DD. Knowledge of insect diversity in Brazil: challenges and advances. *Neotrop Entomol* 2009; 38: 565–570
- [15] Bergmann J, Gonzalez A, Zarbin PHG. Insect pheromone research in South America. *J Brazil Chem Soc* 2009; 20: 1206–1219
- [16] Wilson EO, Holldobler B. Eusociality: origin and consequences. *P Natl Acad Sci USA* 2005; 102: 13367–13371
- [17] Ramadhar TR, Beemelmanns C, Currie CR, Clardy J. Bacterial symbionts in agricultural systems provide a strategic source for antibiotic discovery. *J Antibiot* 2014; 67: 53–58
- [18] Biedermann PHW, Vega FE. Ecology and evolution of insect-fungus mutualisms. *Annu Rev Entomol* 2020; 65: 431–455
- [19] Evans JD, Aronstein K, Chen YP, Hetru C, Imler JL, Jiang H, Kanost M, Thompson GJ, Zou Z, Hultmark D. Immune pathways and defence mechanisms in honey bees *Apis mellifera*. *Insect Mol Biol* 2006; 15: 645–656
- [20] Van Arnem EB, Currie CR, Clardy J. Defense contracts: molecular protection in insect-microbe symbioses. *Chem Soc Rev* 2018; 47: 1638–1651
- [21] Fukuda TTH, Cassilly CD, Gerdt JP, Henke MT, Helfrich EJM, Mevers E. Research tales from the Clardy laboratory: function-driven natural product discovery. *J Nat Prod* 2020; 83: 744–755
- [22] Chevrette MG, Carlson CM, Ortega HE, Thomas C, Ananiev GE, Barns KJ, Book AJ, Cagnazzo J, Carlos C, Flanigan W, Grubbs KJ, Horn HA, Hoffmann FM, Klassen JL, Knack JJ, Lewin GR, McDonald BR, Muller L, Melo WGP, Pinto-Tomas AA, Schmitz A, Wendt-Pienkowski E, Wildman

- S, Zhao M, Zhang F, Bugni TS, Andes DR, Pupo MT, Currie CR. The antimicrobial potential of *Streptomyces* from insect microbiomes. *Nat Commun* 2019; 10: 516
- [23] Wilson EO. *Sociobiology: The new Synthesis*. Cambridge: Harvard University Press; 1975
- [24] Hölldobler B, Wilson EO. *The Ants*. Cambridge: Harvard University Press; 1990: 596
- [25] Branstetter MG, Jesovnik A, Sosa-Calvo J, Lloyd MW, Faircloth BC, Brady SG, Schultz TR. Dry habitats were crucibles of domestication in the evolution of agriculture in ants. *P Roy Soc B-Biol Sci* 2017; 284: 20170095
- [26] Schultz TR, Brady SG. Major evolutionary transitions in ant agriculture. *P Natl Acad Sci USA* 2008; 105: 5435–5440
- [27] Currie CR, Mueller UG, Malloch D. The agricultural pathology of ant fungus gardens. *P Natl Acad Sci USA* 1999; 96: 7998–8002
- [28] Currie CR, Scott JA, Summerbell RC, Malloch D. Fungus-growing ants use antibiotic-producing bacteria to control garden parasites. *Nature* 1999; 398: 701–704
- [29] Cafaro MJ, Currie CR. Phylogenetic analysis of mutualistic filamentous bacteria associated with fungus-growing ants. *Can J Microbiol* 2005; 51: 441–446
- [30] Oh DC, Poulsen M, Currie CR, Clardy J. Dentigerumycin: a bacterial mediator of an ant-fungus symbiosis. *Nat Chem Biol* 2009; 5: 391–393
- [31] Carr G, Derbyshire ER, Caldera E, Currie CR, Clardy J. Antibiotic and antimalarial quinones from fungus-growing ant-associated *Pseudonocardia* sp. *J Nat Prod* 2012; 75: 1806–1809
- [32] Sit CS, Ruzzini AC, Van Arnem EB, Ramadhar TR, Currie CR, Clardy J. Variable genetic architectures produce virtually identical molecules in bacterial symbionts of fungus-growing ants. *P Natl Acad Sci USA* 2015; 112: 13150–13154
- [33] Van Arnem EB, Ruzzini AC, Sit CS, Currie CR, Clardy J. A Rebeccamycin analog provides plasmid-encoded niche defense. *J Am Chem Soc* 2015; 137: 14272–14274
- [34] Van Arnem EB, Ruzzini AC, Sit CS, Horn H, Pinto-Tomas AA, Currie CR, Clardy J. Selvamycin, an atypical antifungal polyene from two alternative genomic contexts. *P Natl Acad Sci USA* 2016; 113: 12940–12945
- [35] Marsh SE, Poulsen M, Gorosito NB, Pinto-Tomas A, Masiulionis VE, Currie CR. Association between *Pseudonocardia* symbionts and *Atta* leaf-cutting ants suggested by improved isolation methods. *Int Microbiol* 2013; 16: 17–25
- [36] Carreiro SC, Pagnocca FC, Bacci M, Bueno OC, Hebling MJA, Middelhoven WJ. Occurrence of killer yeasts in leaf-cutting ant nests. *Folia Microbiol* 2002; 47: 259–262
- [37] Santos AV, Dillon RJ, Dillon VM, Reynolds SE, Samuels RI. Occurrence of the antibiotic producing bacterium *Burkholderia* sp in colonies of the leaf-cutting ant *Atta sexdens rubropilosa*. *FEMS Microbiol Lett* 2004; 239: 319–323
- [38] Silva-Junior EA, Ruzzini AC, Paludo CR, Nascimento FS, Currie CR, Clardy J, Pupo MT. Pyrazines from bacteria and ants: convergent chemistry within an ecological niche. *Sci Rep* 2018; 8: 1–7
- [39] Engl T, Kaltenpoth M. Influence of microbial symbionts on insect pheromones. *Nat Prod Rep* 2018; 35: 386–397
- [40] Silva-Junior EA, Paludo CR, Valadares L, Lopes NP, do Nascimento FS, Pupo MT. Aflatoxins produced by *Aspergillus nomius* ASR3, a pathogen isolated from the leaf-cutter ant *Atta sexdens rubropilosa*. *Rev Bras Farmacogn* 2017; 27: 529–532
- [41] Zucchi TD, Guidolin AS, Consoli FL. Isolation and characterization of actinobacteria ectosymbionts from *Acromyrmex subterraneus brunneus* (Hymenoptera, Formicidae). *Microbiol Res* 2011; 166: 68–76
- [42] Van der Meij A, Worsley SF, Hutchings MI, van Wezel GP. Chemical ecology of antibiotic production by actinomycetes. *FEMS Microbiol Rev* 2017; 41: 392–416
- [43] Mendes TD, Borges WS, Rodrigues A, Solomon SE, Vieira PC, Duarte MCT, Pagnocca FC. Anti-*Candida* properties of urauchimycins from actinobacteria associated with *Trachymyrmex* ants. *Biomed Res Int* 2013; 2013: 1–8
- [44] Pupo MT, Currie CR, Clardy J. Microbial symbionts of insects are the focus of the first international cooperative biodiversity group (ICBG) in Brazil. *J Brazil Chem Soc* 2017; 28: 393–401
- [45] Ortega HE, Batista JM, Melo WGP, de Paula GT, Pupo MT. Structure and absolute configuration of secondary metabolites from two strains of *Streptomyces chartreusis* associated with attine ants. *J Brazil Chem Soc* 2019; 30: 2672–2680
- [46] Ortega HE, Batista JM, Melo WGP, Clardy J, Pupo MT. Absolute configurations of griseorhodins A and C. *Tetrahedron Lett* 2017; 58: 4721–4723
- [47] Ortega HE, Ferreira LLG, Melo WGP, Oliveira ALL, Alvarenga RFR, Lopes NP, Bugni TS, Andricopulo AD, Pupo MT. Antifungal compounds from *Streptomyces* associated with attine ants also inhibit *Leishmania donovani*. *Plos Neglect Trop D* 2019; 13: e0007643
- [48] Moraes PB, Calaça PSST, Rosa CA. Microorganisms associated with stingless Bees. In: Vit P, Pedro SR, Roubik D, eds. *Pot-Honey*: Springer; 2013: 173–186
- [49] Kwong WK, Moran NA. Gut microbial communities of social bees. *Nat Rev Microbiol* 2016; 14: 374
- [50] Moran NA. Genomics of the honey bee microbiome. *Curr Opin Insect Sci* 2015; 10: 22–28
- [51] Mattila HR, Rios D, Walker-Sperling VE, Roeselers G, Newton ILG. Characterization of the active microbiotas associated with honey bees reveals healthier and broader communities when colonies are genetically diverse. *PLoS One* 2012; 7: e32962
- [52] Martinson VG, Danforth BN, Minckley RL, Rueppell O, Tingek S, Moran NA. A simple and distinctive microbiota associated with honey bees and bumble bees. *Mol Ecol* 2011; 20: 619–628
- [53] Crotti E, Rizzi A, Chouaia B, Ricci I, Favia G, Alma A, Sacchi L, Bourtzis K, Mandrioli M, Cherif A, Bandi C, Daffonchio D. Acetic acid bacteria, newly emerging symbionts of insects. *Appl Environ Microb* 2010; 76: 6963–6970
- [54] Florez LV, Biedermann PHW, Engl T, Kaltenpoth M. Defensive symbioses of animals with prokaryotic and eukaryotic microorganisms. *Nat Prod Rep* 2015; 32: 904–936
- [55] Forsgren E, Olofsson TC, Vasquez A, Fries I. Novel lactic acid bacteria inhibiting *Paenibacillus larvae* in honey bee larvae. *Apidologie (Celle)* 2010; 41: 99–108
- [56] Vasquez A, Forsgren E, Fries I, Paxton RJ, Flaberg E, Szekely L, Olofsson TC. Symbionts as major modulators of insect health: lactic acid bacteria and honeybees. *PLoS One* 2012; 7: e33188
- [57] Evans JD, Schwarz RS. Bees brought to their knees: microbes affecting honey bee health. *Trends Microbiol* 2011; 19: 614–620
- [58] White GF. *Bacteria of the apiary*. Technical series (United States Bureau of Entomology) 1906; 14. doi:10.5962/bhl.title.87503
- [59] Ashiralieva A, Genersch E. Reclassification, genotypes and virulence of *Paenibacillus larvae*, the etiological agent of American foulbrood in honeybees—a review. *Apidologie (Celle)* 2006; 37: 411–420
- [60] Crailsheim K, Riessberger-Gallé U. Honey bee age-dependent resistance against American foulbrood. *Apidologie (Celle)* 2001; 32: 91–103
- [61] Genersch E. American foulbrood in honeybees and its causative agent, *Paenibacillus larvae*. *J Invertebr Pathol* 2010; 103: S10–S19
- [62] Waite R, Brown M, Thompson H, Bew M. Control of American foulbrood by eradication of infected colonies. *Apiacta* 2003; 38: 134–136
- [63] Reybroeck W, Daeseleire E, De Brabander HF, Herman L. Antimicrobials in beekeeping. *Vet Microbiol* 2012; 158: 1–11
- [64] Alippi AM, Albo GN, Leniz D, Rivera I, Zanelli ML, Roca AE. Comparative study of tylosin, erythromycin and oxytetracycline to control American foulbrood of honey bees. *J Apicult Res* 1999; 38: 149–158

- [65] Kochansky J, Knox DA, Feldlaufer M, Pettis JS. Screening alternative antibiotics against oxytetracycline-susceptible and-resistant *Paenibacillus larvae*. *Apidologie (Celle)* 2001; 32: 215–222
- [66] Katznelson H. The influence of antibiotics and sulfa drugs on *Bacillus larvae*, cause of American foulbrood of the honeybee, *in vitro* and *in vivo*. *J Bacteriol* 1950; 59: 471
- [67] Kochansky J, Pettis J. Screening additional antibiotics for efficacy against American foulbrood. *J Apicult Res* 2005; 44: 24–28
- [68] Bogdanov S. Contaminants of bee products. *Apidologie (Celle)* 2006; 37: 1–18
- [69] Funfhaus A, Ebeling J, Genersch E. Bacterial pathogens of bees. *Curr Opin Insect Sci* 2018; 26: 89–96
- [70] Koch H, Schmid-Hempel P. Socially transmitted gut microbiota protect bumble bees against an intestinal parasite. *P Natl Acad Sci USA* 2011; 108: 19288–19292
- [71] Michener CD. The Meliponini. In: Vit P, Pedro, editor. *Pot-honey: a legacy of stingless bees*. Springer; 2013: 3–17
- [72] Whitfield CW, Behura SK, Berlocher SH, Clark AG, Johnston JS, Sheppard WS, Smith DR, Suarez AV, Weaver D, Tsutsui ND. Thrice out of Africa: ancient and recent expansions of the honey bee, *Apis mellifera*. *Science* 2006; 314: 642–645
- [73] Purkiss T, Lach L. Pathogen spillover from *Apis mellifera* to a stingless bee. *P Roy Soc B-Biol Sci* 2019; 286: 20191071
- [74] Shanks JL, Haigh AM, Riegler M, Spooner-Hart RN. First confirmed report of a bacterial brood disease in stingless bees. *J Invertebr Pathol* 2017; 144: 7–10
- [75] Ueira-Vieira C, Almeida LO, de Almeida FC, Amaral IMR, Brandeburgo MAM, Bonetti AM. Scientific note on the first molecular detection of the acute bee paralysis virus in Brazilian stingless bees. *Apidologie (Celle)* 2015; 46: 628–630
- [76] Teixeira EW, Ferreira EA, da Luz CFP, Martins MF, Ramos TA, Lourenco AP. European foulbrood in stingless bees (Apidae: Meliponini) in Brazil: old disease, renewed threat. *J Invertebr Pathol* 2020; 172: 107357
- [77] Olofsson TC, Butler E, Markowicz P, Lindholm C, Larsson L, Vasquez A. Lactic acid bacterial symbionts in honeybees—an unknown key to honey's antimicrobial and therapeutic activities. *Int Wound J* 2016; 13: 668–679
- [78] Leonhardt SD, Kaltenpoth M. Microbial communities of three sympatric Australian stingless bee species. *PLoS One* 2014; 9: e105718
- [79] Audisio MC, Torres MJ, Sabate DC, Ibaguren C, Apella MC. Properties of different lactic acid bacteria isolated from *Apis mellifera* L. bee-gut. *Microbiol Res* 2011; 166: 1–13
- [80] Cambroner-Henrichs JC, Matarrita-Carranza B, Murillo-Cruz C, Araya-Valverde E, Chavarria M, Pinto-Tomas AA. Phylogenetic analyses of antibiotic-producing *Streptomyces* sp. isolates obtained from the stingless-bee *Tetragonisca angustula* (Apidae: Meliponini). *Microbiol-Sgm* 2019; 165: 292–301
- [81] Promnuan Y, Kudo T, Chantawannakul P. Actinomycetes isolated from beehives in Thailand. *World J Microb Biot* 2009; 25: 1685–1689
- [82] Menezes C, Vollet-Neto A, Marsaioli AJ, Zampieri D, Fontoura IC, Luchessi AD, Imperatriz-Fonseca VL. A Brazilian social bee must cultivate fungus to survive. *Curr Biol* 2015; 25: 2851–2855
- [83] Paludo CR, Menezes C, Silva-Junior EA, Vollet-Neto A, Andrade-Dominguez A, Pishchany G, Khadempour L, Nascimento FS, Currie CR, Kolter R, Clardy J, Pupo MT. Stingless bee larvae require fungal sterol to pupate. *Sci Rep* 2018; 8: 1122
- [84] Paludo CR, Pishchany G, Andrade-Dominguez A, Silva-Junior EA, Menezes C, Nascimento FS, Currie CR, Kolter R, Clardy J, Pupo MT. Microbial community modulates growth of symbiotic fungus required for stingless bee metamorphosis. *PLoS One* 2019; 14: e0219696
- [85] Barbosa R, Leong S, Vinnere-Pettersson O, Chen A, Souza-Motta C, Frisvad J, Samson R, Oliveira N, Houbraken J. Phylogenetic analysis of *Monascus* and new species from honey, pollen and nests of stingless bees. *Stud Mycol* 2017; 86: 29–51
- [86] Paludo CR, Ruzzini AC, Silva-Junior EA, Pishchany G, Currie CR, Nascimento FS, Kolter RG, Clardy J, Pupo MT. Whole-genome sequence of *Bacillus* sp. SDLI1, isolated from the social bee *Scaptotrigona depilis*. *Genome Announc* 2016; 4: e00174-00116
- [87] Evans JD, Armstrong TN. Inhibition of the American foulbrood bacterium, *Paenibacillus larvae larvae*, by bacteria isolated from honey bees. *J Apicult Res* 2005; 44: 168–171
- [88] Gilliam M. Identification and roles of non-pathogenic microflora associated with honey bees. *FEMS Microbiol Lett* 1997; 155: 1–10
- [89] Cano RJ, Borucki MK, Higbyschweitzer M, Poinar HN, Poinar GO, Pollard KJ. *Bacillus* dna in fossil bees—an ancient symbiosis. *Appl Environ Microb* 1994; 60: 2164–2167
- [90] Menegatti C, Melo WGP, Carrão DB, Oliveira ARM, Nascimento FS, Lopes NP, Pupo MT. *Paenibacillus polymyxa* associated with the stingless bee *Melipona scutellaris* produces antimicrobial compounds against entomopathogens. *J Chem Ecol* 2018; 44: 1158–1169
- [91] Rodriguez-Hernandez D, Melo WGP, Menegatti C, Lourenzon VB, Nascimento FS, Pupo MT. Actinobacteria associated with stingless bee biosynthesize bioactive polyketides against bacterial pathogen. *New J Chem* 2019; 43: 10109–10117
- [92] Menegatti C, Lourenzon VB, Rodríguez-Hernández D, Melo WGP, Ferreira LLG, Andricopulo AD, Nascimento FS, Pupo MT. Meliponamycins: antimicrobials from stingless bee-associated *Streptomyces* sp. *J Nat Prod* 2020; 83: 610–616
- [93] Kaltenpoth M, Goettler W, Dale C, Stubblefield JW, Herzner G, Roeser-Mueller K, Strohm E. “*Candidatus Streptomyces philanthi*”, an endosymbiotic streptomycete in the antennae of *Phantherus digger* wasps. *Int J Syst Evol Micr* 2006; 56: 1403–1411
- [94] Bohart RM, Bohart RM, Menke AS. *Sphecids Wasps of the World: A Generic Revision*. Berkeley: University of California Press; 1976
- [95] Kaltenpoth M, Goettler W, Koehler S, Strohm E. Life cycle and population dynamics of a protective insect symbiont reveal severe bottlenecks during vertical transmission. *Evol Ecol* 2010; 24: 463–477
- [96] Kroiss J, Kaltenpoth M, Schneider B, Schwinger MG, Hertweck C, Maddula RK, Strohm E, Svatos A. Symbiotic streptomycetes provide antibiotic combination prophylaxis for wasp offspring. *Nat Chem Biol* 2010; 6: 261–263
- [97] Kaltenpoth M, Schmitt T, Polidori C, Koedam D, Strohm E. Symbiotic streptomycetes in antennal glands of the South American digger wasp genus *Trachypus* (Hymenoptera, Crabronidae). *Physiol Entomol* 2010; 35: 196–200
- [98] Douglas AE. Lessons from studying insect symbioses. *Cell Host Microbe* 2011; 10: 359–367
- [99] Dedeine F, Vavre F, Fleury F, Loppin B, Hochberg ME, Bouletreau M. Removing symbiotic *Wolbachia* bacteria specifically inhibits oogenesis in a parasitic wasp. *P Natl Acad Sci USA* 2001; 98: 6247–6252
- [100] Kremer N, Charif D, Henri H, Gavory F, Wincker P, Mavingui P, Vavre F. Influence of *Wolbachia* on host gene expression in an obligatory symbiosis. *BMC Microbiol* 2012; 12: S7
- [101] Zchori-Fein E, Gottlieb Y, Kelly SE, Brown JK, Wilson JM, Karr TL, Hunter MS. A newly discovered bacterium associated with parthenogenesis and a change in host selection behavior in parasitoid wasps. *P Natl Acad Sci USA* 2001; 98: 12555–12560
- [102] Zchori-Fein E, Perlman SJ, Kelly SE, Katzir N, Hunter MS. Characterization of a “bacteroidetes” symbiont in *Encarsia* wasps (Hymenoptera: Aphelinidae): proposal of ‘*Candidatus Cardinium hertigii*.’ *Int J Syst Evol Micr* 2004; 54: 961–968
- [103] Chan MS, Godfray HCJ. Host-feeding strategies of parasitoid wasps. *Evol Ecol* 1993; 7: 593–604

- [104] Roossinck MJ. The good viruses: viral mutualistic symbioses. *Nat Rev Microbiol* 2011; 9: 99–108
- [105] Villarreal LP. Virus-host symbiosis mediated by persistence. *Symbiosis* 2007; 44: 1–9
- [106] Yamamura N. Evolution of mutualistic symbiosis: a differential equation model. *Res Popul Ecol* 1996; 38: 211–218
- [107] Whitfield JB. Estimating the age of the polydnavirus/braconid wasp symbiosis. *P Natl Acad Sci USA* 2002; 99: 7508–7513
- [108] Adams AS, Jordan MS, Adams SM, Suen G, Goodwin LA, Davenport KW, Currie CR, Raffa KF. Cellulose-degrading bacteria associated with the invasive woodwasp *Sirex noctilio*. *ISME J* 2011; 5: 1323–1331
- [109] Kukor JJ, Martin MM. Acquisition of digestive enzymes by siricid woodwasps from their fungal symbiont. *Science* 1983; 220: 1161–1163
- [110] Somavilla A, de Oliveira ML, Silveira OT. Diversity and aspects of the ecology of social wasps (Vespidae, Polistinae) in central Amazonian “terra firme” forest. *Rev Bras Entomol* 2014; 58: 349–355
- [111] Adnani N, Rajski SR, Bugni TS. Symbiosis-inspired approaches to antibiotic discovery. *Nat Prod Rep* 2017; 34: 784–814
- [112] Mueller UG, Gerardo N. Fungus-farming insects: multiple origins and diverse evolutionary histories. *P Natl Acad Sci* 2002; 99: 15247–15249
- [113] Aanen DK, Eggleton P, Rouland-Lefevre C, Guldborg-Froslev T, Rosendahl S, Boomsma JJ. The evolution of fungus-growing termites and their mutualistic fungal symbionts. *P Natl Acad Sci USA* 2002; 99: 14887–14892
- [114] Kambhampati S, Eggleton P. Taxonomy and Phylogeny of Termites. In: Abe T, Bignell DE, Higashi M, eds. *Termites: Evolution, Sociality, Symbioses, Ecology*. Dordrecht: Kluwer Academic Publishers; 2000: 1–23
- [115] Lima JT, Costa-Leonardo AM. Recursos alimentares explorados pelos cupins (Insecta: Isoptera). *Biota Neotrop* 2007; bn04007022007
- [116] Grieco MB, Lopes FAC, Oliveira LS, Tschoeke DA, Popov CC, Thompson CC, Goncalves LC, Constantino R, Martins OB, Kruger RH, de Souza W, Thompson FL. Metagenomic analysis of the whole gut microbiota in Brazilian termitidae termites *Cornitermes cumulans*, *Cyrlilotermes stricinus*, *Syntermes dirus*, *Nasutitermes jaraguae*, *Nasutitermes aquilinus*, *Grigiotermes bequaerti*, and *Orthognathotermes mirim*. *Curr Microbiol* 2019; 76: 687–697
- [117] Padilla MA, Rodrigues RAF, Bastos JCS, Martini MC, Barnabe ACD, Kohn LK, Uetanabaro APT, Bomfim GF, Afonso RS, Fantinatti-Garborggini F, Arns CW. Actinobacteria from termite mounds show antiviral activity against bovine viral diarrhoea virus, a surrogate model for hepatitis c virus. *Evid-Based Compl Alt* 2015: 1–9
- [118] Wyche TP, Ruzzini AC, Beemelmans C, Kim KH, Klassen JL, Cao S, Poulsen M, Bugni TS, Currie CR, Clardy J. Linear peptides are the major products of a biosynthetic pathway that encodes for cyclic depsipeptides. *Org Lett* 2017; 19: 1772–1775
- [119] Benndorf R, Guo H, Sommerwerk E, Weigel C, Garcia-Altares M, Martin K, Hu H, Kufner M, De Beer ZW, Poulsen M. Natural products from actinobacteria associated with fungus-growing termites. *Antibiotics* 2018; 7: 83
- [120] Klassen JL, Lee SR, Thomas-Poulsen M, Beemelmans C, Kim KH. Efomycins K and L from a termite-associated *Streptomyces* sp. m56 and their putative biosynthetic origin. *Front Microbiol* 2019; 10: 1739
- [121] Lee SR, Song JH, Song JH, Ko HJ, Baek JY, Trinh TA, Beemelmans C, Yamabe N, Kim KH. Chemical identification of isoflavonoids from a termite-associated *Streptomyces* sp. RB1 and their neuroprotective effects in murine hippocampal HT22 cell line. *Int J Mol Sci* 2018; 19: 2640
- [122] Lee SR, Lee D, Yu JS, Benndorf R, Lee S, Lee DS, Huh J, De Beer ZW, Kim YH, Beemelmans C. Natalenamides A–C, cyclic tripeptides from the termite-associated *Actinomadura* sp. RB99. *Molecules* 2018; 23: 3003
- [123] Beemelmans C, Ramadhar TR, Kim KH, Klassen JL, Cao S, Wyche TP, Hou Y, Poulsen M, Bugni TS, Currie CR. Macrotermycins A–D, glycosylated macrolactams from a termite-associated *Amycolatopsis* sp. M39. *Org Lett* 2017; 19: 1000–1003
- [124] Yoon SY, Lee SR, Hwang JY, Benndorf R, Beemelmans C, Chung SJ, Kim KH. Fridamycin A, a microbial natural product, stimulates glucose uptake without inducing adipogenesis. *Nutrients* 2019; 11: 765
- [125] Carr G, Poulsen M, Klassen JL, Hou Y, Wyche TP, Bugni TS, Currie CR, Clardy J. Microtermolides A and B from termite-associated *Streptomyces* sp. and structural revision of vinylamycin. *Org Lett* 2012; 14: 2822–2825
- [126] Yang Z, Ma M, Yang CH, Gao Y, Zhang Q, Chen Y. Determination of the absolute configurations of microtermolides A and B. *J Nat Prod* 2016; 79: 2408–2412
- [127] Constantino R, Acioli A. Termite diversity in Brazil (Insecta: Isoptera). In: Moreira FMS, Siqueira, JO, editors. *Soil biodiversity in Amazonian and other Brazilian ecosystems*. Wallingford: CAB International; 2006: 117–128
- [128] Hernandez ILC, Godinho MJL, Magalhaes A, Schefer AB, Ferreira AG, Berlinck RGS. N-acetyl-gamma-hydroxyvaline lactone, an unusual amino acid derivative from marine streptomycete. *J Nat Prod* 2000; 63: 664–665
- [129] Chagas FO, Dias LG, Pupo MT. A mixed culture of endophytic fungi increases production of antifungal polyketides. *J Chem Ecol* 2013; 39: 1335–1342
- [130] Sanchez-Bayo F, Wyckhuys KAG. Worldwide decline of the entomofauna: a review of its drivers. *Biol Conserv* 2019; 232: 8–27