

OXYEPOECUS

- bidentatus.** *Oxyepoecus bidentatus* Delsinne & Mackay, in Delsinne, *et al.* 2012: 2, figs. 2, 3 (w.) PARAGUAY.
- browni.** *Oxyepoecus browni* Albuquerque & Brandão, 2004: 60, figs. 1a-c (w.) BRAZIL (Minas Gerais).
Ulysséa & Brandão, 2012: 172 (q.).
Status as species: Albuquerque & Brandão, 2009: 292 (in key); Ulysséa & Brandão, 2012: 172.
- bruchi.** *Oxyepoecus bruchi* Santschi, 1926d: 6, figs. A-D (q.) ARGENTINA (Córdoba).
Santschi, 1929d: 295 (w.).
Combination in *Mitara*: Santschi, 1927d: 246;
combination in *Martia*: Santschi, 1929d: 295;
combination in *Oxyepoecus*: Ettershank, 1966: 146.
Status as species: Santschi, 1927d: 246; Santschi, 1929d: 295; Kusnezov, 1949a: 426; Kusnezov, 1952h: 721; Ettershank, 1966: 146; Kutter, 1968b: 204; Kempf, 1972a: 173; Kempf, 1974b: 478 (redescription); Brandão, 1991: 365; Bolton, 1995b: 301; Albuquerque & Brandão, 2004: 62 (redescription); Wild, 2007b: 34; Albuquerque & Brandão, 2009: 292 (in key); Delsinne, *et al.* 2012: 4.
Senior synonym of minuta: Kempf, 1974b: 478; Brandão, 1991: 365; Bolton, 1995b: 301; Albuquerque & Brandão, 2004: 62.
- crassinodus.** *Oxyepoecus crassinodus* Kempf, 1974b: 482, figs. 2, 8, 15, 21 (w.) BRAZIL (Paraná, Santa Catarina).
Status as species: Brandão, 1991: 364; Bolton, 1995b: 301; Albuquerque & Brandão, 2004: 64 (redescription); Albuquerque & Brandão, 2009: 292 (in key).
- daguerrei.** *Martia daguerrei* Santschi, 1933e: 111 (w.) ARGENTINA (Buenos Aires).
Combination in *Oxyepoecus*: Ettershank, 1966: 146.
Status as species: Kusnezov, 1949a: 426; Kusnezov, 1952h: 721; Ettershank, 1966: 146; Kutter, 1968b: 204; Kempf, 1972a: 173; Kempf, 1974b: 486 (redescription); Bolton, 1995b: 301; Albuquerque & Brandão, 2009: 294 (redescription).
- ephippiatus.** *Oxyepoecus ephippiatus* Albuquerque & Brandão, 2004: 64, figs. 2a-c (w.) BRAZIL (Amazonas).
Status as species: Albuquerque & Brandão, 2009: 292 (in key).
- inquilinus.** *Martia inquilina* Kusnezov, 1952h: 720 (diagnosis in key) (w.) ARGENTINA (Tucumán).
Combination in *Oxyepoecus*: Ettershank, 1966: 146.
Status as species: Kusnezov, 1953b: 337; Ettershank, 1966: 146; Kutter, 1968b: 204; Kempf, 1972a: 173; Kempf, 1974b: 489 (redescription); Brandão, 1991: 365; Bolton, 1995b: 301; Albuquerque & Brandão, 2004: 66 (redescription); Albuquerque & Brandão, 2009: 292 (in key); Delsinne, *et al.* 2012: 4; Fernández & Serna, 2019: 809.
Senior synonym of turgidus: Kempf, 1974b: 489; Brandão, 1991: 365; Bolton, 1995b: 301; Albuquerque & Brandão, 2004: 66.
- kempfi.** *Oxyepoecus kempfi* Albuquerque & Brandão, 2004: 67, figs. 3a-c (w.) BRAZIL (Piauí).
Status as species: Albuquerque & Brandão, 2009: 292 (in key).
- longicephalus.** *Oxyepoecus longicephalus* Albuquerque & Brandão, 2004: 69, figs. 4a-c (w.) BRAZIL (Rio Grande do Sul, São Paulo).
Status as species: Albuquerque & Brandão, 2009: 292 (in key).

luederwaldti. *Monomorium (Martia) rastratum* var. *luederwaldti* Forel, 1913l: 219 (w.) BRAZIL (São Paulo).

Combination in *Martia*: Kusnezov, 1952h: 722;
combination in *Oxyepoecus*: Ettershank, 1966: 146.

Subspecies of *rastratus*: Luederwaldt, 1918: 43; Borgmeier, 1927c: 99; Kusnezov, 1952h: 722; Ettershank, 1966: 146; Kempf, 1972a: 173.

Junior synonym of *rastratus*: Kempf, 1974b: 498; Brandão, 1991: 365; Bolton, 1995b: 301; Albuquerque & Brandão, 2009: 300.

mandibularis. *Monomorium (Martia) mandibulare* Emery, 1913b: 261, fig. 12 (w.) BOLIVIA.

Combination in *Martia*: Kusnezov, 1949a: 426;
combination in *Oxyepoecus*: Ettershank, 1966: 146.

Status as species: Emery, 1922e: 183; Borgmeier, 1927b: 65 (in key); Kusnezov, 1949a: 426; Kusnezov, 1952h: 722; Ettershank, 1966: 146; Kempf, 1972a: 173; Kempf, 1974b: 491; Bolton, 1995b: 301; Albuquerque & Brandão, 2009: 294.

minuta. *Martia minuta* Kusnezov, 1952h: 721 (diagnosis in key) (w.) ARGENTINA (Tucumán).

Combination in *Oxyepoecus*: Ettershank, 1966: 146.

Status as species: Kusnezov, 1953b: 337; Ettershank, 1966: 146; Kutter, 1968b: 204; Kempf, 1972a: 173.

Junior synonym of *bruchi*: Kempf, 1974b: 478; Brandão, 1991: 365; Bolton, 1995b: 301; Albuquerque & Brandão, 2004: 62.

myops. *Oxyepoecus myops* Albuquerque & Brandão, 2009: 295, figs. 1a-c, 6 (w.q.) BRAZIL (São Paulo, Minas Gerais, Paraná, Rio de Janeiro, Santa Catarina).

plaumanni. *Oxyepoecus plaumanni* Kempf, 1974b: 492, figs. 27-30 (w.q.) BRAZIL (Rio Grande do Sul, Santa Catarina, Paraná).

Status as species: Brandão, 1991: 364; Bolton, 1995b: 301; Albuquerque & Brandão, 2009: 297 (redescription).

punctifrons. *Monomorium punctifrons* Borgmeier, 1927b: 63 (w.) BRAZIL (Paraná).

[Also described as new by Borgmeier, 1928a: 39.]

Wheeler, G.C. & Wheeler, J. 1977a: 586 (l.). Albuquerque & Brandão, 2004: 71 (q.).

Combination in *Martia*: Kusnezov, 1949a: 426;
combination in *Oxyepoecus*: Ettershank, 1966: 146.

Status as species: Borgmeier, 1927c: 99; Kusnezov, 1949a: 426; Kusnezov, 1952h: 722; Ettershank, 1966: 146; Kempf, 1972a: 173; Kempf, 1974b: 495 (redescription); Brandão, 1991: 365; Bolton, 1995b: 301; Albuquerque & Brandão, 2004: 71 (redescription); Albuquerque & Brandão, 2009: 292 (in key).

quadratus. *Oxyepoecus quadratus* Albuquerque & Brandão, 2004: 73, figs. 6a-c (w.) ECUADOR, PERU.

Status as species: Albuquerque & Brandão, 2009: 292 (in key); Bezděčková, *et al.* 2015: 118.

rastratus. *Monomorium rastratum* Mayr, 1887: 615 (w.) BRAZIL (Santa Catarina).

Kempf, 1974b: 500 (q.m.); Wheeler, G.C. & Wheeler, J. 1980: 534 (l.).

Combination in *Monomorium (Martia)*: Forel, 1907a: 21; Forel, 1912g: 3;
combination in *Martia*: Kusnezov, 1949a: 426;
combination in *Oxyepoecus*: Ettershank, 1966: 146.

Status as species: Dalla Torre, 1893: 69; Forel, 1895b: 125; Forel, 1907a: 21 (in text); Forel, 1912g: 3; Emery, 1922e: 183; Borgmeier, 1927b: 65 (in key); Borgmeier, 1927c:

- 99; Kusnezov, 1949a: 426; Kusnezov, 1952h: 722; Ettershank, 1966: 146; Kempf, 1972a: 173; Kempf, 1974b: 498 (redescription); Brandão, 1991: 365; Bolton, 1995b: 302; Wild, 2007b: 34; Albuquerque & Brandão, 2009: 299 (redescription); Delsinne, *et al.* 2012: 4.
Senior synonym of *luederwaldti*: Kempf, 1974b: 498; Brandão, 1991: 365; Bolton, 1995b: 302; Albuquerque & Brandão, 2009: 300.
- regularis*.** *Oxyepoecus regularis* Ulysséa & Brandão, 2012: 169, figs. 1, 2 (w.q.) BRAZIL (Bahia).
- reticulatus*.** *Oxyepoecus reticulatus* Kempf, 1974b: 502, figs. 23-26 (w.q.) BRAZIL (Santa Catarina, Paraná, São Paulo).
Status as species: Brandão, 1991: 364; Bolton, 1995b: 302; Wild, 2007b: 34; Albuquerque & Brandão, 2009: 303 (redescription); Delsinne, *et al.* 2012: 5.
- rosai*.** *Oxyepoecus rosai* Albuquerque & Brandão, 2009: 304, figs. 5a-c, 8 (w.q.) BRAZIL (São Paulo, Paraná, Rio de Janeiro, Santa Catarina).
- striatus*.** *Oxyepoecus striatus* Mackay & Delsinne, in Delsinne, *et al.* 2012: 5, fig. 4 (w.) PARAGUAY.
- turgidus*.** *Oxyepoecus turgidus* Kempf, 1969: 278, figs. 1-3 (w.) BRAZIL (São Paulo, Goiás).
Status as species: Kempf, 1972a: 173.
Junior synonym of *inquilinus*: Kempf, 1974b: 489; Brandão, 1991: 365; Bolton, 1995b: 302; Albuquerque & Brandão, 2004: 66.
- vezenyii*.** *Monomorium (Martia) vezenyii* Forel, 1907a: 20 (w.) PARAGUAY.
Kempf, 1969: 277 (q.).
Combination in *Martia*: Kusnezov, 1949a: 426;
combination in *Oxyepoecus*: Ettershank, 1966: 145.
Status as species: Emery, 1922e: 183; Borgmeier, 1927b: 65 (in key); Kusnezov, 1949a: 426; Kusnezov, 1952h: 722; Ettershank, 1966: 146; Kempf, 1969: 277; Kempf, 1972a: 173; Kempf, 1974b: 505 (redescription); Bolton, 1995b: 302; Albuquerque & Brandão, 2004: 75 (redescription); Wild, 2007b: 34; Albuquerque & Brandão, 2009: 292 (in key); Delsinne, *et al.* 2012: 6.
- vivax*.** *Oxyepoecus vivax* Kempf, 1974b: 508, figs. 31-33, 38, 39 (q.) BRAZIL (Minas Gerais).
Status as species: Brandão, 1991: 364; Bolton, 1995b: 302; Albuquerque & Brandão, 2004: 77 (redescription).